



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 16, 2024 – 10:19 PM EDT

PDB ID : 3O10  
Title : Crystal structure of the HEPN domain from human saccin  
Authors : Kozlov, G.; Gehring, K.  
Deposited on : 2010-07-20  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 1.20.1  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	141	 3% 75% 21% ..
1	B	141	 4% 87% 9% ..
1	C	141	 22% 86% 9% ..
1	D	141	 4% 83% 14% .

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4829 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sacsin.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	139	1125	709	201	208	4	3	0	3	0
1	B	139	1102	694	195	206	4	3	0	0	0
1	C	136	1085	684	192	202	4	3	0	0	0
1	D	137	1089	687	192	203	4	3	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	4439	GLY	-	EXPRESSION TAG	UNP Q9NZJ4
B	4439	GLY	-	EXPRESSION TAG	UNP Q9NZJ4
C	4439	GLY	-	EXPRESSION TAG	UNP Q9NZJ4
D	4439	GLY	-	EXPRESSION TAG	UNP Q9NZJ4

- Molecule 2 is MALONATE ION (three-letter code: MLI) (formula: C<sub>3</sub>H<sub>2</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 7 3 4	0	0
2	B	1	Total C O 7 3 4	0	0
2	D	1	Total C O 7 3 4	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	141	Total O 141 141	0	0
3	B	146	Total O 146 146	0	0
3	C	51	Total O 51 51	0	0
3	D	69	Total O 69 69	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Sacsin




- Molecule 1: Sacsin



- Molecule 1: Sacsin



- Molecule 1: Sacsin



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	72.58Å 72.58Å 201.18Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	45.86 – 1.90 45.86 – 1.90	Depositor EDS
% Data completeness (in resolution range)	98.1 (45.86-1.90) 98.1 (45.86-1.90)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.41 (at 1.91Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.184 , 0.225 0.183 , 0.225	Depositor DCC
$R_{free}$ test set	2441 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.0	Xtrriage
Anisotropy	0.099	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 56.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.025 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4829	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.61% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MLI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.74	0/1152	0.82	1/1553 (0.1%)
1	B	0.82	1/1120 (0.1%)	0.78	1/1511 (0.1%)
1	C	0.56	0/1103	0.65	0/1488
1	D	0.62	0/1107	0.65	0/1494
All	All	0.70	1/4482 (0.0%)	0.73	2/6046 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	4561	MSE	SE-CE	-5.66	1.62	1.95

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	4559	ARG	NE-CZ-NH2	-6.69	116.96	120.30
1	A	4559	ARG	NE-CZ-NH2	-5.79	117.40	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1125	0	1132	23	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1102	0	1094	16	0
1	C	1085	0	1077	9	0
1	D	1089	0	1083	18	0
2	A	7	0	2	0	0
2	B	7	0	2	0	0
2	D	7	0	2	0	0
3	A	141	0	0	6	0
3	B	146	0	0	4	0
3	C	51	0	0	0	0
3	D	69	0	0	5	0
All	All	4829	0	4392	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:4559:ARG:NH1	1:C:4562:GLU:OE1	1.89	1.06
1:B:4555:GLU:OE1	3:B:92:HOH:O	1.81	0.98
1:A:4555:GLU:OE1	3:A:91:HOH:O	1.91	0.88
1:A:4501[B]:LYS:HD2	1:A:4502:PRO:HD2	1.59	0.85
3:A:96:HOH:O	1:D:4521:ASN:HB3	1.77	0.84
1:A:4559:ARG:NH2	3:A:49:HOH:O	2.12	0.83
1:C:4479:LEU:HD21	1:D:4537:ARG:HD2	1.64	0.80
1:B:4551:ARG:NH1	3:B:137:HOH:O	2.14	0.76
1:B:4439:GLY:HA3	1:B:4493:ARG:HB3	1.68	0.74
1:D:4517:GLU:OE2	3:D:66:HOH:O	2.06	0.73
1:A:4534:LEU:HB3	1:A:4541:LEU:HD23	1.71	0.72
1:B:4465:LEU:HD11	1:B:4561:MSE:CE	2.21	0.71
1:D:4563:CYS:HB2	3:D:26:HOH:O	1.91	0.70
1:A:4577:GLN:C	3:A:138:HOH:O	2.32	0.68
1:A:4501[B]:LYS:HD2	1:A:4502:PRO:CD	2.24	0.66
1:C:4501:LYS:HD2	1:C:4502:PRO:HD2	1.77	0.66
1:A:4501[B]:LYS:CD	1:A:4502:PRO:HD2	2.26	0.65
1:D:4563:CYS:CB	3:D:26:HOH:O	2.46	0.64
1:C:4444:PRO:O	1:C:4448:ARG:HG2	2.00	0.61
1:A:4497:ASP:N	1:A:4497:ASP:OD1	2.34	0.61
1:A:4455:ARG:NH1	1:A:4572:GLU:OE2	2.33	0.61
1:D:4493:ARG:HD3	1:D:4512:TYR:OH	2.03	0.58
1:A:4442:GLY:O	1:D:4521:ASN:ND2	2.36	0.57

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:4535:LYS:HD3	1:B:4551:ARG:HH21	1.71	0.56
3:A:72:HOH:O	1:D:4524:HIS:HE1	1.87	0.55
1:B:4468:ASN:ND2	3:B:145:HOH:O	2.39	0.55
1:B:4551:ARG:CG	1:B:4551:ARG:HH11	2.21	0.54
1:B:4465:LEU:HD21	1:B:4561:MSE:HE2	1.90	0.52
1:A:4503:THR:O	1:A:4507:GLN:HG3	2.09	0.52
1:B:4465:LEU:HD11	1:B:4561:MSE:HE1	1.91	0.52
1:A:4465:LEU:HD21	1:A:4561:MSE:SE	2.62	0.50
1:D:4570:LYS:HD2	3:D:69:HOH:O	2.13	0.48
1:D:4507:GLN:O	1:D:4511:GLU:HG3	2.14	0.47
1:B:4551:ARG:HD3	1:B:4552:PHE:CZ	2.50	0.47
1:D:4493:ARG:CD	1:D:4512:TYR:OH	2.63	0.47
1:A:4439:GLY:HA3	1:A:4493:ARG:HG2	1.95	0.46
1:B:4517:GLU:HG2	3:B:74:HOH:O	2.14	0.46
1:C:4479:LEU:HD21	1:D:4537:ARG:CD	2.40	0.46
1:A:4485:LEU:HB3	1:A:4505:LEU:HD23	1.97	0.46
1:C:4465:LEU:HD13	1:C:4466:HIS:NE2	2.30	0.46
1:C:4534:LEU:HB3	1:C:4541:LEU:HD23	1.98	0.46
1:A:4508:LYS:HE3	3:A:93:HOH:O	2.15	0.45
1:A:4568:ILE:O	1:A:4572:GLU:HG3	2.15	0.45
1:C:4501:LYS:CD	1:C:4502:PRO:HD2	2.45	0.45
1:B:4574:PHE:O	1:B:4577:GLN:HG3	2.16	0.45
1:D:4489:ASP:OD1	1:D:4493:ARG:HD3	2.17	0.45
1:A:4507:GLN:HG2	1:A:4520:THR:HG21	1.99	0.44
1:D:4493:ARG:HD3	1:D:4512:TYR:HH	1.82	0.44
1:A:4492:VAL:HG21	1:A:4512:TYR:CD2	2.52	0.44
1:B:4551:ARG:HD3	1:B:4552:PHE:CE2	2.53	0.43
1:A:4498:LYS:HE2	1:B:4540:ASP:OD2	2.20	0.42
1:D:4528:ALA:C	1:D:4530:GLY:N	2.72	0.42
1:B:4489:ASP:OD1	1:B:4493:ARG:HD3	2.19	0.42
1:A:4496:SER:C	1:A:4497:ASP:OD1	2.59	0.41
1:C:4451:LEU:HD21	1:C:4572:GLU:HG3	2.02	0.41
1:A:4576:GLN:O	1:A:4577:GLN:C	2.58	0.41
1:A:4509:ILE:HG22	1:A:4516:LEU:HD13	2.03	0.41
1:B:4524:HIS:NE2	1:D:4544:PHE:HB3	2.36	0.40
1:A:4501[B]:LYS:HZ3	1:A:4503:THR:H	1.68	0.40
1:D:4563:CYS:SG	3:D:14:HOH:O	2.62	0.40
1:D:4538:TYR:HA	1:D:4539:PRO:HD3	1.98	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	140/141 (99%)	139 (99%)	1 (1%)	0	100	100
1	B	137/141 (97%)	134 (98%)	3 (2%)	0	100	100
1	C	134/141 (95%)	132 (98%)	2 (2%)	0	100	100
1	D	135/141 (96%)	132 (98%)	2 (2%)	1 (1%)	22	12
All	All	546/564 (97%)	537 (98%)	8 (2%)	1 (0%)	47	38

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	4533	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	122/118 (103%)	115 (94%)	7 (6%)	20	11
1	B	118/118 (100%)	116 (98%)	2 (2%)	60	57
1	C	116/118 (98%)	111 (96%)	5 (4%)	29	19
1	D	117/118 (99%)	114 (97%)	3 (3%)	46	39
All	All	473/472 (100%)	456 (96%)	17 (4%)	35	26

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4440	SER
1	A	4446	GLU
1	A	4497	ASP
1	A	4499	ASP
1	A	4541	LEU
1	A	4545	PRO
1	A	4577	GLN
1	B	4551	ARG
1	B	4559	ARG
1	C	4465	LEU
1	C	4497	ASP
1	C	4499	ASP
1	C	4541	LEU
1	C	4559	ARG
1	D	4558	MSE
1	D	4559	ARG
1	D	4561	MSE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	4470	ASN
1	A	4524	HIS
1	B	4546	GLN
1	C	4470	ASN
1	C	4573	ASN
1	D	4524	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	MLI	A	1	-	6,6,6	1.52	0	7,7,7	0.93	0
2	MLI	B	1	-	6,6,6	1.21	0	7,7,7	1.07	0
2	MLI	D	1	-	6,6,6	1.14	0	7,7,7	1.20	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MLI	A	1	-	-	0/4/4/4	-
2	MLI	B	1	-	-	0/4/4/4	-
2	MLI	D	1	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	136/141 (96%)	0.01	4 (2%) 51 54	9, 17, 37, 46	0
1	B	136/141 (96%)	-0.05	5 (3%) 41 44	9, 18, 34, 46	0
1	C	133/141 (94%)	0.96	31 (23%) 0 0	16, 37, 77, 105	0
1	D	134/141 (95%)	0.21	5 (3%) 41 44	14, 31, 52, 67	0
All	All	539/564 (95%)	0.28	45 (8%) 11 13	9, 25, 56, 105	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	4497	ASP	5.5
1	C	4509	ILE	5.5
1	C	4512	TYR	5.5
1	C	4508	LYS	5.2
1	C	4514	GLN	4.9
1	C	4505	LEU	4.5
1	B	4440	SER	4.4
1	C	4523	VAL	4.0
1	B	4439	GLY	3.9
1	C	4503	THR	3.7
1	C	4501	LYS	3.6
1	C	4522	ASP	3.5
1	D	4531	VAL	3.5
1	C	4500	VAL	3.4
1	C	4502	PRO	3.4
1	B	4441	VAL	3.2
1	C	4511	GLU	3.2
1	C	4524	HIS	3.1
1	C	4516	LEU	3.1
1	C	4507	GLN	3.0
1	C	4504	ALA	3.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	4517	GLU	3.0
1	C	4566	CYS	2.9
1	A	4440	SER	2.8
1	D	4512	TYR	2.8
1	C	4526	LEU	2.8
1	C	4495	LYS	2.7
1	C	4576	GLN	2.7
1	C	4499	ASP	2.6
1	D	4544	PHE	2.5
1	C	4520	THR	2.4
1	B	4577	GLN	2.4
1	A	4497	ASP	2.4
1	A	4439	GLY	2.3
1	D	4535	LYS	2.3
1	C	4529	TYR	2.3
1	C	4521	ASN	2.2
1	C	4563	CYS	2.2
1	C	4506	ALA	2.2
1	A	4577	GLN	2.2
1	B	4512	TYR	2.2
1	D	4517	GLU	2.1
1	C	4496	SER	2.0
1	C	4528	ALA	2.0
1	C	4510	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MLI	D	1	7/7	0.86	0.15	34,35,36,37	0
2	MLI	A	1	7/7	0.93	0.15	23,24,25,26	0
2	MLI	B	1	7/7	0.97	0.07	22,23,24,25	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.