

# Full wwPDB X-ray Structure Validation Report (i)

#### Oct 3, 2023 – 02:33 AM EDT

PDB ID : 6O47

Title: human cGAS core domain (K427E/K428E) bound with RU-521 Authors: Xie, W.; Lama, L.; Adura, C.; Glickman, J.F.; Tuschl, T.; Patel, D.J.

Deposited on : 2019-02-28

Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : FAILED

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : FAILED

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.35.1

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



## 2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 3117 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


• Molecule 1 is a protein called Cyclic GMP-AMP synthase.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	Λ	346	Total	С	N	О	S	0	0	0
1	A	340	2854	1824	493	522	15	0	0	

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual Comment		Reference
A	151	SER	-	expression tag	UNP Q8N884
A	427	GLU	LYS	engineered mutation	UNP Q8N884
A	428	GLU	LYS	engineered mutation	UNP Q8N884

• Molecule 2 is CITRIC ACID (three-letter code: CIT) (formula: C<sub>6</sub>H<sub>8</sub>O<sub>7</sub>).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 13 6 7	0	0
2	A	1	Total C O 13 6 7	0	0



• Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Zn 1 1	0	0


• Molecule 4 is  $(3 \{S\})$ -3- $[1-[4,5-bis(chloranyl)-1 \{H\}-benzimidazol-2-yl]$ -3-methyl-5-oxidanyl-pyrazol-4-yl]-3  $\{H\}$ -2-benzofuran-1-one (three-letter code: LLS) (formula:  $C_{19}H_{12}Cl_2N_4O_3$ ).



Mol	Chain	Residues	${f Atoms}$			ZeroOcc	AltConf		
4	A	1	Total 28	C 19	Cl 2	N 4	O 3	0	1

• Molecule 5 is 2-(4,5-dichloro-1H-benzimidazol-2-yl)-5-methyl-4-[(1R)-3-oxo-1,3-dihyd ro-2-benzofuran-1-yl]-1,2-dihydro-3H-pyrazol-3-one (three-letter code: AEV) (formula:  $C_{19}H_{12}Cl_2N_4O_3$ ).





Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf
5	Λ	1	Total	С	Cl	N	О	0	1
)	A	1	28	19	2	4	3	U	1

#### • Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	180	Total O 180 180	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



# 3 Data and refinement statistics (i)

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 64	Depositor
Cell constants	116.16Å 116.16Å 59.95Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $120.00^{\circ}$	Depositor
Resolution (Å)	29.27 - 2.20	Depositor
% Data completeness	88.3 (29.27-2.20)	Depositor
(in resolution range)	,	-
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.93  (at  2.20Å)	Xtriage
Refinement program	PHENIX (1.14_3260: ???)	Depositor
$R, R_{free}$	0.181 , $0.221$	Depositor
Wilson B-factor $(\mathring{A}^2)$	27.7	Xtriage
Anisotropy	0.051	Xtriage
L-test for twinning <sup>2</sup>	$< L > = 0.48, < L^2> = 0.31$	Xtriage
Estimated twinning fraction	0.053 for h,-h-k,-l	Xtriage
Total number of atoms	3117	wwPDB-VP
Average B, all atoms $(\mathring{A}^2)$	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.68% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $< L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

## 4 Model quality (i)

### 4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

### 4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3 Torsion angles (i)

#### 4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

## 4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

## 4.6 Ligand geometry (i)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trus	Chain	Res	Link	Во	ond leng	ths	Bond angles		
IVIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z  > 2
2	CIT	A	601	-	12,12,12	1.15	0	17,17,17	1.50	1 (5%)
2	CIT	A	602	-	12,12,12	1.18	0	17,17,17	1.46	1 (5%)
5	AEV	A	605[B]	-	28,32,32	2.59	8 (28%)	28,49,49	3.55	7 (25%)
4	LLS	A	604[A]	-	27,32,32	1.80	5 (18%)	27,49,49	3.71	7 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CIT	A	601	-	-	7/16/16/16	-
2	CIT	A	602	-	-	6/16/16/16	-
5	AEV	A	605[B]	-	-	0/3/20/20	0/5/5/5
4	LLS	A	604[A]	-	-	0/0/28/28	0/5/5/5

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	Ideal(A)
5	A	605[B]	AEV	OAC-CAS	7.98	1.39	1.23
5	A	605[B]	AEV	CAS-NBB	-5.56	1.30	1.39
4	A	604[A]	LLS	O12-C11	4.87	1.42	1.36
5	A	605[B]	AEV	OAO-CAP	4.76	1.42	1.36
4	A	604[A]	LLS	O12-C04	-4.22	1.42	1.46
5	A	605[B]	AEV	CAS-CAX	-3.64	1.37	1.46
5	A	605[B]	AEV	CAU-CAW	-3.07	1.34	1.39
4	A	604[A]	LLS	C10-C05	-2.77	1.35	1.39
5	A	605[B]	AEV	CAV-NBB	2.71	1.45	1.40
4	A	604[A]	LLS	O01-C02	2.41	1.39	1.32
5	A	605[B]	AEV	CAW-CBA	2.08	1.55	1.51
5	A	605[B]	AEV	CAU-CAP	2.03	1.52	1.47
4	A	604[A]	LLS	C20-C27	-2.02	1.35	1.42

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
5	A	605[B]	AEV	CBA-OAO-CAP	-14.15	104.42	110.78

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
4	A	604[A]	LLS	C04-O12-C11	-11.50	102.70	110.75
4	A	604[A]	LLS	O12-C11-O13	9.51	129.30	121.19
4	A	604[A]	LLS	O12-C11-C10	-7.46	103.96	108.42
5	A	605[B]	AEV	OAO-CAP-OAB	6.95	127.12	121.19
4	A	604[A]	LLS	C05-C10-C11	-6.11	105.72	108.45
5	A	605[B]	AEV	CAA-CAQ-NAM	5.43	126.16	119.26
4	A	604[A]	LLS	O12-C04-C05	-4.79	100.92	103.49
5	A	605[B]	AEV	CAA-CAQ-CAX	-4.49	126.39	131.66
2	A	601	CIT	O6-C6-C3	4.28	120.48	113.05
5	A	605[B]	AEV	CAW-CAU-CAP	-4.13	106.60	108.45
2	A	602	CIT	O6-C6-C3	3.74	119.55	113.05
5	A	605[B]	AEV	OAO-CAP-CAU	-3.16	106.53	108.42
4	A	604[A]	LLS	C25-C23-CL24	-2.41	118.17	120.54
5	A	605[B]	AEV	CAT-CAR-CLD	-2.33	118.25	120.54
4	A	604[A]	LLS	C06-C05-C04	2.15	133.33	128.82

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	CIT	C2-C3-C6-O5
2	A	601	CIT	C2-C3-C6-O6
2	A	601	CIT	O7-C3-C6-O5
2	A	601	CIT	O7-C3-C6-O6
2	A	602	CIT	O7-C3-C6-O5
2	A	602	CIT	O7-C3-C6-O6
2	A	602	CIT	C4-C3-C6-O5
2	A	602	CIT	C4-C3-C6-O6
2	A	601	CIT	C1-C2-C3-O7
2	A	601	CIT	C1-C2-C3-C4
2	A	602	CIT	C3-C4-C5-O4
2	A	602	CIT	C3-C4-C5-O3
2	A	601	CIT	C1-C2-C3-C6


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.



Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 4.7 Other polymers (i)

There are no such residues in this entry.



# 4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 5 Fit of model and data (i)

#### 5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

### 5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

### 5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

### 5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

## 5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

