



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 9, 2018 – 06:49 PM EDT

PDB ID : 1O5U
Title : Crystal structure of a duf861 family protein (tm1112) from thermotoga maritima at 1.83 Å resolution
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2003-10-06
Resolution : 1.83 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : 1.13
EDS : rb-20031172
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20031172


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.83 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	3313 (1.86-1.82)
Clashscore	122126	3530 (1.86-1.82)
Ramachandran outliers	120053	3495 (1.86-1.82)
Sidechain outliers	120020	3496 (1.86-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	101	 76% 11% 13%
1	B	101	 73% 14% 13%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	UNL	A	401	-	-	X	-
2	UNL	B	402	-	-	X	-
2	UNL	B	403	-	-	X	-

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 1802 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called novel *Thermotoga maritima* enzyme TM1112.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	88	755	495	118	140	2	0	1	0
1	B	88	752	492	118	140	2	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	MET	-	LEADER SEQUENCE	UNP Q9X0J6
A	-10	GLY	-	LEADER SEQUENCE	UNP Q9X0J6
A	-9	SER	-	LEADER SEQUENCE	UNP Q9X0J6
A	-8	ASP	-	LEADER SEQUENCE	UNP Q9X0J6
A	-7	LYS	-	LEADER SEQUENCE	UNP Q9X0J6
A	-6	ILE	-	LEADER SEQUENCE	UNP Q9X0J6
A	-5	HIS	-	LEADER SEQUENCE	UNP Q9X0J6
A	-4	HIS	-	LEADER SEQUENCE	UNP Q9X0J6
A	-3	HIS	-	LEADER SEQUENCE	UNP Q9X0J6
A	-2	HIS	-	LEADER SEQUENCE	UNP Q9X0J6
A	-1	HIS	-	LEADER SEQUENCE	UNP Q9X0J6
A	0	HIS	-	LEADER SEQUENCE	UNP Q9X0J6
B	-11	MET	-	LEADER SEQUENCE	UNP Q9X0J6
B	-10	GLY	-	LEADER SEQUENCE	UNP Q9X0J6
B	-9	SER	-	LEADER SEQUENCE	UNP Q9X0J6
B	-8	ASP	-	LEADER SEQUENCE	UNP Q9X0J6
B	-7	LYS	-	LEADER SEQUENCE	UNP Q9X0J6
B	-6	ILE	-	LEADER SEQUENCE	UNP Q9X0J6
B	-5	HIS	-	LEADER SEQUENCE	UNP Q9X0J6
B	-4	HIS	-	LEADER SEQUENCE	UNP Q9X0J6
B	-3	HIS	-	LEADER SEQUENCE	UNP Q9X0J6
B	-2	HIS	-	LEADER SEQUENCE	UNP Q9X0J6
B	-1	HIS	-	LEADER SEQUENCE	UNP Q9X0J6
B	0	HIS	-	LEADER SEQUENCE	UNP Q9X0J6

- Molecule 2 is UNKNOWN LIGAND (three-letter code: UNL) (formula:).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	2	Total C 8 8	0	0
2	A	1	Total C 5 5	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	139	Total O 139 139	0	0
3	B	143	Total O 143 143	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: novel *Thermotoga maritima* enzyme TM1112

Chain A:  76% 11% 13%



- Molecule 1: novel *Thermotoga maritima* enzyme TM1112

Chain B:  73% 14% 13%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	39.63Å 74.73Å 42.21Å 90.00° 91.02° 90.00°	Depositor
Resolution (Å)	27.98 – 1.83 27.98 – 1.83	Depositor EDS
% Data completeness (in resolution range)	98.2 (27.98-1.83) 98.2 (27.98-1.83)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.62 (at 1.83Å)	Xtrriage
Refinement program	REFMAC 5.1.9999	Depositor
R, R_{free}	0.170 , 0.222 0.184 , 0.174	Depositor DCC
R_{free} test set	1089 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å ²)	27.0	Xtrriage
Anisotropy	0.132	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 45.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.043 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	1802	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: UNL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.87	1/780 (0.1%)	0.87	1/1054 (0.1%)
1	B	0.81	0/773	0.90	3/1044 (0.3%)
All	All	0.84	1/1553 (0.1%)	0.88	4/2098 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	61	GLU	CD-OE1	5.51	1.31	1.25

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	83	ARG	NE-CZ-NH1	6.28	123.44	120.30
1	A	54	ASP	CB-CG-OD1	6.04	123.74	118.30
1	B	64	ASP	CB-CG-OD2	5.77	123.49	118.30
1	B	36	ASP	CB-CG-OD2	5.18	122.97	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	2	GLU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	755	0	753	7	0
1	B	752	0	748	8	0
2	A	5	0	0	7	0
2	B	8	0	0	9	0
3	A	139	0	0	2	0
3	B	143	0	0	2	0
All	All	1802	0	1501	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:402:UNL:C4	2:B:402:UNL:C2	1.74	1.61
2:A:401:UNL:C4	2:A:401:UNL:C2	1.77	1.55
2:A:401:UNL:C2	2:A:401:UNL:C5	1.88	1.48
2:A:401:UNL:C3	2:A:401:UNL:C2	1.96	1.43
2:B:403:UNL:C1	2:B:403:UNL:C2	2.20	1.18
2:A:401:UNL:C1	2:A:401:UNL:C5	2.41	0.98
2:B:402:UNL:C1	2:B:402:UNL:C2	2.45	0.94
2:B:402:UNL:C1	2:B:402:UNL:C5	2.53	0.87
1:A:2:GLU:N	3:A:406:HOH:O	2.20	0.75
2:B:402:UNL:C1	2:B:402:UNL:C3	2.67	0.72
1:A:27:GLU:OE1	3:A:415:HOH:O	2.13	0.66
1:B:23:ILE:H	2:B:403:UNL:C3	2.09	0.66
1:B:16:LEU:O	3:B:543:HOH:O	2.15	0.63
2:A:401:UNL:C4	2:A:401:UNL:C3	2.76	0.63
1:A:48:VAL:CG2	1:A:82[A]:VAL:HG11	2.30	0.61
2:B:402:UNL:C2	2:B:402:UNL:C3	2.82	0.57
1:A:84:LYS:HZ2	2:A:401:UNL:C5	2.18	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:84:LYS:HZ2	2:B:402:UNL:C5	2.17	0.54
1:B:34:TYR:HD1	1:B:73:ARG:HG2	1.73	0.54
1:B:61:GLU:OE2	3:B:460:HOH:O	2.18	0.52
1:A:57:LYS:HZ2	2:B:403:UNL:C1	2.29	0.46
1:B:41:CYS:SG	1:B:84:LYS:NZ	2.74	0.46
2:A:401:UNL:C3	2:A:401:UNL:C1	2.89	0.45
1:A:30:GLU:HG2	1:A:75:ARG:NE	2.35	0.42
1:A:78:VAL:HG13	1:A:82[B]:VAL:HG23	2.01	0.42
1:B:34:TYR:CD1	1:B:73:ARG:HG2	2.54	0.42
1:B:11:GLU:O	1:B:15:GLU:HG3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	87/101 (86%)	86 (99%)	1 (1%)	0	100	100
1	B	86/101 (85%)	85 (99%)	1 (1%)	0	100	100
All	All	173/202 (86%)	171 (99%)	2 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	85/96 (88%)	85 (100%)	0	100	100
1	B	84/96 (88%)	83 (99%)	1 (1%)	74	64
All	All	169/192 (88%)	168 (99%)	1 (1%)	87	84

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	77	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are unknown - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.