



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2018 – 01:20 pm GMT

PDB ID : 3O77
Title : The structure of Ca²⁺ Sensor (Case-16)
Authors : Leder, L.; Stark, W.; Freuler, F.; Marsh, M.; Meyerhofer, M.; Stettler, T.;
Mayr, L.M.; Britanova, O.V.; Strukova, L.A.; Chudakov, D.M.
Deposited on : 2010-07-30
Resolution : 2.35 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtrriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	1015 (2.36-2.36)
Clashscore	122126	1081 (2.36-2.36)
Ramachandran outliers	120053	1066 (2.36-2.36)
Sidechain outliers	120020	1067 (2.36-2.36)
RSRZ outliers	108989	1002 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	415	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3214 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Myosin light chain kinase, smooth muscle, Green fluorescent protein, Green fluorescent protein, Calmodulin-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	389	3056	1916	515	610	15	0	0	0

There are 34 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP P11799
A	2	PRO	-	expression tag	UNP P11799
A	3	GLY	-	expression tag	UNP P11799
A	4	SER	-	expression tag	UNP P11799
A	5	SER	ALA	engineered mutation	UNP P11799
A	10	ASN	GLN	engineered mutation	UNP P11799
A	25	LEU	SER	engineered mutation	UNP P42212
A	26	GLU	HIS	engineered mutation	UNP P42212
A	41	ALA	VAL	engineered mutation	UNP P42212
A	49	VAL	ILE	engineered mutation	UNP P42212
A	81	PHE	THR	engineered mutation	UNP P42212
A	84	VAL	ALA	engineered mutation	UNP P42212
A	109	LEU	HIS	engineered mutation	UNP P42212
A	116	ASN	LYS	engineered mutation	UNP P42212
A	117	VAL	-	linker	UNP P42212
A	118	ASP	-	linker	UNP P42212
A	119	GLY	-	linker	UNP P42212
A	120	GLY	-	linker	UNP P42212
A	121	SER	-	linker	UNP P42212
A	122	GLY	-	linker	UNP P42212
A	123	GLY	-	linker	UNP P42212
A	124	THR	-	linker	UNP P42212
A	125	GLY	-	linker	UNP P42212
A	170	LEU	PHE	engineered mutation	UNP P42212
A	188	LEU	PHE	engineered mutation	UNP P42212
A	190	CR2	SER	chromophore	UNP P42212

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	190	CR2	TYR	chromophore	UNP P42212
A	190	CR2	GLY	chromophore	UNP P42212
A	192	LEU	VAL	engineered mutation	UNP P42212
A	193	LYS	GLN	engineered mutation	UNP P42212
A	196	ALA	SER	see sequence details	UNP P42212
A	253	GLY	ASP	engineered mutation	UNP P42212
A	269	SER	TYR	engineered mutation	UNP P42212
A	270	ARG	ASN	engineered mutation	UNP P42212

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Ca 2 2	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	145	Total	O	0	0
			145	145		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Myosin light chain kinase, smooth muscle, Green fluorescent protein, Green fluorescent protein, Calmodulin-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	92.50Å 106.94Å 43.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	70.01 – 2.35 69.96 – 2.35	Depositor EDS
% Data completeness (in resolution range)	99.9 (70.01-2.35) 99.9 (69.96-2.35)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.99 (at 2.34Å)	Xtrriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.198 , 0.274 0.212 , 0.275	Depositor DCC
R_{free} test set	928 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	41.6	Xtrriage
Anisotropy	0.391	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 59.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3214	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.26% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CR2, CA, SO4, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.88	0/3089	0.88	3/4161 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	381	LEU	CA-CB-CG	7.61	132.80	115.30
1	A	160	ASP	CB-CG-OD1	6.39	124.06	118.30
1	A	375	ARG	NE-CZ-NH1	5.28	122.94	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	67	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3056	0	2922	67	0
2	A	2	0	0	0	0
3	A	10	0	0	0	0
4	A	1	0	0	0	0
5	A	145	0	0	7	0
All	All	3214	0	2922	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:274:THR:HG22	1:A:277:GLN:H	1.19	1.06
1:A:58:ASP:HB2	5:A:506:HOH:O	1.56	1.05
1:A:67:GLY:HA3	1:A:68:ASP:HB2	1.37	1.02
1:A:30:ILE:HD11	1:A:79:LEU:HD12	1.44	1.00
1:A:45:ILE:HD13	1:A:59:HIS:CE1	2.06	0.90
1:A:297:THR:HG22	1:A:299:LYS:H	1.42	0.84
1:A:344:LYS:O	1:A:345:MET:HB3	1.79	0.79
1:A:192:LEU:HD21	1:A:245:ASN:HB2	1.67	0.76
1:A:348:THR:O	1:A:349:ASP:HB2	1.87	0.73
1:A:30:ILE:CD1	1:A:79:LEU:HD12	2.20	0.72
1:A:43:PHE:HE2	1:A:45:ILE:HD11	1.56	0.70
1:A:72:LEU:HD13	1:A:73:LEU:H	1.57	0.70
1:A:297:THR:HG22	1:A:299:LYS:N	2.05	0.70
1:A:6:ARG:NH1	5:A:535:HOH:O	2.26	0.68
1:A:43:PHE:CE2	1:A:45:ILE:HD11	2.29	0.68
1:A:67:GLY:CA	1:A:68:ASP:HB2	2.21	0.65
1:A:62:GLN:NE2	5:A:481:HOH:O	2.29	0.64
1:A:295:THR:HG23	1:A:331:THR:HB	1.80	0.63
1:A:101:PHE:CE1	1:A:165:LYS:HG3	2.35	0.62
1:A:391:ASP:HB3	1:A:395:ARG:NH2	2.14	0.62
1:A:313:THR:HG22	1:A:316:GLU:H	1.64	0.62
1:A:321:ILE:O	1:A:324:VAL:HB	2.00	0.61
1:A:274:THR:HG22	1:A:277:GLN:N	2.03	0.61
1:A:30:ILE:HD11	1:A:79:LEU:CD1	2.26	0.61
1:A:287:LEU:HD23	1:A:288:PHE:CE2	2.35	0.61
1:A:72:LEU:HD13	1:A:73:LEU:N	2.15	0.60
1:A:19:ILE:HD13	1:A:19:ILE:N	2.16	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:30:ILE:CD1	1:A:79:LEU:CD1	2.82	0.57
1:A:293:ASP:HB3	1:A:295:THR:HB	1.85	0.57
1:A:391:ASP:HB3	1:A:395:ARG:HH21	1.69	0.56
1:A:202:MET:O	1:A:205:HIS:HB2	2.06	0.56
1:A:348:THR:O	1:A:349:ASP:CB	2.52	0.56
1:A:354:ILE:HG21	1:A:411:VAL:HG22	1.87	0.55
1:A:310:GLN:HA	1:A:310:GLN:OE1	2.06	0.55
1:A:85:LEU:H	1:A:272:GLN:HE22	1.58	0.52
1:A:274:THR:H	1:A:277:GLN:NE2	2.08	0.51
1:A:274:THR:H	1:A:277:GLN:HE21	1.59	0.51
1:A:295:THR:HG22	5:A:497:HOH:O	2.11	0.50
1:A:227:ASP:OD1	1:A:254:PHE:HA	2.12	0.50
1:A:244:VAL:CG1	1:A:246:ARG:HG3	2.44	0.47
1:A:259:ASN:HA	1:A:264:LYS:HB2	1.97	0.46
1:A:313:THR:HG23	1:A:315:ALA:H	1.79	0.46
1:A:61:GLN:HE22	1:A:193:LYS:HE3	1.81	0.46
1:A:190:CR2:N2	1:A:190:CR2:HD1	2.30	0.46
1:A:313:THR:HG22	1:A:316:GLU:HG3	1.97	0.45
1:A:142:LEU:HD23	1:A:142:LEU:C	2.35	0.45
1:A:25:LEU:HA	1:A:81:PHE:O	2.16	0.45
1:A:12:THR:HG23	1:A:381:LEU:CD2	2.48	0.44
1:A:75:ASP:OD2	1:A:205:HIS:HE1	2.01	0.44
1:A:287:LEU:HD23	1:A:288:PHE:CD2	2.53	0.43
1:A:141:GLU:HG3	5:A:509:HOH:O	2.18	0.43
1:A:300:GLU:O	1:A:304:VAL:HG23	2.20	0.42
1:A:29:TYR:CZ	1:A:76:ASN:HB3	2.55	0.42
1:A:85:LEU:H	1:A:272:GLN:NE2	2.17	0.42
1:A:180:PRO:HG2	1:A:265:LEU:HD12	2.01	0.42
1:A:337:PHE:CZ	1:A:341:MET:CE	3.03	0.42
1:A:85:LEU:N	1:A:85:LEU:HD12	2.34	0.42
1:A:12:THR:HG23	1:A:381:LEU:HD21	2.01	0.41
1:A:337:PHE:CZ	1:A:341:MET:HE2	2.56	0.41
1:A:44:LYS:HD2	5:A:522:HOH:O	2.18	0.41
1:A:61:GLN:HB2	1:A:220:ARG:HG2	2.03	0.41
1:A:284:ALA:HB3	1:A:308:LEU:HD21	2.02	0.41
1:A:408:GLU:H	1:A:408:GLU:CD	2.24	0.41
1:A:270:ARG:H	1:A:270:ARG:HG2	1.48	0.41
1:A:7:ARG:NH2	5:A:538:HOH:O	2.54	0.41
1:A:58:ASP:O	1:A:222:ILE:HA	2.21	0.40
1:A:38:GLY:HA3	1:A:63:ASN:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	378/415 (91%)	363 (96%)	8 (2%)	7 (2%)	9 7

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	272	GLN
1	A	349	ASP
1	A	200	ASP
1	A	68	ASP
1	A	204	GLN
1	A	345	MET
1	A	19	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	326/351 (93%)	300 (92%)	26 (8%)	13 13

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	24	SER
1	A	72	LEU
1	A	73	LEU
1	A	129	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	200	ASP
1	A	201	HIS
1	A	204	GLN
1	A	233	ARG
1	A	257	ASP
1	A	269	SER
1	A	270	ARG
1	A	271	ASP
1	A	272	GLN
1	A	274	THR
1	A	293	ASP
1	A	295	THR
1	A	305	MET
1	A	308	LEU
1	A	310	GLN
1	A	311	ASN
1	A	313	THR
1	A	314	GLU
1	A	349	ASP
1	A	350	SER
1	A	356	GLU
1	A	394	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	14	HIS
1	A	48	ASN
1	A	205	HIS
1	A	272	GLN
1	A	277	GLN
1	A	311	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CR2	A	190	1	20,20,21	3.45	5 (25%)	25,27,29	4.48	14 (56%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CR2	A	190	1	-	0/6/25/26	0/2/2/2

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	190	CR2	CA2-C2	-7.74	1.40	1.48
1	A	190	CR2	C2-N3	-4.26	1.29	1.39
1	A	190	CR2	O2-C2	2.02	1.27	1.23
1	A	190	CR2	C1-N2	3.77	1.40	1.32
1	A	190	CR2	CB2-CA2	11.50	1.44	1.35

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	190	CR2	O2-C2-CA2	-12.27	124.45	130.97
1	A	190	CR2	CA2-N2-C1	-4.80	102.18	105.76
1	A	190	CR2	C1-CA1-N1	-4.71	102.91	112.96
1	A	190	CR2	CB2-CA2-C2	-4.67	116.57	122.28
1	A	190	CR2	CG2-CB2-CA2	-3.87	125.69	130.10
1	A	190	CR2	CE1-CD1-CG2	-3.83	116.23	121.27
1	A	190	CR2	O3-C3-CA3	-2.99	116.50	126.38
1	A	190	CR2	CE2-CD2-CG2	-2.85	117.53	121.27
1	A	190	CR2	CA1-C1-N3	-2.53	119.00	122.67
1	A	190	CR2	CB2-CA2-N2	2.97	133.12	128.82
1	A	190	CR2	CA3-N3-C1	3.05	132.19	127.85
1	A	190	CR2	CA1-C1-N2	3.28	128.98	124.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	190	CR2	CD1-CG2-CD2	4.57	124.51	117.63
1	A	190	CR2	CA2-C2-N3	13.00	108.91	103.30

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	190	CR2	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	420	-	4,4,4	0.34	0	6,6,6	0.35	0
3	SO4	A	421	-	4,4,4	0.21	0	6,6,6	0.10	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	A	420	-	-	0/0/0/0	0/0/0/0
3	SO4	A	421	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	388/415 (93%)	-0.07	11 (2%) 53 65	20, 43, 87, 130	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	202	MET	4.5
1	A	203	LYS	3.7
1	A	416	ALA	3.4
1	A	1	GLY	3.1
1	A	25	LEU	3.0
1	A	24	SER	2.9
1	A	381	LEU	2.7
1	A	201	HIS	2.2
1	A	342	ALA	2.2
1	A	314	GLU	2.2
1	A	324	VAL	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CR2	A	190	19/20	0.96	0.11	23,28,33,33	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	CL	A	422	1/1	0.74	0.40	119,119,119,119	0
3	SO4	A	421	5/5	0.91	0.22	149,149,151,156	0
3	SO4	A	420	5/5	0.92	0.29	135,136,138,141	0
2	CA	A	419	1/1	0.94	0.17	51,51,51,51	0
2	CA	A	418	1/1	0.99	0.16	51,51,51,51	0

6.5 Other polymers [i](#)

There are no such residues in this entry.