



wwPDB NMR Structure Validation Summary Report ⓘ

Jun 13, 2024 – 01:01 PM EDT

PDB ID : 1O9A
Title : Solution structure of the complex of 1F12F1 from fibronectin with B3 from FnBB from *S. dysgalactiae*
Authors : Schwarz-Linek, U.; Werner, J.M.; Pickford, A.R.; Pilka, E.S.; Gurusiddappa, S.; Briggs, J.A.G.; Hook, M.; Campbell, I.D.; Potts, J.R.
Deposited on : 2002-12-11

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	93	
2	B	36	

2 Ensemble composition and analysis i

This entry contains 15 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:20-A:96, A:100-A:109, B:13-B:32 (107)	2.50	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 2, 3, 4, 5, 6, 7, 8, 9, 13, 14
2	10, 11, 12
Single-model clusters	15

3 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 1785 atoms, of which 852 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called FIBRONECTIN.

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		S
1	A	93	1404	447	671	132	145	9	0

- Molecule 2 is a protein called FIBRONECTIN BINDING PROTEIN.


Mol	Chain	Residues	Atoms					Trace
			Total	C	H	N	O	
2	B	24	381	124	181	31	45	0

4 Residue-property plots [i](#)


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: FIBRONECTIN




- Molecule 2: FIBRONECTIN BINDING PROTEIN




4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 1. Colouring as in section 4.1 above.

- Molecule 1: FIBRONECTIN



- Molecule 2: FIBRONECTIN BINDING PROTEIN



GLU	GLU	GLY	S20	S20	S20	D29	N30	N32	P33	K34	E35	D36
GLU	GLU	GLY	K21	K21	K21	D29	N30	N32	P33	K34	E35	D36
SER	SER	SER	P22	P22	P22	D29	N30	N32	P33	K34	E35	D36
LEU	LEU	LEU	K23	K23	K23	D29	N30	N32	P33	K34	E35	D36
PRO	PRO	PRO	L24	L24	L24	D29	N30	N32	P33	K34	E35	D36
THR	THR	THR	S25	S25	S25	D29	N30	N32	P33	K34	E35	D36
GLU	GLU	GLU	I26	I26	I26	D29	N30	N32	P33	K34	E35	D36
GLN	GLN	GLN	H27	H27	H27	D29	N30	N32	P33	K34	E35	D36
GLN	GLN	GLN	F28	F28	F28	D29	N30	N32	P33	K34	E35	D36
GLN	GLN	GLN	D29	D29	D29	D29	N30	N32	P33	K34	E35	D36
SER	SER	SER	N30	N30	N30	D29	N30	N32	P33	K34	E35	D36
SER	SER	SER	E31	E31	E31	D29	N30	N32	P33	K34	E35	D36
GLY	GLY	GLY	N32	N32	N32	D29	N30	N32	P33	K34	E35	D36
S13	S13	S13	P33	P33	P33	D29	N30	N32	P33	K34	E35	D36
T14	T14	T14	K34	K34	K34	D29	N30	N32	P33	K34	E35	D36
T15	T15	T15	E35	E35	E35	D29	N30	N32	P33	K34	E35	D36
E16	E16	E16	D36	D36	D36	D29	N30	N32	P33	K34	E35	D36
V17	V17	V17				D29	N30	N32	P33	K34	E35	D36

5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 15 were deposited, based on the following criterion: *LEAST RESTRAINT VIOLATION*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	refinement	
XPLOR	structure solution	3.8

No chemical shift data was provided.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	691	626	625	35±7
2	B	166	151	150	11±2
All	All	12855	11655	11625	536

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 22.

5 of 196 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:103:SER:HA	2:B:17:VAL:HB	0.82	1.49	9	4
1:A:43:LEU:HD22	1:A:43:LEU:H	0.80	1.37	9	10
1:A:80:THR:HG22	1:A:93:THR:HG22	0.76	1.55	15	11
1:A:101:ARG:HD3	1:A:102:ILE:N	0.73	1.98	12	1
1:A:43:LEU:HG	2:B:26:ILE:HD11	0.71	1.61	14	3

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR

entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	86/93 (92%)	67±2 (78±2%)	14±2 (16±2%)	5±1 (5±2%)	3	23
2	B	19/36 (53%)	10±1 (53±7%)	7±2 (37±8%)	2±2 (10±9%)	1	10
All	All	1575/1935 (81%)	1163 (74%)	314 (20%)	98 (6%)	3	19

5 of 32 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	40	GLY	11
1	A	24	ASN	11
1	A	96	GLY	8
1	A	87	SER	7
1	A	25	GLY	5

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	74/78 (95%)	46±4 (63±5%)	28±4 (37±5%)	1	7
2	B	20/34 (59%)	13±1 (67±6%)	7±1 (33±6%)	1	11
All	All	1410/1680 (84%)	895 (63%)	515 (37%)	1	8

5 of 78 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	43	LEU	15
1	A	69	LYS	15
1	A	89	ILE	15
1	A	106	ILE	15
1	A	88	MET	14

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided