



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 6, 2020 – 08:59 AM BST

PDB ID : 5O9X  
Title : Crystal structure of *Aspergillus fumigatus* N-acetylphosphoglucosamine mutase S69A in complex with glucose1,6bisphosphate  
Authors : Raimi, O.G.; Hurtado-Guerrero, R.  
Deposited on : 2017-06-20  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	549	

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 4657 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Phosphoacetylglucosamine mutase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	541	4152	2612	722	802	16	0	1	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	35	LYS	ARG	conflict	UNP A0A0S7E9S6
A	69	ALA	SER	engineered mutation	UNP A0A0S7E9S6
A	142	ILE	VAL	conflict	UNP A0A0S7E9S6
A	161	PHE	TYR	conflict	UNP A0A0S7E9S6
A	276	ILE	VAL	conflict	UNP A0A0S7E9S6
A	291	LEU	VAL	conflict	UNP A0A0S7E9S6
A	371	MET	LEU	conflict	UNP A0A0S7E9S6
A	467	GLU	ASP	conflict	UNP A0A0S7E9S6

- Molecule 2 is 1,6-di-O-phosphono-alpha-D-glucopyranose (three-letter code: G16) (formula:  $C_6H_{13}O_{12}P_2$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	O	P	0	0
			20	6	12	2		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Cl	0	0
			2	2		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	471	Total	O	0	0
			471	471		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphoacetylglucosamine mutase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	80.86Å 87.57Å 91.87Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.90 19.80 – 1.90	Depositor EDS
% Data completeness (in resolution range)	97.5 (20.00-1.90) 97.6 (19.80-1.90)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.64 (at 1.90Å)	Xtrriage
Refinement program	REFMAC 5.8.0158	Depositor
R, $R_{free}$	0.165 , 0.201 0.166 , 0.200	Depositor DCC
$R_{free}$ test set	519 reflections (1.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.4	Xtrriage
Anisotropy	0.037	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 48.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.011 for -h,l,k	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4657	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.89% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CL, G16

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.12	6/4225 (0.1%)	1.18	20/5723 (0.3%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	293	TYR	CG-CD1	6.06	1.47	1.39
1	A	346	GLU	CD-OE1	5.50	1.31	1.25
1	A	169	TYR	CG-CD1	5.42	1.46	1.39
1	A	226	ARG	CD-NE	-5.38	1.37	1.46
1	A	132	ARG	CZ-NH1	5.19	1.39	1.33
1	A	503	PHE	CG-CD1	5.05	1.46	1.38

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	226	ARG	NE-CZ-NH2	-21.30	109.65	120.30
1	A	140	ARG	NE-CZ-NH2	-17.50	111.55	120.30
1	A	226	ARG	NE-CZ-NH1	16.04	128.32	120.30
1	A	291	LEU	CB-CG-CD1	-15.51	84.63	111.00
1	A	140	ARG	NE-CZ-NH1	14.89	127.74	120.30
1	A	132	ARG	NE-CZ-NH2	-12.36	114.12	120.30
1	A	132	ARG	NE-CZ-NH1	11.95	126.27	120.30
1	A	522	ARG	NE-CZ-NH2	-11.64	114.48	120.30
1	A	522	ARG	NE-CZ-NH1	8.79	124.70	120.30
1	A	291	LEU	CA-CB-CG	7.94	133.56	115.30
1	A	432	MET	CG-SD-CE	-7.69	87.89	100.20
1	A	226	ARG	CG-CD-NE	-6.88	97.36	111.80
1	A	122	MET	CG-SD-CE	-6.86	89.23	100.20
1	A	132	ARG	CD-NE-CZ	6.86	133.20	123.60
1	A	251	ASP	CB-CG-OD1	5.97	123.67	118.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	167	LEU	CA-CB-CG	-5.89	101.76	115.30
1	A	226	ARG	CD-NE-CZ	5.86	131.80	123.60
1	A	140	ARG	CD-NE-CZ	5.83	131.77	123.60
1	A	291	LEU	CD1-CG-CD2	5.06	125.69	110.50
1	A	417	LEU	CB-CG-CD2	-5.05	102.41	111.00

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4152	0	4167	39	0
2	A	20	0	10	0	0
3	A	2	0	0	1	0
4	A	12	0	16	0	0
5	A	471	0	0	5	2
All	All	4657	0	4193	40	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:603:CL:CL	5:A:1116:HOH:O	2.15	0.99
1:A:25:PHE:H	1:A:101:ASN:HD21	1.05	0.98
1:A:218:ASN:HD22	1:A:246:ASP:H	1.15	0.92
1:A:61:TRP:H	1:A:175:ASN:HD21	1.24	0.85
1:A:132:ARG:NE	5:A:701:HOH:O	1.89	0.75
1:A:85:GLY:O	1:A:168:HIS:HD2	1.69	0.74
1:A:146:SER:O	1:A:150:THR:HG23	1.91	0.70
1:A:538:ASP:OD2	5:A:703:HOH:O	2.10	0.69
1:A:36:ALA:HA	1:A:39:LEU:HD12	1.79	0.64
1:A:43:VAL:HG22	1:A:78:VAL:HG21	1.81	0.63

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:386:THR:HB	1:A:426:GLY:HA2	1.85	0.58
1:A:61:TRP:N	1:A:175:ASN:HD21	1.98	0.58
1:A:49:LEU:HD22	1:A:96:ALA:HB2	1.86	0.56
1:A:218:ASN:ND2	1:A:246:ASP:H	1.96	0.56
1:A:43:VAL:HG22	1:A:78:VAL:CG2	2.37	0.54
1:A:172:ARG:HH12	1:A:423:GLN:HB2	1.72	0.54
1:A:132:ARG:NH2	5:A:701:HOH:O	2.38	0.54
1:A:79:LYS:NZ	1:A:168:HIS:HE1	2.08	0.51
1:A:501[A]:ARG:NH1	1:A:518:GLU:OE2	2.44	0.51
1:A:105:GLU:H	1:A:105:GLU:CD	2.16	0.49
1:A:132:ARG:HD3	1:A:142:ILE:HG12	1.94	0.48
1:A:323:SER:O	1:A:407:GLN:HG2	2.15	0.47
1:A:66:VAL:HG21	1:A:142:ILE:HD13	1.97	0.47
1:A:421:ILE:HA	1:A:421:ILE:HD13	1.52	0.45
1:A:421:ILE:HD11	1:A:434:LEU:HB2	1.99	0.45
1:A:112:ASP:HA	1:A:115:VAL:HG13	2.00	0.44
1:A:79:LYS:NZ	1:A:168:HIS:CE1	2.86	0.43
1:A:335:VAL:HB	1:A:378:TYR:HB3	2.00	0.43
1:A:537:ARG:O	1:A:541:THR:HB	2.19	0.42
1:A:334:VAL:O	1:A:356:SER:HA	2.20	0.42
1:A:291:LEU:HD13	1:A:291:LEU:N	2.35	0.42
1:A:316:PHE:CD2	1:A:438:ILE:HG12	2.56	0.41
1:A:46:VAL:HG13	1:A:80:LEU:HD11	2.03	0.41
1:A:218:ASN:HD22	1:A:246:ASP:N	1.98	0.41
1:A:140:ARG:NH2	5:A:718:HOH:O	2.52	0.41
1:A:164:THR:HB	1:A:165:PRO:HD3	2.01	0.41
1:A:310:ALA:HB2	1:A:379:PHE:HB2	2.03	0.41
1:A:66:VAL:HG21	1:A:142:ILE:CD1	2.51	0.41
1:A:220:VAL:O	1:A:224:LYS:HG2	2.20	0.41
1:A:326:ILE:HA	1:A:328:GLN:HE22	1.87	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:1092:HOH:O	5:A:1093:HOH:O[2_565]	2.11	0.09
5:A:1092:HOH:O	5:A:1115:HOH:O[2_565]	2.13	0.07

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	540/549 (98%)	521 (96%)	17 (3%)	2 (0%)	34 24

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	424	ALA
1	A	182	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	436/442 (99%)	426 (98%)	10 (2%)	50 45

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	LEU
1	A	142	ILE
1	A	234	GLU
1	A	235	ASP
1	A	291	LEU
1	A	467	GLU
1	A	468	ARG
1	A	501[A]	ARG
1	A	501[B]	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	541	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	101	ASN
1	A	168	HIS
1	A	175	ASN
1	A	218	ASN
1	A	367	HIS
1	A	403	GLN
1	A	498	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	GOL	A	605	-	5,5,5	0.36	0	5,5,5	0.33	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	GOL	A	604	-	5,5,5	0.57	0	5,5,5	0.55	0
2	G16	A	601	-	19,20,20	1.76	4 (21%)	30,31,31	1.64	4 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	605	-	-	2/4/4/4	-
4	GOL	A	604	-	-	2/4/4/4	-
2	G16	A	601	-	-	1/11/31/31	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	G16	O5-C1	4.72	1.53	1.41
2	A	601	G16	O4-C4	2.94	1.49	1.43
2	A	601	G16	P'-O1	2.63	1.64	1.59
2	A	601	G16	O3-C3	-2.26	1.37	1.43

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	G16	O5-C1-O1	-5.55	104.11	111.36
2	A	601	G16	O5-C5-C6	-3.53	99.55	106.67
2	A	601	G16	O3X-P'-O1X	3.03	119.21	107.64
2	A	601	G16	O4-C4-C3	2.02	115.02	110.35

There are no chirality outliers.


All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	605	GOL	O1-C1-C2-C3
4	A	604	GOL	O1-C1-C2-O2
4	A	604	GOL	O1-C1-C2-C3
4	A	605	GOL	O1-C1-C2-O2
2	A	601	G16	O5-C1-O1-P'

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	541/549 (98%)	-0.32	11 (2%) 65 68	13, 23, 49, 75	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	181	TYR	5.4
1	A	15	LEU	3.4
1	A	180	GLN	3.2
1	A	5	ALA	3.2
1	A	4	PRO	3.0
1	A	541	THR	2.8
1	A	178	GLY	2.8
1	A	234	GLU	2.4
1	A	48	LEU	2.1
1	A	235	ASP	2.1
1	A	21	GLU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	GOL	A	605	6/6	0.90	0.12	54,55,62,62	0
4	GOL	A	604	6/6	0.90	0.16	29,36,40,52	0
3	CL	A	602	1/1	0.98	0.13	26,26,26,26	0
2	G16	A	601	20/20	0.99	0.07	15,19,27,31	0
3	CL	A	603	1/1	1.00	0.04	20,20,20,20	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.