



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2018 – 05:48 pm GMT

PDB ID : 3OAU
Title : Antibody 2G12 Recognizes Di-Mannose Equivalently in Domain- and Non-Domain-Exchanged Forms, but only binds the HIV-1 Glycan Shield if Domain-Exchanged
Authors : Doores, K.J.; Fulton, Z.; Huber, M.; Wilson, I.A.; Burton, D.R.
Deposited on : 2010-08-05
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	5502 (1.90-1.90)
Clashscore	122126	6115 (1.90-1.90)
Ramachandran outliers	120053	6048 (1.90-1.90)
Sidechain outliers	120020	6048 (1.90-1.90)
RSRZ outliers	108989	5379 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	225	
2	L	212	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3514 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fab 2G12, heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	H	218	1651	1043	282	319	7	0	3	0

- Molecule 2 is a protein called Fab 2G12, light chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	L	212	1629	1025	273	326	5	0	2	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
3	H	1	5	4	1	0	0
3	H	1	5	4	1	0	0

- Molecule 4 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	H	1	Total	C	O	0	0
			12	6	6		
4	H	1	Total	C	O	0	0
			11	6	5		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	L	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	H	76	Total 76	O 76	0	0
6	L	119	Total 119	O 119	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Fab 2G12, heavy chain



- Molecule 2: Fab 2G12, light chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	39.30Å 56.63Å 57.65Å 92.91° 105.05° 92.74°	Depositor
Resolution (Å)	19.75 – 1.90 18.81 – 1.90	Depositor EDS
% Data completeness (in resolution range)	(Not available) (19.75-1.90) 85.8 (18.81-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.82 (at 1.90Å)	Xtrriage
Refinement program	BUSTER-TNT BUSTER 2.8.0, BUSTER 2.8.0	Depositor
R, R_{free}	0.212 , 0.243 0.214 , 0.251	Depositor DCC
R_{free} test set	1633 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	28.6	Xtrriage
Anisotropy	0.547	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 64.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3514	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	H	0.45	0/1698	0.68	0/2312
2	L	0.48	0/1671	0.67	0/2269
All	All	0.46	0/3369	0.67	0/4581

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1651	0	1633	19	0
2	L	1629	0	1592	9	0
3	H	10	0	0	0	0
4	H	23	0	21	0	0
5	L	6	0	7	0	0
6	H	76	0	0	0	0
6	L	119	0	0	0	0
All	All	3514	0	3253	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:198:HIS:CD2	2:L:200:GLY:H	2.14	0.66
2:L:20:THR:HG23	2:L:72:THR:HG23	1.79	0.62
1:H:173:VAL:HG22	1:H:192[B]:VAL:HG22	1.83	0.61
1:H:42:GLY:H	1:H:43:GLY:HA2	1.64	0.61
1:H:11:LEU:HD21	1:H:156:PHE:HE1	1.65	0.60
1:H:152:VAL:HG11	1:H:160:VAL:HG11	1.82	0.60
2:L:155:GLN:HE21	2:L:158:ASN:HD21	1.49	0.60
1:H:154:ASP:HB3	1:H:185[B]:LEU:HD23	1.88	0.56
1:H:129:PRO:HB3	1:H:155:TYR:HB3	1.93	0.51
1:H:205:ILE:HG12	1:H:220:LYS:HG2	1.91	0.51
2:L:198:HIS:HD2	2:L:200:GLY:H	1.60	0.50
2:L:141:PRO:O	2:L:198:HIS:HE1	1.94	0.50
1:H:154:ASP:HA	1:H:185[B]:LEU:HB3	1.94	0.50
1:H:11:LEU:HD21	1:H:156:PHE:CE1	2.46	0.49
2:L:146:VAL:HG13	2:L:194:CYS:SG	2.52	0.49
1:H:136:PRO:HD3	1:H:148:LEU:HB3	1.96	0.48
1:H:148:LEU:HD21	1:H:204:TYR:CD2	2.50	0.46
1:H:133:PRO:HD3	1:H:219:LYS:HE2	1.99	0.44
1:H:3:GLN:HB3	1:H:25:SER:HB2	1.99	0.44
2:L:19:ILE:HD11	2:L:75:ILE:HD12	1.98	0.44
2:L:24:ARG:NE	2:L:70:GLU:OE2	2.50	0.44
1:H:42:GLY:N	1:H:43:GLY:HA2	2.33	0.43
1:H:60:TYR:HB2	1:H:65:LYS:HG2	2.01	0.43
1:H:136:PRO:HG3	1:H:199:LEU:HD22	2.02	0.41
1:H:40:VAL:O	1:H:43:GLY:HA2	2.20	0.41
1:H:177:PRO:HD2	2:L:162:SER:OG	2.21	0.40
1:H:39:ARG:HB2	1:H:95:TYR:HE2	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	217/225 (96%)	210 (97%)	6 (3%)	1 (0%)	31	20
2	L	212/212 (100%)	206 (97%)	6 (3%)	0	100	100
All	All	429/437 (98%)	416 (97%)	12 (3%)	1 (0%)	49	40

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	42	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	186/190 (98%)	182 (98%)	4 (2%)	55	49
2	L	184/183 (100%)	178 (97%)	6 (3%)	41	32
All	All	370/373 (99%)	360 (97%)	10 (3%)	48	40

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	102	ASP
1	H	161	THR
1	H	202	GLN
1	H	216	LYS
2	L	33	LEU
2	L	81	ASP
2	L	105	GLU
2	L	147	GLN
2	L	152	ASN
2	L	188	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	H	84	HIS
1	H	181	GLN
1	H	202	GLN
2	L	89	GLN
2	L	90	HIS
2	L	155	GLN
2	L	198	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	H	226	-	4,4,4	0.24	0	6,6,6	0.18	0
3	SO4	H	227	-	4,4,4	0.13	0	6,6,6	0.12	0
4	MAN	H	228	4	12,12,12	0.48	0	17,17,17	0.77	0
4	MAN	H	229	4	11,11,12	0.38	0	15,15,17	0.98	0
5	GOL	L	1	-	5,5,5	1.52	1 (20%)	5,5,5	1.22	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	H	226	-	-	0/0/0/0	0/0/0/0
3	SO4	H	227	-	-	0/0/0/0	0/0/0/0
4	MAN	H	228	4	-	0/2/22/22	0/1/1/1
4	MAN	H	229	4	-	0/2/19/22	0/1/1/1
5	GOL	L	1	-	-	0/4/4/4	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	L	1	GOL	O2-C2	-3.01	1.34	1.43

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	L	1	GOL	O2-C2-C3	-2.30	98.48	109.00

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	H	218/225 (96%)	0.59	24 (11%) 5 6	29, 49, 72, 101	0
2	L	212/212 (100%)	0.16	12 (5%) 24 27	25, 39, 64, 90	0
All	All	430/437 (98%)	0.38	36 (8%) 11 13	25, 44, 68, 101	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	144	GLY	5.2
1	H	41	PRO	4.9
1	H	40	VAL	4.6
1	H	200	GLY	4.3
1	H	137	SER	4.3
1	H	224	LYS	4.3
1	H	103	ARG	4.2
1	H	145	THR	4.0
2	L	212	GLY	4.0
2	L	154	LEU	3.8
1	H	42	GLY	3.8
1	H	223	PRO	3.7
1	H	168	ALA	3.5
1	H	196	SER	3.4
2	L	168	SER	3.3
2	L	169	LYS	2.8
1	H	1	GLU	2.7
1	H	211	LYS	2.7
2	L	188	LYS	2.6
1	H	68	PHE	2.6
1	H	182	SER	2.5
1	H	170	THR	2.5
1	H	166	SER	2.3
2	L	153	ALA	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	L	35	TRP	2.3
2	L	86	TYR	2.3
1	H	57	TYR	2.2
1	H	14	ALA	2.2
2	L	57	GLY	2.2
1	H	49	ALA	2.2
2	L	213	GLU	2.1
2	L	136	LEU	2.1
1	H	26	ASN	2.1
1	H	102	ASP	2.0
1	H	62	ASP	2.0
2	L	139	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GOL	L	1	6/6	0.82	0.16	38,48,50,53	0
4	MAN	H	228	12/12	0.84	0.15	48,61,72,82	0
4	MAN	H	229	11/12	0.97	0.11	32,38,44,50	0
3	SO4	H	227	5/5	0.98	0.13	73,77,78,78	0
3	SO4	H	226	5/5	0.98	0.13	71,75,76,76	0

6.5 Other polymers [i](#)

There are no such residues in this entry.