



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 17, 2024 – 02:08 AM EDT

PDB ID : 5OAU
Title : Penicillin-Binding Protein 2X (PBP2X) from Streptococcus pneumoniae
Authors : Bernardo-Garcia, N.; Hermoso, J.A.
Deposited on : 2017-06-23
Resolution : 2.67 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.67 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3863 (2.70-2.66)
Clashscore	141614	4210 (2.70-2.66)
Ramachandran outliers	138981	4141 (2.70-2.66)
Sidechain outliers	138945	4141 (2.70-2.66)
RSRZ outliers	127900	3780 (2.70-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	702	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5360 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Penicillin-binding protein 2X.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	667	5131	3216	856	1036	23	0	0	0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	229	Total	O	0	0
			229	229		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Penicillin-binding protein 2X



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	100.26Å 100.26Å 189.81Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	34.78 – 2.67 34.78 – 2.67	Depositor EDS
% Data completeness (in resolution range)	99.9 (34.78-2.67) 100.0 (34.78-2.67)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	23.18 (at 2.68Å)	Xtrriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.173 , 0.225 0.180 , 0.226	Depositor DCC
R_{free} test set	1506 reflections (4.69%)	wwPDB-VP
Wilson B-factor (Å ²)	54.9	Xtrriage
Anisotropy	0.023	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 52.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.001 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5360	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.80	2/5221 (0.0%)	0.88	8/7072 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	338	THR	CB-CG2	-5.87	1.32	1.52
1	A	528	TYR	CE1-CZ	-5.14	1.31	1.38

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	82	ARG	NE-CZ-NH2	-6.78	116.91	120.30
1	A	573	VAL	CB-CA-C	-5.72	100.53	111.40
1	A	426	ARG	NE-CZ-NH2	-5.53	117.53	120.30
1	A	82	ARG	NE-CZ-NH1	5.39	122.99	120.30
1	A	512	ARG	NE-CZ-NH1	5.37	122.98	120.30
1	A	648	ASP	CB-CG-OD1	5.31	123.08	118.30
1	A	384	ARG	NE-CZ-NH1	-5.26	117.67	120.30
1	A	426	ARG	NE-CZ-NH1	5.04	122.82	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	596	SER	Peptide
1	A	715	ASN	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5131	0	5051	51	0
2	A	229	0	0	3	0
All	All	5360	0	5051	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:678:LYS:HD3	1:A:679:ASN:H	1.11	1.13
1:A:678:LYS:HD3	1:A:679:ASN:N	1.73	1.01
1:A:76:ARG:NH1	1:A:220:GLU:OE1	2.20	0.75
1:A:140:GLN:HA	1:A:143:GLN:HE21	1.52	0.75
1:A:678:LYS:CD	1:A:679:ASN:H	1.98	0.67
1:A:123:ALA:HA	1:A:137:VAL:HG21	1.80	0.63
1:A:233:ILE:HD11	1:A:252:SER:HB3	1.81	0.62
1:A:95:ASN:HB2	1:A:181:THR:HG23	1.81	0.62
1:A:338:THR:HG21	1:A:465:PHE:HZ	1.63	0.62
1:A:338:THR:HG23	1:A:573:VAL:CG2	2.31	0.61
1:A:671:ASN:ND2	1:A:713:TRP:CE2	2.69	0.60
1:A:527:MET:CE	1:A:546:LEU:HB3	2.31	0.59
1:A:466:THR:HG21	1:A:475:LEU:HB2	1.85	0.59
1:A:71:THR:HG22	1:A:233:ILE:HG22	1.84	0.58
1:A:234:ILE:HD13	1:A:251:VAL:HG12	1.84	0.57
1:A:338:THR:CG2	1:A:465:PHE:HZ	2.16	0.57
1:A:518:VAL:HG12	1:A:527:MET:HG3	1.87	0.55
1:A:338:THR:CG2	1:A:465:PHE:CZ	2.91	0.53
1:A:257:ASP:H	1:A:487:ASN:HD21	1.57	0.53
1:A:234:ILE:CD1	1:A:251:VAL:HG12	2.38	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:689:LEU:HD12	1:A:713:TRP:CE3	2.44	0.52
1:A:671:ASN:ND2	1:A:713:TRP:NE1	2.58	0.52
1:A:76:ARG:NH2	1:A:224:ASN:OD1	2.43	0.51
1:A:544:VAL:HG12	2:A:801:HOH:O	2.10	0.51
1:A:527:MET:HE2	1:A:546:LEU:HB3	1.91	0.50
1:A:250:GLN:O	1:A:251:VAL:HG22	2.12	0.50
1:A:95:ASN:HB2	1:A:181:THR:CG2	2.43	0.49
1:A:527:MET:HE3	1:A:546:LEU:HD13	1.95	0.48
1:A:494:SER:HB3	1:A:657:LEU:HD22	1.95	0.47
1:A:527:MET:HE3	1:A:546:LEU:HB3	1.94	0.47
1:A:456:VAL:HG12	1:A:457:THR:N	2.29	0.47
1:A:102:GLU:O	1:A:104:TYR:N	2.48	0.47
1:A:159:THR:HG22	2:A:846:HOH:O	2.15	0.47
1:A:186:ARG:HD3	1:A:201:ALA:HB3	1.97	0.47
1:A:223:LEU:HD23	1:A:655:ARG:HG2	1.97	0.47
1:A:396:SER:O	1:A:398:VAL:N	2.48	0.47
1:A:492:ARG:HB2	1:A:657:LEU:HD12	1.97	0.47
1:A:340:LYS:CD	1:A:400:MET:HG3	2.45	0.46
1:A:206:ASN:O	1:A:208:ASP:N	2.49	0.46
1:A:526:THR:HG21	2:A:903:HOH:O	2.16	0.46
1:A:456:VAL:HG11	1:A:461:MET:HG2	1.98	0.45
1:A:234:ILE:CG2	1:A:249:GLU:HG3	2.46	0.45
1:A:518:VAL:HG12	1:A:527:MET:CG	2.46	0.45
1:A:67:GLN:HA	1:A:236:TYR:O	2.19	0.42
1:A:340:LYS:HD2	1:A:400:MET:HG3	2.00	0.42
1:A:93:SER:HB2	1:A:157:GLY:HA2	2.02	0.41
1:A:491:ALA:O	1:A:633:GLN:HA	2.20	0.41
1:A:95:ASN:HD22	1:A:181:THR:HG23	1.83	0.41
1:A:526:THR:O	1:A:526:THR:HG22	2.20	0.41
1:A:96:VAL:HG12	1:A:180:PHE:CE2	2.56	0.41
1:A:667:THR:HG22	1:A:667:THR:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	661/702 (94%)	614 (93%)	38 (6%)	9 (1%)	11	25

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	105	LYS
1	A	251	VAL
1	A	207	GLU
1	A	715	ASN
1	A	440	ASP
1	A	108	THR
1	A	285	LYS
1	A	176	LYS
1	A	252	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	564/590 (96%)	553 (98%)	11 (2%)	55	79

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	81	ASP
1	A	130	LEU
1	A	205	GLU
1	A	284	VAL
1	A	427	PHE
1	A	506	ASP
1	A	573	VAL
1	A	595	TYR
1	A	600	LEU
1	A	678	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	689	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	103	ASN
1	A	143	GLN
1	A	148	GLN
1	A	394	HIS
1	A	487	ASN
1	A	529	ASN
1	A	599	GLN
1	A	671	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	667/702 (95%)	0.16	44 (6%) 18 16	33, 55, 118, 171	0

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	236	TYR	14.5
1	A	319	THR	10.0
1	A	740	ASP	6.7
1	A	594	HIS	6.1
1	A	248	THR	6.0
1	A	593	GLU	5.9
1	A	320	GLU	5.4
1	A	66	HIS	4.8
1	A	318	ILE	4.8
1	A	249	GLU	4.6
1	A	742	LYS	4.3
1	A	67	GLN	4.2
1	A	68	THR	4.1
1	A	743	LYS	4.1
1	A	207	GLU	4.0
1	A	237	GLU	3.9
1	A	314	THR	3.8
1	A	103	ASN	3.8
1	A	238	LYS	3.4
1	A	316	GLU	3.4
1	A	208	ASP	3.3
1	A	625	LYS	3.2
1	A	321	ASP	3.2
1	A	322	PHE	3.1
1	A	741	ILE	3.1
1	A	739	LYS	3.1
1	A	250	GLN	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	210	SER	3.1
1	A	104	TYR	3.0
1	A	206	ASN	2.9
1	A	323	VAL	2.8
1	A	145	ASN	2.8
1	A	70	ARG	2.5
1	A	599	GLN	2.5
1	A	142	SER	2.4
1	A	592	PRO	2.3
1	A	251	VAL	2.3
1	A	750	ASP	2.3
1	A	287	LYS	2.3
1	A	719	GLU	2.2
1	A	234	ILE	2.2
1	A	716	ILE	2.1
1	A	440	ASP	2.1
1	A	675	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.