



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 4, 2023 – 12:48 PM EDT

PDB ID : 6OAG  
Title : Crystal structure of human FPPS in complex with an allosteric inhibitor YF-02-82  
Authors : Park, J.; Berghuis, A.M.  
Deposited on : 2019-03-16  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.35.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	F	375	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PO4	F	403[B]	-	-	X	-

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2890 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Farnesyl pyrophosphate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	F	343	2722	1761	441	509	11	0	2	0

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	-21	MET	-	initiating methionine	UNP P14324
F	-20	GLY	-	expression tag	UNP P14324
F	-19	SER	-	expression tag	UNP P14324
F	-18	SER	-	expression tag	UNP P14324
F	-17	HIS	-	expression tag	UNP P14324
F	-16	HIS	-	expression tag	UNP P14324
F	-15	HIS	-	expression tag	UNP P14324
F	-14	HIS	-	expression tag	UNP P14324
F	-13	HIS	-	expression tag	UNP P14324
F	-12	HIS	-	expression tag	UNP P14324
F	-11	SER	-	expression tag	UNP P14324
F	-10	SER	-	expression tag	UNP P14324
F	-9	GLY	-	expression tag	UNP P14324
F	-8	ARG	-	expression tag	UNP P14324
F	-7	GLU	-	expression tag	UNP P14324
F	-6	ASN	-	expression tag	UNP P14324
F	-5	LEU	-	expression tag	UNP P14324
F	-4	TYR	-	expression tag	UNP P14324
F	-3	PHE	-	expression tag	UNP P14324
F	-2	GLN	-	expression tag	UNP P14324
F	-1	GLY	-	expression tag	UNP P14324
F	0	HIS	-	expression tag	UNP P14324

- Molecule 2 is [(1S)-1-{{6-(3-chloro-4-methylphenyl)thieno[2,3-d]pyrimidin-4-yl}amino}-2-phenylethyl]phosphonic acid (three-letter code: M2Y) (formula: C<sub>21</sub>H<sub>19</sub>ClN<sub>3</sub>O<sub>3</sub>PS).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	
			Total	C	Cl	N	O	P			S
2	F	1	30	21	1	3	3	1	1	0	0
2	F	1	30	21	1	3	3	1	1	0	0

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O P		
3	F	1	10	8 2	0	1


- Molecule 4 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	F	98	Total	O	0	0
			98	98		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Farnesyl pyrophosphate synthase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	111.35Å 111.35Å 76.86Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	55.74 – 2.30 55.67 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.3 (55.74-2.30) 99.4 (55.67-2.30)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.36 (at 2.29Å)	Xtrriage
Refinement program	REFMAC 5.8.0238	Depositor
R, $R_{free}$	0.193 , 0.241 0.205 , 0.254	Depositor DCC
$R_{free}$ test set	1047 reflections (4.78%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	64.7	Xtrriage
Anisotropy	0.034	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 65.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	2890	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	88.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.97% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: M2Y, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	F	1.19	20/2782 (0.7%)	0.91	4/3782 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	F	0	1

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	13	GLU	CD-OE2	16.11	1.43	1.25
1	F	13	GLU	CD-OE1	13.75	1.40	1.25
1	F	273	GLN	CD-NE2	11.44	1.61	1.32
1	F	93	GLU	CG-CD	11.36	1.69	1.51
1	F	80	ASP	CG-OD2	11.17	1.51	1.25
1	F	70	GLU	CD-OE1	6.70	1.33	1.25
1	F	11	ALA	C-O	6.46	1.35	1.23
1	F	180	GLN	CD-NE2	6.27	1.48	1.32
1	F	114	GLY	C-O	5.63	1.32	1.23
1	F	39	GLY	C-O	5.61	1.32	1.23
1	F	63	THR	C-O	5.57	1.33	1.23
1	F	215	TYR	C-O	5.55	1.33	1.23
1	F	62	LEU	C-O	5.51	1.33	1.23
1	F	177	THR	C-O	5.37	1.33	1.23
1	F	318	GLU	CD-OE2	5.32	1.31	1.25
1	F	89	GLY	CA-C	5.20	1.60	1.51
1	F	264	ASP	C-O	5.17	1.33	1.23
1	F	111	THR	CB-OG1	5.11	1.53	1.43

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	70	GLU	CD-OE2	5.05	1.31	1.25
1	F	199	TYR	C-O	5.05	1.32	1.23

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	80	ASP	CB-CG-OD2	-10.46	108.88	118.30
1	F	273	GLN	CG-CD-NE2	-5.69	103.05	116.70
1	F	11	ALA	O-C-N	5.17	130.97	122.70
1	F	13	GLU	OE1-CD-OE2	5.07	129.38	123.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	F	264	ASP	Mainchain

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	2722	0	2621	30	0
2	F	60	0	0	0	0
3	F	10	0	0	4	0
4	F	98	0	0	3	0
All	All	2890	0	2621	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:23:GLN:HE22	1:F:83:GLN:NE2	1.65	0.95
3:F:403[A]:PO4:O1	4:F:501:HOH:O	2.09	0.69

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:271:VAL:HG11	1:F:290:TYR:CZ	2.27	0.69
1:F:271:VAL:HG12	1:F:305:TYR:OH	1.94	0.68
1:F:23:GLN:HE22	1:F:83:GLN:HE22	1.45	0.65
1:F:271:VAL:CG1	1:F:305:TYR:OH	2.45	0.64
1:F:273:GLN:OE1	1:F:277:ARG:NH1	2.32	0.61
1:F:257:LYS:HA	4:F:583:HOH:O	2.04	0.57
1:F:289:ASN:HB3	1:F:297:LYS:O	2.07	0.54
1:F:59:ASN:ND2	3:F:403[B]:PO4:O3	2.37	0.54
1:F:58:TYR:N	3:F:403[B]:PO4:O1	2.40	0.54
1:F:166:GLN:NE2	1:F:203[A]:PHE:CE1	2.76	0.54
1:F:285:ILE:O	1:F:289:ASN:ND2	2.42	0.52
1:F:58:TYR:HA	1:F:93:GLU:OE1	2.12	0.49
1:F:60:ARG:NH1	3:F:403[B]:PO4:O2	2.46	0.48
1:F:99:PHE:CE2	1:F:167:THR:HG21	2.48	0.48
1:F:124:VAL:O	1:F:127:ASP:HB2	2.14	0.48
1:F:310:LEU:N	1:F:311:PRO:CD	2.77	0.47
1:F:245:TYR:HE1	1:F:311:PRO:HA	1.83	0.44
1:F:271:VAL:HG13	1:F:305:TYR:OH	2.18	0.43
1:F:68:PHE:CE1	1:F:72:VAL:HG21	2.53	0.43
1:F:96:GLN:O	1:F:100:LEU:HG	2.18	0.43
1:F:322:TYR:CE1	1:F:326:MET:HE2	2.53	0.43
1:F:23:GLN:HG3	1:F:26:ARG:NH2	2.33	0.43
1:F:252:PRO:HB3	1:F:258:ILE:HG12	2.01	0.42
1:F:235:MET:HE2	1:F:235:MET:HB3	1.84	0.42
1:F:60:ARG:NH1	1:F:96:GLN:HE22	2.17	0.41
1:F:201:THR:HG22	4:F:513:HOH:O	2.19	0.41
1:F:29:THR:O	1:F:32:GLU:HG3	2.22	0.40
1:F:235:MET:CE	1:F:344:LEU:HB3	2.51	0.40
1:F:212:ALA:O	1:F:216:MET:HG3	2.20	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	F	343/375 (92%)	325 (95%)	18 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	F	275/322 (85%)	268 (98%)	7 (2%)	47	65

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	30	GLU
1	F	81	SER
1	F	186	VAL
1	F	204	TYR
1	F	243	ASP
1	F	253	SER
1	F	335	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	F	20	HIS
1	F	83	GLN
1	F	96	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	PO4	F	403[A]	-	4,4,4	0.60	0	6,6,6	0.47	0
2	M2Y	F	402	-	28,33,33	3.69	11 (39%)	36,48,48	2.12	10 (27%)
2	M2Y	F	401	-	28,33,33	3.25	11 (39%)	36,48,48	2.15	8 (22%)
3	PO4	F	403[B]	-	4,4,4	0.63	0	6,6,6	0.47	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	M2Y	F	402	-	-	1/18/18/18	0/4/4/4
2	M2Y	F	401	-	-	3/18/18/18	0/4/4/4

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	402	M2Y	CAJ-SAK	-10.59	1.59	1.72
2	F	402	M2Y	PBA-OB	8.22	1.63	1.49
2	F	401	M2Y	PBA-OB	7.42	1.62	1.49
2	F	402	M2Y	C4-C5	-7.24	1.35	1.44

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	401	M2Y	C4-C5	-7.01	1.36	1.44
2	F	401	M2Y	PBA-OBC	6.01	1.64	1.54
2	F	401	M2Y	CAS-CAP	-5.54	1.39	1.51
2	F	402	M2Y	CAS-CAP	-5.48	1.40	1.51
2	F	401	M2Y	C2-N1	4.96	1.40	1.32
2	F	401	M2Y	CAL-CAT	-4.73	1.39	1.51
2	F	402	M2Y	CAL-CAT	-4.64	1.40	1.51
2	F	402	M2Y	PBA-OBC	4.52	1.62	1.54
2	F	402	M2Y	C2-N1	4.49	1.39	1.32
2	F	401	M2Y	PBA-OBB	-4.32	1.48	1.54
2	F	401	M2Y	CAJ-SAK	-4.06	1.67	1.72
2	F	402	M2Y	C6-C5	-4.00	1.35	1.42
2	F	401	M2Y	C6-C5	-3.76	1.36	1.42
2	F	402	M2Y	PBA-OBB	-3.75	1.48	1.54
2	F	402	M2Y	CAB-NAA	3.38	1.48	1.45
2	F	401	M2Y	C2-N3	3.30	1.40	1.33
2	F	402	M2Y	C2-N3	2.89	1.39	1.33
2	F	401	M2Y	CAB-NAA	2.85	1.47	1.45

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	402	M2Y	N1-C2-N3	-7.43	117.07	128.68
2	F	401	M2Y	N1-C2-N3	-7.23	117.37	128.68
2	F	401	M2Y	C5-C6-SAK	-4.56	107.55	112.24
2	F	402	M2Y	C2-N3-C4	4.39	120.35	116.59
2	F	401	M2Y	C5-C4-NAA	-4.35	116.52	120.63
2	F	402	M2Y	C5-C6-SAK	-4.28	107.84	112.24
2	F	401	M2Y	C2-N3-C4	3.99	120.01	116.59
2	F	402	M2Y	OBD-PBA-CAB	-3.74	106.39	113.80
2	F	401	M2Y	OBC-PBA-OBD	-3.68	104.21	113.45
2	F	402	M2Y	OBC-PBA-OBD	-3.57	104.47	113.45
2	F	401	M2Y	C2-N1-C6	2.93	120.31	113.45
2	F	402	M2Y	C2-N1-C6	2.89	120.22	113.45
2	F	401	M2Y	OBB-PBA-OBC	2.68	114.83	107.64
2	F	402	M2Y	C5-C4-NAA	-2.44	118.33	120.63
2	F	402	M2Y	OBB-PBA-OBC	2.38	114.05	107.64
2	F	402	M2Y	CAL-CAB-NAA	2.19	114.09	111.40
2	F	401	M2Y	NAA-C4-N3	2.09	121.24	118.06
2	F	402	M2Y	PBA-CAB-CAL	-2.02	108.13	111.30

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	F	401	M2Y	NAA-CAB-PBA-OBD
2	F	401	M2Y	CAL-CAB-PBA-OBD
2	F	401	M2Y	NAA-CAB-PBA-OBB
2	F	402	M2Y	NAA-CAB-CAL-CAT


There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	403[A]	PO4	1	0
3	F	403[B]	PO4	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	F	343/375 (91%)	0.66	32 (9%) <b>8</b> <b>11</b>	51, 81, 147, 167	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	33	MET	9.7
1	F	278	ALA	7.3
1	F	280	PRO	4.7
1	F	34	GLY	4.6
1	F	283	TYR	4.4
1	F	277	ARG	4.4
1	F	275	LEU	3.8
1	F	284	GLN	3.7
1	F	286	LEU	3.6
1	F	185	LEU	3.6
1	F	279	THR	3.4
1	F	276	GLN	3.2
1	F	301	VAL	3.2
1	F	35	HIS	3.1
1	F	254	VAL	3.1
1	F	273	GLN	2.9
1	F	281	GLU	2.9
1	F	285	ILE	2.8
1	F	272	VAL	2.7
1	F	82	LEU	2.7
1	F	298	VAL	2.5
1	F	32	GLU	2.5
1	F	290	TYR	2.4
1	F	296	GLU	2.2
1	F	308	LEU	2.2
1	F	313	VAL	2.1
1	F	287	LYS	2.1

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	F	186	VAL	2.1
1	F	303	ALA	2.0
1	F	101	VAL	2.0
1	F	10	TYR	2.0
1	F	258	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	M2Y	F	401	30/30	0.76	0.27	99,146,154,163	0
2	M2Y	F	402	30/30	0.88	0.23	123,147,168,171	0
3	PO4	F	403[A]	5/5	0.89	0.28	75,79,83,83	5
3	PO4	F	403[B]	5/5	0.89	0.28	88,88,92,96	5




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around M2Y F 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around M2Y F 402:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.