



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 10, 2023 – 08:25 pm GMT

PDB ID : 1ODZ  
Title : Expansion of the glycosynthase repertoire to produce defined manno-oligosaccharides  
Authors : Jahn, M.; Stoll, D.; Warren, R.A.J.; Szabo, L.; Singh, P.; Gilbert, H.J.; Ducros, V.M.A.; Davies, G.J.; Withers, S.G.  
Deposited on : 2003-03-17  
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*





The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1714 (1.40-1.40)
Clashscore	141614	1812 (1.40-1.40)
Ramachandran outliers	138981	1763 (1.40-1.40)
Sidechain outliers	138945	1762 (1.40-1.40)
RSRZ outliers	127900	1674 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	386	 2% 93% 5%
1	B	386	 3% 92% 5%
2	C	2	 100%
2	D	2	 50% 50%

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 7200 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mannan endo-1,4-beta-mannosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	379	3051	1947	522	578	4	0	6	0
1	B	379	3050	1947	522	577	4	0	6	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	38	MET	-	initiating methionine	UNP P49424
A	320	GLY	GLU	engineered mutation	UNP P49424
B	38	MET	-	initiating methionine	UNP P49424
B	320	GLY	GLU	engineered mutation	UNP P49424

- Molecule 2 is an oligosaccharide called beta-D-mannopyranose-(1-4)-alpha-D-mannopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	C	2	23	12	11	0	0	0
2	D	2	23	12	11	0	0	0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Zn	0	0
			1	1		

- Molecule 4 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: C<sub>4</sub>H<sub>12</sub>NO<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			8	4	1	3		
4	B	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Na	0	0
			1	1		
5	B	1	Total	Na	0	0
			1	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	494	Total	O	0	0
			494	494		

Continued on next page...


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
6	B	539	Total 539	O 539	0	0

### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Mannan endo-1,4-beta-mannosidase

Chain A: 



- Molecule 1: Mannan endo-1,4-beta-mannosidase

Chain B: 



- Molecule 2: beta-D-mannopyranose-(1-4)-alpha-D-mannopyranose

Chain C: 



- Molecule 2: beta-D-mannopyranose-(1-4)-alpha-D-mannopyranose

Chain D: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	131.85Å 131.85Å 54.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.40 14.94 – 1.40	Depositor EDS
% Data completeness (in resolution range)	93.7 (30.00-1.40) 93.8 (14.94-1.40)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.40 (at 1.40Å)	Xtrriage
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.151 , 0.177 0.162 , 0.188	Depositor DCC
$R_{free}$ test set	8545 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	13.6	Xtrriage
Anisotropy	0.046	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 51.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.025 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	7200	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 73.80 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.7193e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, TRS, NA, ZN, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.52	0/3166	0.81	11/4323 (0.3%)
1	B	0.54	0/3165	0.81	7/4322 (0.2%)
All	All	0.53	0/6331	0.81	18/8645 (0.2%)

There are no bond length outliers.

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	123[A]	ASP	CB-CG-OD2	8.04	125.53	118.30
1	B	123[B]	ASP	CB-CG-OD2	8.04	125.53	118.30
1	B	103	ASP	CB-CG-OD2	7.17	124.75	118.30
1	A	103	ASP	CB-CG-OD2	6.79	124.41	118.30
1	A	293[A]	ASP	CB-CG-OD2	6.76	124.38	118.30
1	A	293[B]	ASP	CB-CG-OD2	6.76	124.38	118.30
1	B	170	ASP	CB-CG-OD2	6.68	124.31	118.30
1	A	170	ASP	CB-CG-OD2	6.65	124.29	118.30
1	A	145	ASP	CB-CG-OD2	6.30	123.97	118.30
1	A	123[A]	ASP	CB-CG-OD2	6.22	123.89	118.30
1	A	123[B]	ASP	CB-CG-OD2	6.22	123.89	118.30
1	B	145	ASP	CB-CG-OD2	6.04	123.74	118.30
1	A	91	ASP	CB-CG-OD2	5.93	123.64	118.30
1	B	91	ASP	CB-CG-OD2	5.82	123.54	118.30
1	A	261	ASP	CB-CG-OD2	5.55	123.30	118.30
1	A	402	ASP	CB-CG-OD2	5.34	123.11	118.30
1	A	197	ASP	CB-CG-OD2	5.08	122.87	118.30
1	B	410	ASP	CB-CG-OD2	5.05	122.84	118.30

There are no chirality outliers.

There are no planarity outliers.



## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3051	0	2894	15	0
1	B	3050	0	2894	17	0
2	C	23	0	21	0	0
2	D	23	0	21	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	8	0	11	0	0
4	B	8	0	11	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	494	0	0	7	0
6	B	539	0	0	8	0
All	All	7200	0	5852	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:123[A]:ASP:OD2	6:B:2162:HOH:O	1.53	1.19
1:A:123[B]:ASP:OD1	6:A:2150:HOH:O	1.59	1.17
1:B:201:ARG:CZ	6:B:2310:HOH:O	2.05	1.03
1:B:201:ARG:NE	6:B:2310:HOH:O	2.02	0.92
1:A:123[A]:ASP:OD2	6:A:2151:HOH:O	1.92	0.88
1:A:293[B]:ASP:OD2	6:A:2368:HOH:O	1.98	0.79
1:A:190[B]:GLU:OE1	6:A:2267:HOH:O	2.08	0.71
1:A:190[B]:GLU:OE1	6:A:2268:HOH:O	2.09	0.69
1:B:190[B]:GLU:OE1	6:B:2291:HOH:O	2.11	0.67
1:B:123[A]:ASP:HB2	6:B:2164:HOH:O	1.96	0.64
1:A:123[A]:ASP:HB2	6:A:2150:HOH:O	1.98	0.63
1:B:79:HIS:HD2	1:B:82:THR:OG1	1.85	0.60
1:A:79:HIS:HD2	1:A:82:THR:OG1	1.85	0.59
1:B:89:ARG:HD2	6:B:2088:HOH:O	2.06	0.55
1:B:66:GLN:HG2	1:B:315:ILE:HD11	1.92	0.51

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:79:HIS:HE1	1:B:121:GLU:OE1	1.93	0.51
1:A:293[A]:ASP:OD2	1:A:296:ARG:NH2	2.43	0.50
1:B:293[A]:ASP:OD1	1:B:296:ARG:NH2	2.41	0.50
1:A:79:HIS:HE1	1:A:121:GLU:OE1	1.96	0.49
1:A:390:ASN:HD22	1:A:390:ASN:C	2.15	0.47
1:A:79:HIS:CD2	1:A:82:THR:OG1	2.67	0.47
1:A:66:GLN:HA	1:A:66:GLN:OE1	2.14	0.47
1:B:201:ARG:CD	6:B:2310:HOH:O	2.54	0.45
1:B:79:HIS:CD2	1:B:82:THR:OG1	2.69	0.45
1:B:201:ARG:HD2	6:B:2310:HOH:O	2.17	0.44
1:B:390:ASN:C	1:B:390:ASN:HD22	2.21	0.43
1:A:293[B]:ASP:CG	6:A:2368:HOH:O	2.50	0.43
1:A:390:ASN:ND2	1:A:392:THR:H	2.17	0.41
1:B:104:PHE:CD1	1:B:134:ARG:HD3	2.54	0.41
1:B:81:THR:HA	1:B:94:GLN:O	2.21	0.40
1:B:390:ASN:ND2	1:B:392:THR:H	2.19	0.40
1:A:198:GLU:HG2	1:A:199:GLN:NE2	2.37	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	381/386 (99%)	374 (98%)	6 (2%)	1 (0%)	41	18
1	B	381/386 (99%)	374 (98%)	7 (2%)	0	100	100
All	All	762/772 (99%)	748 (98%)	13 (2%)	1 (0%)	51	23

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	154	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	321/322 (100%)	318 (99%)	3 (1%)	78	58
1	B	321/322 (100%)	318 (99%)	3 (1%)	78	58
All	All	642/644 (100%)	636 (99%)	6 (1%)	86	58

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	319[A]	SER
1	A	319[B]	SER
1	A	390	ASN
1	B	123[A]	ASP
1	B	123[B]	ASP
1	B	390	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	79	HIS
1	A	152	GLN
1	A	164	GLN
1	A	199	GLN
1	A	389	ASN
1	A	390	ASN
1	A	408	ASN
1	B	79	HIS
1	B	164	GLN
1	B	389	ASN
1	B	390	ASN
1	B	408	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	MAN	C	1	2	12,12,12	0.78	0	17,17,17	1.80	5 (29%)
2	BMA	C	2	2	11,11,12	0.84	0	15,15,17	1.07	2 (13%)
2	MAN	D	1	2	12,12,12	0.79	0	17,17,17	1.85	5 (29%)
2	BMA	D	2	2	11,11,12	0.68	0	15,15,17	0.69	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MAN	C	1	2	-	0/2/22/22	0/1/1/1
2	BMA	C	2	2	-	0/2/19/22	0/1/1/1
2	MAN	D	1	2	-	0/2/22/22	0/1/1/1
2	BMA	D	2	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	1	MAN	C1-O5-C5	-4.29	105.57	113.66
2	C	1	MAN	C1-O5-C5	-3.69	106.70	113.66
2	D	1	MAN	O1-C1-O5	3.19	119.97	110.38
2	C	2	BMA	O5-C1-C2	-2.95	106.22	110.77
2	C	1	MAN	O1-C1-C2	2.87	117.11	109.03

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	MAN	O5-C1-C2	2.85	115.38	110.28
2	D	1	MAN	O4-C4-C5	-2.83	102.27	109.30
2	C	1	MAN	O4-C4-C5	-2.77	102.42	109.30
2	D	1	MAN	O1-C1-C2	2.62	116.41	109.03
2	C	1	MAN	C1-C2-C3	-2.40	105.34	110.31
2	C	2	BMA	C1-O5-C5	2.29	115.30	112.19
2	D	1	MAN	C1-C2-C3	-2.00	106.16	110.31


There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	TRS	B	1426	3	7,7,7	0.33	0	9,9,9	0.49	0
4	TRS	A	1424	3	7,7,7	0.47	0	9,9,9	0.47	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	TRS	B	1426	3	-	1/9/9/9	-
4	TRS	A	1424	3	-	0/9/9/9	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	1426	TRS	C1-C-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	379/386 (98%)	-0.32	9 (2%) 59 58	10, 14, 25, 39	1 (0%)
1	B	379/386 (98%)	-0.27	11 (2%) 51 50	10, 14, 25, 39	2 (0%)
All	All	758/772 (98%)	-0.30	20 (2%) 56 55	10, 14, 25, 39	3 (0%)

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	372	GLY	7.8
1	A	422	VAL	5.4
1	B	372	GLY	4.9
1	B	373	THR	4.9
1	A	373	THR	4.5
1	B	369	GLY	4.5
1	B	422	VAL	4.1
1	A	369	GLY	4.1
1	B	198	GLU	3.5
1	B	42	VAL	3.3
1	A	42	VAL	3.0
1	A	198	GLU	2.8
1	B	385	PRO	2.4
1	A	199	GLN	2.4
1	B	199	GLN	2.4
1	A	89	ARG	2.2
1	B	389	ASN	2.2
1	A	389	ASN	2.1
1	B	88	THR	2.1
1	B	290	ASP	2.1



## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.


## 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MAN	D	1	12/12	0.96	0.07	11,12,15,15	0
2	BMA	D	2	11/12	0.96	0.06	11,12,14,14	0
2	MAN	C	1	12/12	0.97	0.05	11,12,14,15	0
2	BMA	C	2	11/12	0.97	0.06	11,12,14,14	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	TRS	A	1424	8/8	0.96	0.06	13,15,15,17	0
4	TRS	B	1426	8/8	0.96	0.07	14,15,16,18	0
3	ZN	A	1423	1/1	1.00	0.02	13,13,13,13	0
3	ZN	B	1423	1/1	1.00	0.03	13,13,13,13	0
5	NA	A	1427	1/1	1.00	0.03	12,12,12,12	0
5	NA	B	1427	1/1	1.00	0.03	13,13,13,13	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.