



# Full wwPDB X-ray Structure Validation Report i

Nov 5, 2023 – 11:59 pm GMT

PDB ID : 8OG1  
Title : Exostosin-like 3 apo enzyme  
Authors : Sammon, D.; Hohenester, E.  
Deposited on : 2023-03-17  
Resolution : 1.58 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 1.58 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5534 (1.60-1.56)
Clashscore	141614	5861 (1.60-1.56)
Ramachandran outliers	138981	5708 (1.60-1.56)
RSRZ outliers	127900	5431 (1.60-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	C	5	-	-	-	X

## 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 11911 atoms, of which 5720 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Exostosin-like 3.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	715	11372	3716	5622	978	1024	32	0	5	0

There are 23 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	29	ALA	-	expression tag	UNP O43909
A	30	PRO	-	expression tag	UNP O43909
A	31	LEU	-	expression tag	UNP O43909
A	32	VAL	-	expression tag	UNP O43909
A	33	HIS	-	expression tag	UNP O43909
A	34	HIS	-	expression tag	UNP O43909
A	35	HIS	-	expression tag	UNP O43909
A	36	HIS	-	expression tag	UNP O43909
A	37	HIS	-	expression tag	UNP O43909
A	38	HIS	-	expression tag	UNP O43909
A	39	ALA	-	expression tag	UNP O43909
A	40	LEU	-	expression tag	UNP O43909
A	41	ASP	-	expression tag	UNP O43909
A	42	GLU	-	expression tag	UNP O43909
A	43	ASN	-	expression tag	UNP O43909
A	44	LEU	-	expression tag	UNP O43909
A	45	TYR	-	expression tag	UNP O43909
A	46	PHE	-	expression tag	UNP O43909
A	47	GLN	-	expression tag	UNP O43909
A	48	GLY	-	expression tag	UNP O43909
A	49	ALA	-	expression tag	UNP O43909
A	50	LEU	-	expression tag	UNP O43909
A	51	ALA	-	expression tag	UNP O43909

- Molecule 2 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
2	B	3	73	22	34	2	15	0	0	0

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
3	C	6	139	42	64	3	30	0	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	327	Total O		0	0
			327	327		


### 3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Exostosin-like 3



- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	120.90Å 90.00°    120.90Å 90.00°    127.45Å 120.00°	Depositor
Resolution (Å)	34.76 – 1.58 34.90 – 1.58	Depositor EDS
% Data completeness (in resolution range)	99.9 (34.76-1.58) 99.9 (34.90-1.58)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	1.25 (at 1.58Å)	Xtriage
Refinement program	PHENIX 1.18rc1_3769, PHENIX 1.18rc1_3769	Depositor
$R$ , $R_{free}$	0.184 , 0.211 0.184 , 0.211	Depositor DCC
$R_{free}$ test set	7425 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.1	Xtriage
Anisotropy	0.267	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.43 , 44.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.019 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	11911	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.24% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [\(i\)](#)

### 5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, CSO, MAN, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.52	2/5922 (0.0%)	0.70	4/8064 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	763	GLU	CB-CG	-6.51	1.39	1.52
1	A	396	VAL	CB-CG1	-5.23	1.41	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	854	ARG	NE-CZ-NH2	7.15	123.88	120.30
1	A	475[A]	MET	CG-SD-CE	6.42	110.47	100.20
1	A	475[B]	MET	CG-SD-CE	6.42	110.47	100.20
1	A	854	ARG	NE-CZ-NH1	-5.52	117.54	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	520	PHE	Peptide

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5750	5622	5633	51	1
2	B	39	34	34	0	0
3	C	75	64	64	1	0
4	A	327	0	0	8	0
All	All	6191	5720	5731	51	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:851:VAL:HG12	1:A:852:THR:HG23	1.57	0.87
1:A:854:ARG:HH21	1:A:854:ARG:HG3	1.54	0.73
1:A:308:TYR:OH	4:A:1002:HOH:O	2.08	0.71
1:A:447[B]:CYS:SG	4:A:1288:HOH:O	2.50	0.69
1:A:698:SER:O	1:A:719:ARG:NH1	2.27	0.67
1:A:589:TYR:OH	4:A:1003:HOH:O	2.13	0.61
1:A:411:GLU:OE1	1:A:444:SER:OG	2.09	0.59
1:A:205:VAL:HG21	1:A:259[B]:GLN:HG3	1.85	0.57
1:A:205:VAL:HG22	1:A:259[A]:GLN:HG2	1.86	0.57
1:A:283:ASP:OD2	1:A:405:LYS:HD3	2.09	0.52
1:A:337:VAL:HG22	1:A:506:LEU:HD22	1.90	0.52
1:A:756:PHE:CE2	1:A:851:VAL:HG13	2.44	0.51
1:A:646:SER:OG	1:A:647:GLY:N	2.45	0.50
1:A:395:LEU:C	1:A:395:LEU:HD23	2.32	0.50
1:A:537:ARG:NH1	4:A:1006:HOH:O	2.30	0.50
1:A:620:ASP:OD1	1:A:621:PRO:HD2	2.12	0.50
1:A:763:GLU:CG	4:A:1044:HOH:O	2.59	0.50
1:A:280:ARG:NH2	1:A:281:LYS:HD2	2.27	0.49
1:A:745:ASP:OD2	4:A:1004:HOH:O	2.19	0.49
1:A:205:VAL:CG2	1:A:259[B]:GLN:HG3	2.43	0.49
1:A:209:TYR:CE2	1:A:251:LEU:HD13	2.49	0.48
1:A:763:GLU:HG2	4:A:1044:HOH:O	2.12	0.48
1:A:349:LYS:HE3	1:A:410:THR:O	2.14	0.48

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:550:ALA:HB3	1:A:734:GLU:CD	2.35	0.47
1:A:550:ALA:HB3	1:A:734:GLU:OE1	2.14	0.47
1:A:475[A]:MET:HG3	1:A:533:MET:HG2	1.97	0.47
1:A:672:ARG:HB3	1:A:675:VAL:HG22	1.96	0.47
1:A:168:LEU:N	1:A:169:PRO:HD2	2.29	0.47
1:A:701:LEU:HB3	1:A:702:PRO:CD	2.46	0.46
1:A:221:THR:HG23	1:A:524:ASP:OD1	2.16	0.45
1:A:743:ILE:CG1	1:A:747:ALA:HB3	2.47	0.45
1:A:475[A]:MET:HE3	1:A:533:MET:HA	2.00	0.44
1:A:550:ALA:HB3	1:A:734:GLU:OE2	2.18	0.44
1:A:763:GLU:HG2	1:A:763:GLU:O	2.18	0.43
1:A:557:SER:HA	1:A:708:TRP:O	2.18	0.43
1:A:670:TYR:CD1	1:A:698:SER:HA	2.54	0.43
1:A:697:ASN:OD1	1:A:720:THR:HG21	2.19	0.43
1:A:340:LYS:HE2	1:A:393:GLN:OE1	2.19	0.42
1:A:396:VAL:CG1	1:A:398:PHE:CZ	3.02	0.42
1:A:739:ALA:HB2	1:A:761:TRP:CZ3	2.53	0.42
1:A:789:SER:OG	3:C:1:NAG:H81	2.20	0.42
1:A:396:VAL:CG1	1:A:398:PHE:CE2	3.02	0.42
1:A:780:ILE:HB	1:A:781:PRO:HD3	2.01	0.41
1:A:748:HIS:NE2	1:A:859:CYS:HB3	2.35	0.41
1:A:706:LEU:HD12	1:A:706:LEU:HA	1.86	0.41
1:A:396:VAL:HG12	1:A:398:PHE:CE2	2.55	0.41
1:A:442:VAL:HG12	1:A:469:GLN:HB2	2.03	0.41
1:A:668:LEU:HD12	1:A:695:VAL:HB	2.02	0.41
1:A:588[B]:ARG:NH2	4:A:1015:HOH:O	2.54	0.40
1:A:213:LEU:HD12	1:A:213:LEU:HA	1.91	0.40
1:A:691:LYS:HD3	1:A:716:MET:HG3	2.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:763:GLU:OE2	1:A:895:THR:HG1[6_555]	1.59	0.01

## 5.3 Torsion angles (i)

### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	709/891 (80%)	682 (96%)	27 (4%)	0	100 100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains (i)

There are no protein residues with a non-rotameric sidechain to report in this entry.

### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	CSO	A	606	1	3,6,7	0.93	0	0,6,8	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSO	A	606	1	-	0/1/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [\(i\)](#)

9 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	B	1	2,1	14,14,15	0.56	0	17,19,21	0.67	0
2	NAG	B	2	2	14,14,15	0.54	0	17,19,21	0.41	0
2	BMA	B	3	2	11,11,12	1.30	2 (18%)	15,15,17	0.92	0
3	NAG	C	1	3,1	14,14,15	0.35	0	17,19,21	0.60	0
3	NAG	C	2	3	14,14,15	0.87	1 (7%)	17,19,21	0.50	0
3	BMA	C	3	3	11,11,12	1.18	1 (9%)	15,15,17	0.92	0
3	MAN	C	4	3	11,11,12	1.23	1 (9%)	15,15,17	1.59	4 (26%)
3	NAG	C	5	3	14,14,15	0.23	0	17,19,21	0.50	0
3	MAN	C	6	3	11,11,12	1.19	1 (9%)	15,15,17	1.35	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	0/6/23/26	0/1/1/1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1
2	BMA	B	3	2	-	2/2/19/22	0/1/1/1
3	NAG	C	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	C	2	3	-	0/6/23/26	0/1/1/1
3	BMA	C	3	3	-	0/2/19/22	0/1/1/1
3	MAN	C	4	3	-	2/2/19/22	0/1/1/1
3	NAG	C	5	3	-	2/6/23/26	0/1/1/1
3	MAN	C	6	3	-	1/2/19/22	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	2	NAG	O5-C1	-2.96	1.39	1.43
2	B	3	BMA	C2-C3	2.85	1.56	1.52
3	C	6	MAN	C1-C2	2.75	1.58	1.52
3	C	3	BMA	O5-C1	-2.49	1.39	1.43
2	B	3	BMA	C1-C2	2.27	1.57	1.52
3	C	4	MAN	O6-C6	2.18	1.51	1.42

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	6	MAN	C1-O5-C5	4.05	117.68	112.19
3	C	4	MAN	O2-C2-C1	2.83	114.95	109.15
3	C	4	MAN	C1-C2-C3	-2.60	106.47	109.67
3	C	4	MAN	O5-C5-C6	2.39	110.96	107.20
3	C	4	MAN	O2-C2-C3	-2.19	105.75	110.14

There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	C	4	MAN	C4-C5-C6-O6
3	C	5	NAG	C1-C2-N2-C7
3	C	4	MAN	O5-C5-C6-O6
2	B	3	BMA	C4-C5-C6-O6
3	C	5	NAG	O5-C5-C6-O6
2	B	3	BMA	O5-C5-C6-O6
3	C	6	MAN	O5-C5-C6-O6


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry (i)

There are no ligands in this entry.

## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

## 6 Fit of model and data [\(i\)](#)

### 6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	714/891 (80%)	0.27	64 (8%) <span style="background-color: red;">9</span> <span style="background-color: red;">9</span>	20, 31, 68, 105	0

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	705	ASP	7.7
1	A	700	LYS	7.5
1	A	350	ILE	7.5
1	A	168	LEU	7.3
1	A	707	LEU	7.2
1	A	170	PRO	6.8
1	A	580	THR	6.6
1	A	169	PRO	6.5
1	A	173	ALA	6.3
1	A	703	SER	6.2
1	A	706	LEU	5.9
1	A	171	PRO	5.8
1	A	701	LEU	5.7
1	A	704	GLU	5.4
1	A	250	VAL	5.1
1	A	163	LYS	5.0
1	A	559	LYS	4.8
1	A	172	LYS	4.6
1	A	910	HIS	4.5
1	A	160	LEU	4.4
1	A	403	GLN	4.3
1	A	374	ASP	4.2
1	A	581	GLU	3.9
1	A	328	PHE	3.9
1	A	860	PRO	3.9
1	A	699	PRO	3.8
1	A	579	GLU	3.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	908	LEU	3.6
1	A	869	ASP	3.5
1	A	918	PHE	3.5
1	A	289	TYR	3.4
1	A	917	LYS	3.3
1	A	719	ARG	3.3
1	A	863	PRO	3.3
1	A	551	ALA	3.2
1	A	306	VAL	3.2
1	A	290	ASN	3.2
1	A	550	ALA	3.1
1	A	857	PHE	3.0
1	A	175	ARG	2.9
1	A	254	ALA	2.9
1	A	402	ASN	2.9
1	A	856	THR	2.8
1	A	671	GLU	2.8
1	A	558	GLY	2.8
1	A	674	GLU	2.7
1	A	582	PRO	2.6
1	A	859	CYS	2.6
1	A	722	LYS	2.6
1	A	557	SER	2.5
1	A	761	TRP	2.5
1	A	174	THR	2.4
1	A	743	ILE	2.4
1	A	708	TRP	2.3
1	A	768	ILE	2.3
1	A	909	PRO	2.3
1	A	804	PHE	2.2
1	A	913	THR	2.2
1	A	702	PRO	2.2
1	A	251	LEU	2.2
1	A	670	TYR	2.2
1	A	870	ASP	2.2
1	A	710	ASP	2.1
1	A	740	ILE	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	CSO	A	606	7/8	0.98	0.04	19,24,31,37	0


### 6.3 Carbohydrates [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	NAG	C	5	14/15	0.52	0.65	83,116,147,162	0
2	BMA	B	3	11/12	0.54	0.39	77,99,128,129	0
3	MAN	C	6	11/12	0.63	0.26	60,82,101,121	0
3	MAN	C	4	11/12	0.75	0.21	44,65,99,102	0
3	BMA	C	3	11/12	0.88	0.14	43,53,71,72	0
2	NAG	B	2	14/15	0.90	0.15	39,57,81,84	0
3	NAG	C	1	14/15	0.94	0.07	32,39,45,48	0
3	NAG	C	2	14/15	0.94	0.11	34,47,59,68	0
2	NAG	B	1	14/15	0.96	0.07	26,34,44,44	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.