



Full wwPDB X-ray Structure Validation Report i

Dec 6, 2023 – 07:36 am GMT

PDB ID : 8OIZ
Title : Crystal structure of human CRBN-DDB1 in complex with Pomalidomide
Authors : Le Bihan, Y.-V.; Cabry, M.P.; van Montfort, R.L.M.
Deposited on : 2023-03-23
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 11765 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA damage-binding protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	A	1116	Total	C 8329	N 5313	O 1382	S 1586	48	6	7	1

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1141	TRP	-	expression tag	UNP Q16531
A	1142	SER	-	expression tag	UNP Q16531
A	1143	HIS	-	expression tag	UNP Q16531
A	1144	PRO	-	expression tag	UNP Q16531
A	1145	GLN	-	expression tag	UNP Q16531
A	1146	PHE	-	expression tag	UNP Q16531
A	1147	GLU	-	expression tag	UNP Q16531
A	1148	LYS	-	expression tag	UNP Q16531


- Molecule 2 is a protein called Protein cereblon.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
2	B	379	Total	C 2900	N 1865	O 480	S 530	25	0	3	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	36	GLY	-	expression tag	UNP Q96SW2
B	37	PRO	-	expression tag	UNP Q96SW2
B	38	HIS	-	expression tag	UNP Q96SW2
B	39	MET	-	expression tag	UNP Q96SW2

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total Zn 1 1	0	0

- Molecule 5 is S-Pomalidomide (three-letter code: Y70) (formula: C₁₃H₁₁N₃O₄) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total 20	C 13	N 3	O 4	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	377	Total 379		O 379	0
6	B	112	Total 112		O 112	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA damage-binding protein 1



- Molecule 2: Protein cereblon



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	71.84 Å 128.51 Å 198.38 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.59 – 2.50 49.59 – 2.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.59-2.50) 100.0 (49.59-2.50)	Depositor EDS
R_{merge}	0.24	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	1.31 (at 2.51 Å)	Xtriage
Refinement program	BUSTER 2.10.4 (17-FEB-2023)	Depositor
R , R_{free}	0.213 , 0.262 0.208 , 0.260	Depositor DCC
R_{free} test set	3287 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å ²)	55.7	Xtriage
Anisotropy	0.425	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 71.7	EDS
L-test for twinning ²	$< L > = 0.52$, $< L^2 > = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	11765	wwPDB-VP
Average B, all atoms (Å ²)	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, EDO, Y70

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.38	0/8487	0.60	0/11516
2	B	0.35	0/2972	0.52	0/4045
All	All	0.37	0/11459	0.58	0/15561

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8329	0	7749	62	0
2	B	2900	0	2661	16	0
3	A	24	0	36	1	0
4	B	1	0	0	0	0
5	B	20	0	11	0	0
6	A	379	0	0	2	0
6	B	112	0	0	0	0
All	All	11765	0	10457	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:69:PRO:HG2	1:A:72:GLU:HG3	1.52	0.90
1:A:908:ASN:HD21	1:A:947:ARG:HH12	1.19	0.86
1:A:596:PHE:HB3	1:A:661:SER:HB2	1.57	0.84
1:A:545:PRO:HA	1:A:553:SER:HB2	1.65	0.75
1:A:984:THR:HG23	1:A:985:THR:HG22	1.79	0.65
1:A:47:GLU:H	1:A:47:GLU:CD	2.00	0.65
1:A:449:MET:HG2	1:A:484:LYS:O	1.97	0.64
1:A:812:TYR:CZ	2:B:241:PRO:HB3	2.32	0.64
1:A:217:SER:HB2	2:B:204:LYS:HB2	1.80	0.63
1:A:482:GLU:CB	1:A:483:PRO:HD3	2.29	0.62
1:A:1012:LEU:HD23	1:A:1140:HIS:HA	1.80	0.62
2:B:146:GLU:OE2	2:B:148:GLN:NE2	2.33	0.62
2:B:143:ALA:HB3	2:B:158:LYS:HB2	1.82	0.61
1:A:545:PRO:CA	1:A:553:SER:HB2	2.31	0.61
1:A:181:VAL:HG22	1:A:190:VAL:HG22	1.84	0.60
1:A:908:ASN:HD21	1:A:947:ARG:NH1	1.96	0.59
1:A:595:THR:HG22	1:A:600:HIS:CD2	2.38	0.58
1:A:239:TYR:HB3	1:A:246:LEU:HB2	1.85	0.58
1:A:1055:GLN:HG2	1:A:1093:LEU:HD23	1.86	0.58
1:A:840:GLU:HG3	6:A:1385:HOH:O	2.06	0.56
1:A:837:TYR:HB2	1:A:840:GLU:HG2	1.87	0.56
1:A:280:LEU:HD23	1:A:347:VAL:HG21	1.88	0.55
1:A:11:LYS:HD3	1:A:38:ARG:HD2	1.88	0.55
2:B:75:ASP:HB3	2:B:186:PRO:HB3	1.87	0.55
1:A:165:ILE:HG13	2:B:207:ILE:HD12	1.89	0.55
2:B:290:ILE:HA	2:B:424:PRO:HG3	1.90	0.53
1:A:434:ARG:HD3	1:A:445:GLU:OE1	2.08	0.53
1:A:546:LEU:HD22	1:A:595:THR:HG23	1.91	0.53
1:A:471:ILE:HG12	1:A:476:VAL:HG13	1.89	0.52
1:A:358:PRO:HD2	1:A:380:GLY:HA2	1.93	0.51
1:A:161:GLU:HA	3:A:1205:EDO:H11	1.92	0.51
1:A:791:LEU:HD23	1:A:858:LEU:HD21	1.94	0.49
1:A:315:THR:HG22	1:A:323:PHE:HB3	1.95	0.49
1:A:393:GLY:HA2	1:A:708:GLN:HE21	1.77	0.49
1:A:595:THR:CG2	1:A:600:HIS:CD2	2.96	0.49
2:B:61:GLY:HA3	2:B:145:ARG:NH2	2.26	0.49
1:A:480:SER:HB2	1:A:483:PRO:HD2	1.94	0.49
1:A:143:ILE:HG12	1:A:154:ALA:HB2	1.95	0.49
1:A:25:SER:HA	1:A:74:LYS:HD3	1.95	0.48
1:A:725[A]:CYS:SG	1:A:829:PHE:CE1	3.06	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:414:ARG:HG2	1:A:422:ASP:HA	1.96	0.48
1:A:492:GLU:HG3	1:A:512:VAL:HG21	1.95	0.48
2:B:199:LEU:HD13	2:B:237:LEU:HD11	1.96	0.48
1:A:889:ARG:HE	1:A:904:ASN:HD21	1.62	0.48
1:A:962:ASP:O	1:A:979:LYS:NZ	2.46	0.47
1:A:213:GLU:OE2	1:A:215:GLU:HB2	2.15	0.47
2:B:146:GLU:HG2	2:B:155:VAL:HG22	1.97	0.47
1:A:883:SER:HB2	1:A:911:ALA:CB	2.45	0.46
1:A:883:SER:HB2	1:A:911:ALA:HB3	1.95	0.46
1:A:984:THR:HG23	1:A:985:THR:N	2.30	0.46
1:A:216:ALA:HA	1:A:233:GLY:HA2	1.97	0.46
1:A:725[A]:CYS:SG	1:A:829:PHE:HE1	2.38	0.46
1:A:189:HIS:HB3	1:A:210:GLU:HA	1.98	0.45
1:A:710:LEU:HD21	1:A:1141:TRP:CD2	2.51	0.45
1:A:412:PRO:HD3	1:A:680:CYS:HB2	1.98	0.45
1:A:190:VAL:HG23	1:A:212:VAL:HG11	1.99	0.44
1:A:979:LYS:HA	6:A:1302:HOH:O	2.16	0.44
1:A:432:GLN:HG2	1:A:433:THR:N	2.32	0.44
1:A:762:SER:O	1:A:803:HIS:HA	2.18	0.43
1:A:385:GLY:HA3	1:A:719:GLU:O	2.18	0.43
1:A:671:VAL:HG12	1:A:673:LEU:HB2	2.00	0.43
2:B:323:CYS:HB3	2:B:326:CYS:HB2	2.01	0.43
1:A:970:ASN:HA	1:A:971:ALA:HA	1.85	0.42
1:A:58:TYR:HB3	1:A:1073:TRP:HB2	2.02	0.42
1:A:486:LEU:HD21	1:A:489:GLU:HB2	2.01	0.42
2:B:99:LEU:HB2	2:B:157:VAL:HG22	2.02	0.41
1:A:1054:MET:HG3	1:A:1112:LEU:HD11	2.01	0.41
2:B:57:HIS:CE1	2:B:98:PRO:HG3	2.55	0.41
2:B:60:LEU:HD23	2:B:60:LEU:HA	1.80	0.41
1:A:650:PHE:CD1	1:A:679:MET:HG2	2.55	0.41
1:A:230:ILE:HD11	1:A:285:LEU:HD21	2.03	0.41
2:B:84:LEU:HD23	2:B:109:MET:HE3	2.03	0.40
2:B:388:VAL:HG13	2:B:397:HIS:HD2	1.86	0.40
1:A:741:GLU:HG2	1:A:751:ALA:HA	2.04	0.40
1:A:935:TYR:O	1:A:937:PRO:HD3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	1109/1148 (97%)	1066 (96%)	43 (4%)	0	100 100
2	B	374/407 (92%)	366 (98%)	8 (2%)	0	100 100
All	All	1483/1555 (95%)	1432 (97%)	51 (3%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	812/1007 (81%)	781 (96%)	31 (4%)	33 58
2	B	279/369 (76%)	270 (97%)	9 (3%)	39 65
All	All	1091/1376 (79%)	1051 (96%)	40 (4%)	34 60

All (40) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	SER
1	A	38	ARG
1	A	101	ILE
1	A	127	GLU
1	A	159	LEU
1	A	198	ARG
1	A	203	ASN
1	A	243	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	352	THR
1	A	391	ARG
1	A	394	ILE
1	A	401	SER
1	A	421	THR
1	A	427	LEU
1	A	460	CYS
1	A	598	SER
1	A	670	ASN
1	A	691	LEU
1	A	706	GLU
1	A	722	ARG
1	A	749	THR
1	A	761	LEU
1	A	767	SER
1	A	845	GLN
1	A	896	GLU
1	A	928	ARG
1	A	931	LEU
1	A	941	ASN
1	A	957	VAL
1	A	991	HIS
1	A	1122	ARG
2	B	89	MET
2	B	91	LEU
2	B	199	LEU
2	B	203	ASN
2	B	266	GLU
2	B	324	LYS
2	B	340	LEU
2	B	373	ARG
2	B	403	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	203	ASN
1	A	209	GLN
1	A	255	GLN
1	A	456	GLN
1	A	600	HIS
1	A	648	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	708	GLN
1	A	904	ASN
1	A	907	ASN
1	A	908	ASN
1	A	941	ASN
1	A	1056	ASN
2	B	86	GLN
2	B	198	GLN
2	B	203	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

Of 8 ligands modelled in this entry, 1 is monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	EDO	A	1201	-	3,3,3	0.27	0	2,2,2	0.06	0
3	EDO	A	1206	-	3,3,3	0.48	0	2,2,2	0.06	0
3	EDO	A	1203	-	3,3,3	0.20	0	2,2,2	0.30	0
3	EDO	A	1204	-	3,3,3	0.42	0	2,2,2	0.19	0
3	EDO	A	1202	-	3,3,3	0.30	0	2,2,2	0.23	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	Y70	B	502	-	22,22,22	0.40	0	31,33,33	0.62	2 (6%)
3	EDO	A	1205	-	3,3,3	0.36	0	2,2,2	0.11	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	1201	-	-	1/1/1/1	-
3	EDO	A	1206	-	-	1/1/1/1	-
3	EDO	A	1203	-	-	1/1/1/1	-
3	EDO	A	1204	-	-	0/1/1/1	-
3	EDO	A	1202	-	-	0/1/1/1	-
5	Y70	B	502	-	-	0/4/33/33	0/3/3/3
3	EDO	A	1205	-	-	1/1/1/1	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	502	Y70	C14-C12-N8	-2.13	108.68	113.85
5	B	502	Y70	C15-C12-N8	2.05	110.95	109.08

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1206	EDO	O1-C1-C2-O2
3	A	1203	EDO	O1-C1-C2-O2
3	A	1205	EDO	O1-C1-C2-O2
3	A	1201	EDO	O1-C1-C2-O2


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1205	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1116/1148 (97%)	0.17	50 (4%) 33 36	30, 61, 116, 129	3 (0%)
2	B	379/407 (93%)	0.12	14 (3%) 41 45	45, 67, 97, 131	0
All	All	1495/1555 (96%)	0.16	64 (4%) 35 38	30, 63, 112, 131	3 (0%)

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	984	THR	5.4
1	A	596	PHE	5.1
2	B	440	LEU	4.9
1	A	745	THR	4.9
1	A	582	LEU	4.8
1	A	614	PHE	4.8
1	A	518	TYR	4.5
1	A	539	ALA	4.5
1	A	572	PRO	4.4
1	A	612	PHE	4.2
1	A	566	ALA	4.2
2	B	44	PRO	4.0
1	A	561	TRP	3.8
1	A	571	LEU	3.8
1	A	370	GLN	3.7
2	B	195	SER	3.6
1	A	573	SER	3.6
1	A	338	VAL	3.5
1	A	746	SER	3.3
1	A	487	VAL	3.3
1	A	550	ASN	3.2
2	B	272	SER	3.1
1	A	668	PHE	3.1
1	A	613	TYR	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	624	SER	3.0
1	A	493	PRO	3.0
2	B	427	PRO	3.0
1	A	517	TYR	3.0
1	A	607	GLY	2.9
1	A	601	TYR	2.8
1	A	536	HIS	2.8
2	B	196	ALA	2.7
1	A	490	TRP	2.7
2	B	264	TRP	2.6
1	A	747	GLY	2.5
1	A	562	THR	2.5
1	A	563	ASP	2.5
1	A	512	VAL	2.5
1	A	1012	LEU	2.4
1	A	288	GLU	2.4
2	B	68	HIS	2.4
1	A	583	GLY	2.4
1	A	554	PRO	2.3
1	A	1033	VAL	2.3
2	B	267	ASN	2.3
1	A	576	LEU	2.3
1	A	635	PRO	2.2
2	B	197	VAL	2.2
1	A	534	MET	2.2
1	A	95	GLY	2.2
1	A	981	SER	2.2
2	B	194	MET	2.2
2	B	206	GLN	2.2
1	A	530	SER	2.2
1	A	431	GLY	2.2
1	A	476	VAL	2.2
1	A	708	GLN	2.2
1	A	575	GLU	2.1
2	B	273	LEU	2.1
1	A	636	THR	2.1
2	B	265	ASP	2.1
1	A	633	THR	2.1
1	A	368	GLU	2.0
1	A	666	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)


There are no monosaccharides in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	EDO	A	1204	4/4	0.70	0.30	76,76,77,77	0
3	EDO	A	1205	4/4	0.81	0.26	71,71,71,71	0
3	EDO	A	1206	4/4	0.87	0.25	65,65,65,65	0
3	EDO	A	1201	4/4	0.89	0.25	56,57,58,58	0
3	EDO	A	1202	4/4	0.94	0.16	47,47,48,48	0
3	EDO	A	1203	4/4	0.95	0.15	59,59,59,59	0
5	Y70	B	502	20/20	0.95	0.14	48,51,51,51	0
4	ZN	B	501	1/1	0.99	0.14	62,62,62,62	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.