



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 13, 2024 – 12:28 am BST

PDB ID : 1OJ9
Title : HUMAN MONOAMINE OXIDASE B IN COMPLEX WITH 1,4-DIPHENYL
L-2-BUTENE
Authors : Binda, C.; Edmondson, D.E.; Mattevi, A.
Deposited on : 2003-07-08
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5963 (2.30-2.30)
Clashscore	180529	6698 (2.30-2.30)
Ramachandran outliers	177936	6640 (2.30-2.30)
Sidechain outliers	177891	6640 (2.30-2.30)
RSRZ outliers	164620	5963 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	520	
1	B	520	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 8279 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called AMINE OXIDASE [FLAVIN-CONTAINING] B.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	499	Total 3971	C 2538	N 681	O 728	S 24	0	0	0
1	B	494	Total 3940	C 2519	N 676	O 721	S 24	0	0	0

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total 53	C 27	N 9	O 15	P 2	0	0
2	B	1	Total 53	C 27	N 9	O 15	P 2	0	0

- Molecule 3 is 1,4-DIPHENYL-2-BUTENE (three-letter code: 1PB) (formula: $C_{16}H_{16}$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C 16 16	0	0
3	B	1	Total C 16 16	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	100	Total O 100 100	0	0
4	B	130	Total O 130 130	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: AMINE OXIDASE [FLAVIN-CONTAINING] B



- Molecule 1: AMINE OXIDASE [FLAVIN-CONTAINING] B



4 Data and refinement statistics i

Property	Value	Source
Space group	C 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	132.10Å 224.02Å 86.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 2.30 40.00 – 2.30	Depositor EDS
% Data completeness (in resolution range)	90.6 (40.00-2.30) 90.2 (40.00-2.30)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.88 (at 2.29Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.212 , 0.253 0.206 , 0.243	Depositor DCC
R_{free} test set	1332 reflections (2.57%)	wwPDB-VP
Wilson B-factor (Å ²)	19.8	Xtrriage
Anisotropy	0.583	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 52.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.003 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.015 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	8279	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 1PB, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.41	0/4068	0.64	4/5522 (0.1%)
1	B	0.41	0/4037	0.65	6/5479 (0.1%)
All	All	0.41	0/8105	0.64	10/11001 (0.1%)

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	6	ASP	CB-CG-OD2	5.47	123.23	118.30
1	A	55	ASP	CB-CG-OD2	5.43	123.19	118.30
1	B	153	ASP	CB-CG-OD2	5.27	123.05	118.30
1	B	123	ASP	CB-CG-OD2	5.15	122.94	118.30
1	A	330	ASP	CB-CG-OD2	5.15	122.93	118.30
1	B	223	ASP	CB-CG-OD2	5.15	122.93	118.30
1	A	471	ASP	CB-CG-OD2	5.08	122.87	118.30
1	A	6	ASP	CB-CG-OD2	5.06	122.86	118.30
1	B	227	ASP	CB-CG-OD2	5.03	122.82	118.30
1	B	310	ASP	CB-CG-OD2	5.01	122.81	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3971	0	3967	21	0
1	B	3940	0	3937	22	0
2	A	53	0	29	0	0
2	B	53	0	29	0	0
3	A	16	0	16	0	0
3	B	16	0	16	0	0
4	A	100	0	0	1	0
4	B	130	0	0	0	0
All	All	8279	0	7994	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:117:ASN:HD22	1:B:120:ARG:HH21	1.31	0.77
1:A:353:ALA:HB1	1:B:280:MET:HE1	1.71	0.71
1:A:117:ASN:HD22	1:A:120:ARG:HH21	1.37	0.71
1:A:280:MET:HE1	1:B:353:ALA:HB1	1.77	0.66
1:A:28:LEU:HD11	1:A:454:MET:HE1	1.88	0.55
1:A:3:ASN:HB3	1:A:4:LYS:HG2	1.89	0.55
1:B:3:ASN:HB3	1:B:4:LYS:HG2	1.90	0.54
1:A:389:CYS:HB2	1:B:280:MET:HG3	1.93	0.50
1:B:117:ASN:HD22	1:B:120:ARG:NH2	2.04	0.50
1:B:445:ARG:HD2	1:B:463:TRP:CZ2	2.48	0.48
1:A:91:HIS:HE1	4:A:2067:HOH:O	1.96	0.48
1:A:445:ARG:HD2	1:A:463:TRP:CZ2	2.50	0.47
1:B:278:LEU:HD13	1:B:282:ARG:HG2	1.98	0.46
1:A:117:ASN:HD22	1:A:120:ARG:NH2	2.09	0.45
1:B:28:LEU:HD11	1:B:454:MET:CE	2.46	0.45
1:A:28:LEU:HD11	1:A:454:MET:CE	2.46	0.45
1:A:278:LEU:HD13	1:A:282:ARG:HG2	1.98	0.45
1:A:116:ASN:HD21	1:A:483:GLU:HG2	1.82	0.45
1:B:308:LYS:HB2	1:B:308:LYS:HE2	1.85	0.45
1:A:445:ARG:HD2	1:A:463:TRP:CH2	2.52	0.44
1:B:28:LEU:HD21	1:B:456:LYS:HE3	1.99	0.44
1:A:28:LEU:HD21	1:A:456:LYS:HE3	2.00	0.44
1:B:28:LEU:HD11	1:B:454:MET:HE1	2.00	0.43
1:B:281:MET:HB3	1:B:413:VAL:HG21	2.00	0.43
1:A:108:ASN:HA	1:A:109:PRO:HD3	1.89	0.43
1:A:280:MET:HG3	1:B:389:CYS:HB2	2.01	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:249:THR:CG2	1:A:253:GLU:HB2	2.49	0.43
1:B:445:ARG:HD2	1:B:463:TRP:CH2	2.53	0.42
1:A:489:VAL:N	1:A:490:PRO:HD2	2.34	0.42
1:A:23:LEU:HD22	1:A:28:LEU:HD12	2.02	0.42
1:A:451:LEU:HA	1:A:454:MET:HE2	2.00	0.42
1:B:55:ASP:O	1:B:296:LYS:HE2	2.20	0.41
1:B:416:GLN:HA	1:B:417:PRO:HD3	1.93	0.41
1:A:237:TYR:HB3	1:A:248:GLU:HB3	2.02	0.41
1:B:237:TYR:HB3	1:B:248:GLU:HB3	2.02	0.41
1:B:489:VAL:N	1:B:490:PRO:HD2	2.35	0.41
1:B:108:ASN:HA	1:B:109:PRO:HD3	1.87	0.41
1:B:238:ILE:HG21	1:B:421:ILE:HD11	2.02	0.41
1:B:249:THR:CG2	1:B:253:GLU:HB2	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	497/520 (96%)	483 (97%)	14 (3%)	0	100	100
1	B	492/520 (95%)	481 (98%)	11 (2%)	0	100	100
All	All	989/1040 (95%)	964 (98%)	25 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	427/444 (96%)	413 (97%)	14 (3%)	33	48
1	B	424/444 (96%)	410 (97%)	14 (3%)	33	48
All	All	851/888 (96%)	823 (97%)	28 (3%)	33	48

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	ASN
1	A	49	GLN
1	A	209	LYS
1	A	242	ARG
1	A	249	THR
1	A	280	MET
1	A	318	ASP
1	A	321	GLU
1	A	350	ARG
1	A	379	GLU
1	A	397	CYS
1	A	445	ARG
1	A	464	GLN
1	A	498	LEU
1	B	3	ASN
1	B	49	GLN
1	B	190	LYS
1	B	209	LYS
1	B	242	ARG
1	B	249	THR
1	B	280	MET
1	B	318	ASP
1	B	321	GLU
1	B	350	ARG
1	B	379	GLU
1	B	397	CYS
1	B	445	ARG
1	B	464	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	116	ASN
1	A	117	ASN
1	A	170	ASN
1	A	431	HIS
1	B	116	ASN
1	B	117	ASN
1	B	170	ASN
1	B	431	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	1PB	B	1498	-	17,17,17	1.09	2 (11%)	20,20,20	1.29	3 (15%)
2	FAD	A	1502	1	53,58,58	1.13	4 (7%)	68,89,89	1.51	9 (13%)
3	1PB	A	1503	-	17,17,17	1.06	2 (11%)	20,20,20	1.32	3 (15%)
2	FAD	B	1497	1	53,58,58	1.17	4 (7%)	68,89,89	1.45	9 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	1PB	B	1498	-	-	5/7/7/7	0/2/2/2
2	FAD	A	1502	1	-	1/30/50/50	0/6/6/6
3	1PB	A	1503	-	-	5/7/7/7	0/2/2/2
2	FAD	B	1497	1	-	1/30/50/50	0/6/6/6

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1497	FAD	C2A-N3A	4.14	1.38	1.32
2	A	1502	FAD	C4X-N5	4.00	1.38	1.30
2	B	1497	FAD	C4X-N5	3.95	1.38	1.30
2	A	1502	FAD	C2A-N3A	3.76	1.38	1.32
2	A	1502	FAD	C10-N1	3.02	1.39	1.33
2	B	1497	FAD	C10-N1	3.02	1.39	1.33
2	A	1502	FAD	C2A-N1A	2.57	1.38	1.33
3	B	1498	1PB	CD-CC	-2.57	1.39	1.52
3	A	1503	1PB	CD-CC	-2.55	1.39	1.52
3	A	1503	1PB	CA-CB	2.53	1.39	1.31
3	B	1498	1PB	CA-CB	2.51	1.39	1.31
2	B	1497	FAD	C2A-N1A	2.43	1.38	1.33

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1502	FAD	N3A-C2A-N1A	-5.71	119.75	128.68
2	B	1497	FAD	N3A-C2A-N1A	-5.68	119.80	128.68
2	A	1502	FAD	C4X-C10-N1	-4.19	115.02	124.73
2	B	1497	FAD	C4X-C10-N1	-3.90	115.67	124.73
2	A	1502	FAD	O4-C4-C4X	-3.68	116.84	126.60
2	B	1497	FAD	O4-C4-C4X	-3.46	117.42	126.60
3	A	1503	1PB	CC-CD-C1	3.42	121.89	113.18
3	B	1498	1PB	CC-CD-C1	3.41	121.85	113.18
2	A	1502	FAD	C9A-C5X-N5	-3.32	118.82	122.43
3	A	1503	1PB	CC-CB-CA	-3.22	119.39	125.45
3	B	1498	1PB	CC-CB-CA	-3.19	119.45	125.45
2	B	1497	FAD	C9A-C5X-N5	-3.07	119.10	122.43
2	A	1502	FAD	C4-N3-C2	-3.00	120.11	125.64
2	B	1497	FAD	C4-N3-C2	-2.89	120.30	125.64
2	A	1502	FAD	P-O3P-PA	-2.82	123.16	132.83

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1502	FAD	C10-C4X-N5	-2.78	118.95	124.86
2	B	1497	FAD	P-O3P-PA	-2.76	123.37	132.83
2	B	1497	FAD	C10-C4X-N5	-2.73	119.06	124.86
3	A	1503	1PB	CG-CA-CB	-2.36	121.08	126.99
2	B	1497	FAD	C4X-C4-N3	2.33	119.10	113.19
2	A	1502	FAD	C4X-C4-N3	2.30	119.03	113.19
3	B	1498	1PB	CG-CA-CB	-2.18	121.53	126.99
2	A	1502	FAD	C4-C4X-C10	2.18	120.46	116.79
2	B	1497	FAD	C10-N1-C2	2.12	121.14	116.90

There are no chirality outliers.


All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	1498	1PB	CB-CA-CG-CL
3	B	1498	1PB	CB-CA-CG-CH
3	A	1503	1PB	CB-CA-CG-CL
3	A	1503	1PB	CB-CA-CG-CH
3	B	1498	1PB	CA-CB-CC-CD
2	A	1502	FAD	O4B-C4B-C5B-O5B
2	B	1497	FAD	O4B-C4B-C5B-O5B
3	A	1503	1PB	CN-C1-CD-CC
3	B	1498	1PB	CN-C1-CD-CC
3	A	1503	1PB	CA-CB-CC-CD
3	A	1503	1PB	CR-C1-CD-CC
3	B	1498	1PB	CR-C1-CD-CC

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	499/520 (95%)	0.68	55 (11%) 12 13	30, 42, 58, 82	0
1	B	494/520 (95%)	0.68	45 (9%) 16 18	30, 41, 56, 73	0
All	All	993/1040 (95%)	0.68	100 (10%) 14 15	30, 42, 57, 82	0

All (100) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	501	ILE	7.4
1	B	496	ILE	6.2
1	A	498	LEU	5.2
1	B	107	TRP	4.9
1	A	500	THR	4.7
1	B	163	GLN	4.3
1	B	460	ASP	4.1
1	B	354	ARG	3.8
1	A	460	ASP	3.7
1	A	497	GLY	3.7
1	A	93	LYS	3.7
1	A	163	GLN	3.6
1	B	3	ASN	3.6
1	A	95	LYS	3.6
1	B	92	VAL	3.6
1	A	495	LEU	3.6
1	B	495	LEU	3.6
1	A	49	GLN	3.5
1	A	321	GLU	3.5
1	B	318	ASP	3.5
1	A	107	TRP	3.4
1	A	481	PHE	3.4
1	B	49	GLN	3.3
1	B	493	LEU	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	3	ASN	3.3
1	A	499	THR	3.3
1	A	308	LYS	3.2
1	B	321	GLU	3.2
1	A	318	ASP	3.1
1	B	479	THR	3.1
1	A	494	ARG	3.0
1	B	106	VAL	2.9
1	A	242	ARG	2.9
1	B	476	PRO	2.9
1	B	141	GLU	2.9
1	B	242	ARG	2.8
1	B	494	ARG	2.8
1	A	470	VAL	2.7
1	B	224	LEU	2.7
1	B	243	GLU	2.7
1	B	455	GLY	2.6
1	A	224	LEU	2.6
1	A	29	ASN	2.6
1	B	26	SER	2.6
1	B	227	ASP	2.6
1	A	320	GLU	2.6
1	B	249	THR	2.6
1	A	370	LYS	2.6
1	B	93	LYS	2.5
1	A	468	GLU	2.5
1	A	319	GLY	2.5
1	B	246	LEU	2.5
1	B	226	GLY	2.5
1	B	414	LEU	2.5
1	A	412	ARG	2.5
1	A	302	LYS	2.5
1	A	243	GLU	2.4
1	B	370	LYS	2.4
1	B	319	GLY	2.4
1	A	492	LEU	2.4
1	B	482	LEU	2.4
1	A	483	GLU	2.4
1	A	474	ALA	2.4
1	A	280	MET	2.3
1	B	254	MET	2.3
1	A	477	ILE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	226	GLY	2.3
1	A	493	LEU	2.3
1	B	374	SER	2.3
1	A	154	LYS	2.3
1	A	254	MET	2.3
1	A	128	GLU	2.2
1	A	227	ASP	2.2
1	A	109	PRO	2.2
1	B	103	PHE	2.2
1	A	475	GLN	2.2
1	B	223	ASP	2.2
1	A	101	GLY	2.2
1	A	482	LEU	2.2
1	A	480	THR	2.2
1	B	27	GLY	2.2
1	B	303	GLU	2.2
1	A	354	ARG	2.2
1	A	484	ARG	2.2
1	B	485	HIS	2.2
1	B	475	GLN	2.1
1	B	481	PHE	2.1
1	A	141	GLU	2.1
1	A	104	PRO	2.1
1	B	52	LYS	2.1
1	B	302	LYS	2.1
1	A	249	THR	2.1
1	B	478	THR	2.1
1	A	159	GLU	2.1
1	A	478	THR	2.1
1	A	252	HIS	2.0
1	A	251	ASN	2.0
1	B	480	THR	2.0
1	B	483	GLU	2.0
1	A	463	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	1PB	A	1503	16/16	0.86	0.18	59,60,61,61	0
3	1PB	B	1498	16/16	0.89	0.18	59,60,61,61	0
2	FAD	A	1502	53/53	0.97	0.08	36,38,45,46	0
2	FAD	B	1497	53/53	0.97	0.08	36,38,45,46	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.