



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 29, 2023 – 12:30 AM EDT

PDB ID : 3OL7
Title : Poliovirus polymerase elongation complex with CTP
Authors : Gong, P.; Peersen, O.B.
Deposited on : 2010-08-25
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)
RNA backbone	3102	1159 (3.00-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	471	
1	E	471	
1	I	471	
1	M	471	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	B	26	
2	F	26	
2	J	26	
2	N	26	
3	C	15	
3	G	15	
3	K	15	
3	O	15	
4	D	9	
4	H	9	
4	L	9	
4	P	9	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
10	GOL	I	8002	-	-	X	-
10	GOL	I	8003	-	-	X	-
10	GOL	I	8004	-	-	X	-
10	GOL	I	8006	-	-	X	-
7	IPA	E	6014	-	-	X	-
7	IPA	I	6023	-	-	-	X
7	IPA	M	6016	-	-	X	-
7	IPA	M	6019	-	-	X	-
8	PEG	A	7001	-	-	X	-
8	PEG	E	7002	-	-	X	-

2 Entry composition [i](#)

There are 11 unique types of molecules in this entry. The entry contains 18525 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	461	3697	2370	610	695	22	0	0	0
1	E	461	3697	2370	610	695	22	0	0	0
1	I	461	3697	2370	610	695	22	0	0	0
1	M	461	3697	2370	610	695	22	0	0	0

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	446	ASP	LEU	engineered mutation	UNP B3VQP5
A	462	GLY	-	expression tag	UNP B3VQP5
A	463	SER	-	expression tag	UNP B3VQP5
A	464	SER	-	expression tag	UNP B3VQP5
A	465	SER	-	expression tag	UNP B3VQP5
A	466	HIS	-	expression tag	UNP B3VQP5
A	467	HIS	-	expression tag	UNP B3VQP5
A	468	HIS	-	expression tag	UNP B3VQP5
A	469	HIS	-	expression tag	UNP B3VQP5
A	470	HIS	-	expression tag	UNP B3VQP5
A	471	HIS	-	expression tag	UNP B3VQP5
E	446	ASP	LEU	engineered mutation	UNP B3VQP5
E	462	GLY	-	expression tag	UNP B3VQP5
E	463	SER	-	expression tag	UNP B3VQP5
E	464	SER	-	expression tag	UNP B3VQP5
E	465	SER	-	expression tag	UNP B3VQP5
E	466	HIS	-	expression tag	UNP B3VQP5
E	467	HIS	-	expression tag	UNP B3VQP5
E	468	HIS	-	expression tag	UNP B3VQP5
E	469	HIS	-	expression tag	UNP B3VQP5
E	470	HIS	-	expression tag	UNP B3VQP5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	471	HIS	-	expression tag	UNP B3VQP5
I	446	ASP	LEU	engineered mutation	UNP B3VQP5
I	462	GLY	-	expression tag	UNP B3VQP5
I	463	SER	-	expression tag	UNP B3VQP5
I	464	SER	-	expression tag	UNP B3VQP5
I	465	SER	-	expression tag	UNP B3VQP5
I	466	HIS	-	expression tag	UNP B3VQP5
I	467	HIS	-	expression tag	UNP B3VQP5
I	468	HIS	-	expression tag	UNP B3VQP5
I	469	HIS	-	expression tag	UNP B3VQP5
I	470	HIS	-	expression tag	UNP B3VQP5
I	471	HIS	-	expression tag	UNP B3VQP5
M	446	ASP	LEU	engineered mutation	UNP B3VQP5
M	462	GLY	-	expression tag	UNP B3VQP5
M	463	SER	-	expression tag	UNP B3VQP5
M	464	SER	-	expression tag	UNP B3VQP5
M	465	SER	-	expression tag	UNP B3VQP5
M	466	HIS	-	expression tag	UNP B3VQP5
M	467	HIS	-	expression tag	UNP B3VQP5
M	468	HIS	-	expression tag	UNP B3VQP5
M	469	HIS	-	expression tag	UNP B3VQP5
M	470	HIS	-	expression tag	UNP B3VQP5
M	471	HIS	-	expression tag	UNP B3VQP5

- Molecule 2 is a RNA chain called RNA (5'-R(*AP*AP*GP*UP*CP*UP*CP*CP*AP*GP*GP*UP*CP*UP*CP*UP*CP*UP*CP*GP*UP*CP*CP*GP*GP*AP*AP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	B	18	Total 361	C 159	N 59	O 125	P 18	0	0	0
2	F	18	Total 361	C 159	N 59	O 125	P 18	0	0	0
2	J	16	Total 337	C 150	N 56	O 115	P 16	0	0	0
2	N	16	Total 337	C 150	N 56	O 115	P 16	0	0	0

- Molecule 3 is a RNA chain called RNA (5'-R(*GP*CP*CP*CP*GP*GP*AP*CP*GP*AP*GP*AP*GP*AP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	15	Total	C	N	O	P	0	0	0
			323	145	65	99	14			
3	G	15	Total	C	N	O	P	0	0	0
			323	145	65	99	14			
3	K	15	Total	C	N	O	P	0	0	0
			323	145	65	99	14			
3	O	15	Total	C	N	O	P	0	0	0
			323	145	65	99	14			


- Molecule 4 is a RNA chain called RNA (5'-R(*GP*GP*GP*AP*GP*AP*UP*GP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	3	Total	C	N	O	P	0	0	0
			68	30	15	20	3			
4	H	3	Total	C	N	O	P	0	0	0
			68	30	15	20	3			
4	L	4	Total	C	N	O	P	0	0	0
			91	40	20	27	4			
4	P	4	Total	C	N	O	P	0	0	0
			91	40	20	27	4			

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Mg	0	0
			2	2		
5	E	2	Total	Mg	0	0
			2	2		
5	I	1	Total	Mg	0	0
			1	1		
5	M	1	Total	Mg	0	0
			1	1		

- Molecule 6 is PYROPHOSPHATE 2- (three-letter code: POP) (formula: H₂O₇P₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total O P 9 7 2	0	0
6	E	1	Total O P 9 7 2	0	0
6	K	1	Total O P 9 7 2	0	0
6	M	1	Total O P 9 7 2	0	0

- Molecule 7 is ISOPROPYL ALCOHOL (three-letter code: IPA) (formula: C₃H₈O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			4	3	1		
7	A	1	Total	C	O	0	0
			4	3	1		
7	A	1	Total	C	O	0	0
			4	3	1		
7	E	1	Total	C	O	0	0
			4	3	1		
7	E	1	Total	C	O	0	0
			4	3	1		
7	E	1	Total	C	O	0	0
			4	3	1		
7	E	1	Total	C	O	0	0
			4	3	1		
7	I	1	Total	C	O	0	0
			4	3	1		
7	I	1	Total	C	O	0	0
			4	3	1		
7	M	1	Total	C	O	0	0
			4	3	1		
7	M	1	Total	C	O	0	0
			4	3	1		
7	M	1	Total	C	O	0	0
			4	3	1		
7	N	1	Total	C	O	0	0
			4	3	1		
7	O	1	Total	C	O	0	0
			4	3	1		

- Molecule 8 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total C O 7 4 3	0	0
8	E	1	Total C O 7 4 3	0	0
8	J	1	Total C O 7 4 3	0	0
8	N	1	Total C O 7 4 3	0	0

- Molecule 9 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	1	Total Zn 1 1	0	0
9	E	1	Total Zn 1 1	0	0
9	I	1	Total Zn 1 1	0	0
9	M	1	Total Zn 1 1	0	0

- Molecule 10 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	E	1	Total C O 6 3 3	0	0
10	I	1	Total C O 6 3 3	0	0
10	I	1	Total C O 6 3 3	0	0
10	I	1	Total C O 6 3 3	0	0
10	I	1	Total C O 6 3 3	0	0
10	J	1	Total C O 6 3 3	0	0
10	M	1	Total C O 6 3 3	0	0
10	N	1	Total C O 6 3 3	0	0

- Molecule 11 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	A	122	Total O 122 122	0	0
11	B	19	Total O 19 19	0	0
11	C	14	Total O 14 14	0	0
11	D	1	Total O 1 1	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	E	100	Total 100	O 100	0	0
11	F	25	Total 25	O 25	0	0
11	G	9	Total 9	O 9	0	0
11	H	1	Total 1	O 1	0	0
11	I	95	Total 95	O 95	0	0
11	J	19	Total 19	O 19	0	0
11	K	11	Total 11	O 11	0	0
11	M	105	Total 105	O 105	0	0
11	N	21	Total 21	O 21	0	0
11	O	7	Total 7	O 7	0	0


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Polymerase




- Molecule 1: Polymerase







• Molecule 1: Polymerase



• Molecule 1: Polymerase







● Molecule 2: RNA (5'-R(*AP*AP*GP*UP*CP*UP*CP*CP*AP*GP*GP*UP*CP*UP*CP*UP*CP*GP*UP*CP*CP*GP*GP*AP*AP*A)-3')




● Molecule 2: RNA (5'-R(*AP*AP*GP*UP*CP*UP*CP*CP*AP*GP*GP*UP*CP*UP*CP*UP*CP*GP*UP*CP*CP*GP*GP*AP*AP*A)-3')




● Molecule 2: RNA (5'-R(*AP*AP*GP*UP*CP*UP*CP*CP*AP*GP*GP*UP*CP*UP*CP*UP*CP*GP*UP*CP*CP*GP*GP*AP*AP*A)-3')




● Molecule 2: RNA (5'-R(*AP*AP*GP*UP*CP*UP*CP*CP*AP*GP*GP*UP*CP*UP*CP*UP*CP*GP*UP*CP*CP*GP*GP*AP*AP*A)-3')




● Molecule 3: RNA (5'-R(*GP*CP*CP*CP*GP*GP*AP*CP*GP*AP*GP*AP*GP*AP*C)-3')




- Molecule 3: RNA (5'-R(*GP*CP*CP*CP*GP*GP*AP*CP*GP*AP*GP*AP*GP*AP*C)-3')




- Molecule 3: RNA (5'-R(*GP*CP*CP*CP*GP*GP*AP*CP*GP*AP*GP*AP*GP*AP*C)-3')




- Molecule 3: RNA (5'-R(*GP*CP*CP*CP*GP*GP*AP*CP*GP*AP*GP*AP*GP*AP*C)-3')




- Molecule 4: RNA (5'-R(*GP*GP*GP*AP*GP*AP*UP*GP*A)-3')




- Molecule 4: RNA (5'-R(*GP*GP*GP*AP*GP*AP*UP*GP*A)-3')



- Molecule 4: RNA (5'-R(*GP*GP*GP*AP*GP*AP*UP*GP*A)-3')



- Molecule 4: RNA (5'-R(*GP*GP*GP*AP*GP*AP*UP*GP*A)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	60.72Å 60.77Å 192.97Å 83.01° 82.93° 77.12°	Depositor
Resolution (Å)	48.08 – 2.70 48.09 – 2.40	Depositor EDS
% Data completeness (in resolution range)	97.5 (48.08-2.70) 97.4 (48.09-2.40)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.37 (at 2.39Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.6.4_486)	Depositor
R, R_{free}	0.226 , 0.290 0.218 , 0.281	Depositor DCC
R_{free} test set	5054 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	60.8	Xtrriage
Anisotropy	0.445	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 40.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.61$, $\langle L^2 \rangle = 0.46$	Xtrriage
Estimated twinning fraction	0.448 for -k,-h,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	18525	wwPDB-VP
Average B, all atoms (Å ²)	69.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 43.74 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.6883e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, IPA, POP, ZN, MG, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/3787	0.60	0/5122
1	E	0.45	0/3787	0.60	0/5122
1	I	0.43	0/3787	0.62	0/5122
1	M	0.44	0/3787	0.62	1/5122 (0.0%)
2	B	0.79	2/400 (0.5%)	1.21	2/621 (0.3%)
2	F	0.79	2/400 (0.5%)	1.20	3/621 (0.5%)
2	J	0.62	0/374	1.01	0/580
2	N	0.63	0/374	1.01	0/580
3	C	0.66	0/362	1.19	0/564
3	G	0.68	0/362	1.15	0/564
3	K	0.64	0/362	1.01	0/564
3	O	0.66	0/362	0.99	0/564
4	D	0.28	0/76	0.54	0/117
4	H	0.31	0/76	0.54	0/117
4	L	0.29	0/102	0.62	0/158
4	P	0.32	0/102	0.63	0/158
All	All	0.49	4/18500 (0.0%)	0.73	6/25696 (0.0%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	613	A	P-OP1	6.71	1.60	1.49
2	F	613	A	P-OP1	6.70	1.60	1.49
2	F	613	A	P-OP2	6.36	1.59	1.49
2	B	613	A	P-OP2	6.31	1.59	1.49

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	599	G	C4-N9-C1'	5.72	133.94	126.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	599	G	C4-N9-C1'	5.66	133.86	126.50
2	B	597	C	C2-N1-C1'	5.48	124.83	118.80
2	F	597	C	C2-N1-C1'	5.41	124.76	118.80
1	M	45	LYS	CD-CE-NZ	5.25	123.77	111.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3697	0	3659	161	0
1	E	3697	0	3659	177	0
1	I	3697	0	3658	177	0
1	M	3697	0	3658	167	0
2	B	361	0	183	34	0
2	F	361	0	183	33	0
2	J	337	0	173	17	0
2	N	337	0	173	19	0
3	C	323	0	167	10	0
3	G	323	0	167	9	0
3	K	323	0	167	9	0
3	O	323	0	167	10	0
4	D	68	0	34	3	0
4	H	68	0	34	2	0
4	L	91	0	45	2	0
4	P	91	0	45	2	0
5	A	2	0	0	0	0
5	E	2	0	0	0	0
5	I	1	0	0	0	0
5	M	1	0	0	0	0
6	A	9	0	0	0	0
6	E	9	0	0	0	0
6	K	9	0	0	0	0
6	M	9	0	0	0	0
7	A	16	0	32	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	E	16	0	32	16	0
7	I	8	0	16	2	0
7	M	12	0	24	9	0
7	N	4	0	8	2	0
7	O	4	0	8	0	0
8	A	7	0	10	7	0
8	E	7	0	10	9	0
8	J	7	0	10	0	0
8	N	7	0	10	1	0
9	A	1	0	0	0	0
9	E	1	0	0	0	0
9	I	1	0	0	0	0
9	M	1	0	0	0	0
10	E	6	0	8	0	0
10	I	24	0	32	27	0
10	J	6	0	8	3	0
10	M	6	0	8	2	0
10	N	6	0	8	3	0
11	A	122	0	0	12	0
11	B	19	0	0	1	0
11	C	14	0	0	0	0
11	D	1	0	0	0	0
11	E	100	0	0	6	0
11	F	25	0	0	2	0
11	G	9	0	0	0	0
11	H	1	0	0	1	0
11	I	95	0	0	6	0
11	J	19	0	0	2	0
11	K	11	0	0	0	0
11	M	105	0	0	16	0
11	N	21	0	0	2	0
11	O	7	0	0	1	0
All	All	18525	0	16396	783	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

The worst 5 of 783 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:404:THR:HB	8:E:7002:PEG:H32	1.35	1.09

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:N:6020:IPA:H31	8:N:7004:PEG:H42	1.33	1.06
1:A:410:THR:HA	8:A:7001:PEG:H41	1.35	1.05
1:I:337:GLU:HG2	10:I:8003:GOL:H31	1.41	1.02
1:A:404:THR:HB	8:A:7001:PEG:H32	1.38	1.01

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	459/471 (98%)	426 (93%)	29 (6%)	4 (1%)	17	40
1	E	459/471 (98%)	425 (93%)	30 (6%)	4 (1%)	17	40
1	I	459/471 (98%)	400 (87%)	51 (11%)	8 (2%)	9	23
1	M	459/471 (98%)	396 (86%)	56 (12%)	7 (2%)	10	26
All	All	1836/1884 (98%)	1647 (90%)	166 (9%)	23 (1%)	12	30

5 of 23 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	66	LYS
1	E	66	LYS
1	I	6	MET
1	I	12	VAL
1	M	6	MET

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	403/412 (98%)	372 (92%)	31 (8%)	13	30
1	E	403/412 (98%)	372 (92%)	31 (8%)	13	30
1	I	403/412 (98%)	370 (92%)	33 (8%)	11	26
1	M	403/412 (98%)	370 (92%)	33 (8%)	11	26
All	All	1612/1648 (98%)	1484 (92%)	128 (8%)	12	28

5 of 128 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	M	281	CYS
1	M	352	LEU
1	E	218	TRP
1	E	212	CYS
1	M	366	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
1	I	398	HIS
1	M	4	GLN
1	I	424	ASN
1	M	170	GLN
1	E	170	GLN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	B	16/26 (61%)	4 (25%)	1 (6%)
2	F	16/26 (61%)	4 (25%)	1 (6%)
2	J	15/26 (57%)	2 (13%)	1 (6%)
2	N	15/26 (57%)	2 (13%)	1 (6%)
3	C	14/15 (93%)	2 (14%)	0
3	G	14/15 (93%)	2 (14%)	0
3	K	14/15 (93%)	0	0
3	O	14/15 (93%)	0	0
4	D	2/9 (22%)	1 (50%)	0
4	H	2/9 (22%)	1 (50%)	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
4	L	3/9 (33%)	1 (33%)	0
4	P	3/9 (33%)	1 (33%)	0
All	All	128/200 (64%)	20 (15%)	4 (3%)

5 of 20 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	B	598	A
2	B	599	G
2	B	600	G
2	B	602	C
3	C	690	C

All (4) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	B	599	G
2	F	599	G
2	J	598	A
2	N	598	A

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 41 ligands modelled in this entry, 10 are monoatomic - leaving 31 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	IPA	E	6014	-	3,3,3	0.40	0	3,3,3	0.76	0
7	IPA	N	6020	-	3,3,3	0.63	0	3,3,3	0.16	0
6	POP	M	5001	5	6,8,8	0.73	0	13,13,13	1.40	3 (23%)
7	IPA	E	6013	-	3,3,3	0.59	0	3,3,3	0.19	0
10	GOL	I	8003	-	5,5,5	0.46	0	5,5,5	0.40	0
7	IPA	M	6016	-	3,3,3	0.60	0	3,3,3	0.31	0
10	GOL	E	8008	-	5,5,5	0.39	0	5,5,5	0.35	0
7	IPA	A	6025	-	3,3,3	0.59	0	3,3,3	0.24	0
7	IPA	M	6007	-	3,3,3	0.60	0	3,3,3	0.25	0
7	IPA	A	6008	-	3,3,3	0.57	0	3,3,3	0.24	0
10	GOL	M	8009	-	5,5,5	0.41	0	5,5,5	0.56	0
6	POP	E	5003	5	6,8,8	0.72	0	13,13,13	1.58	2 (15%)
8	PEG	E	7002	-	6,6,6	0.45	0	5,5,5	0.33	0
10	GOL	I	8006	-	5,5,5	0.35	0	5,5,5	0.58	0
7	IPA	E	6015	-	3,3,3	0.61	0	3,3,3	0.33	0
6	POP	K	5002	5	6,8,8	0.65	0	13,13,13	1.34	1 (7%)
7	IPA	M	6019	-	3,3,3	0.55	0	3,3,3	0.30	0
7	IPA	I	6012	-	3,3,3	0.59	0	3,3,3	0.31	0
8	PEG	J	7003	-	6,6,6	0.63	0	5,5,5	0.24	0
7	IPA	O	6021	-	3,3,3	0.65	0	3,3,3	0.13	0
7	IPA	I	6023	-	3,3,3	0.55	0	3,3,3	0.35	0
8	PEG	A	7001	-	6,6,6	0.47	0	5,5,5	0.54	0
7	IPA	A	6011	-	3,3,3	0.61	0	3,3,3	0.21	0
8	PEG	N	7004	-	6,6,6	0.56	0	5,5,5	0.29	0
7	IPA	A	6009	-	3,3,3	0.63	0	3,3,3	0.19	0
6	POP	A	5004	5	6,8,8	0.70	0	13,13,13	1.35	1 (7%)
7	IPA	E	6010	-	3,3,3	0.56	0	3,3,3	0.35	0
10	GOL	I	8002	-	5,5,5	0.30	0	5,5,5	0.55	0
10	GOL	J	8005	-	5,5,5	0.48	0	5,5,5	0.64	0
10	GOL	I	8004	-	5,5,5	0.46	0	5,5,5	0.37	0
10	GOL	N	8010	-	5,5,5	0.38	0	5,5,5	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	PEG	N	7004	-	-	1/4/4/4	-
6	POP	M	5001	5	-	0/6/6/6	-
6	POP	K	5002	5	-	0/6/6/6	-
6	POP	A	5004	5	-	0/6/6/6	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	GOL	M	8009	-	-	2/4/4/4	-
10	GOL	I	8002	-	-	3/4/4/4	-
8	PEG	J	7003	-	-	1/4/4/4	-
6	POP	E	5003	5	-	0/6/6/6	-
10	GOL	J	8005	-	-	2/4/4/4	-
10	GOL	I	8004	-	-	2/4/4/4	-
8	PEG	E	7002	-	-	2/4/4/4	-
10	GOL	I	8003	-	-	2/4/4/4	-
10	GOL	N	8010	-	-	2/4/4/4	-
10	GOL	I	8006	-	-	4/4/4/4	-
8	PEG	A	7001	-	-	2/4/4/4	-
10	GOL	E	8008	-	-	2/4/4/4	-

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	E	5003	POP	P2-O-P1	-3.97	119.22	132.83
6	K	5002	POP	P2-O-P1	-3.63	120.36	132.83
6	M	5001	POP	P2-O-P1	-3.32	121.44	132.83
6	A	5004	POP	P2-O-P1	-3.22	121.78	132.83
6	E	5003	POP	O3-P1-O	2.51	113.04	104.64

There are no chirality outliers.

5 of 25 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
10	I	8002	GOL	O1-C1-C2-C3
10	I	8004	GOL	O1-C1-C2-C3
10	I	8006	GOL	O1-C1-C2-C3
10	I	8006	GOL	C1-C2-C3-O3
10	I	8006	GOL	O2-C2-C3-O3

There are no ring outliers.

21 monomers are involved in 84 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	E	6014	IPA	10	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	N	6020	IPA	2	0
7	E	6013	IPA	1	0
10	I	8003	GOL	5	0
7	M	6016	IPA	4	0
7	A	6025	IPA	1	0
10	M	8009	GOL	2	0
8	E	7002	PEG	9	0
10	I	8006	GOL	10	0
7	E	6015	IPA	2	0
7	M	6019	IPA	5	0
7	I	6023	IPA	2	0
8	A	7001	PEG	7	0
7	A	6011	IPA	2	0
8	N	7004	PEG	1	0
7	A	6009	IPA	1	0
7	E	6010	IPA	3	0
10	I	8002	GOL	8	0
10	J	8005	GOL	3	0
10	I	8004	GOL	4	0
10	N	8010	GOL	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	461/471 (97%)	0.17	30 (6%) 18 17	33, 57, 94, 113	0
1	E	461/471 (97%)	0.12	32 (6%) 16 15	32, 57, 93, 113	0
1	I	461/471 (97%)	0.29	38 (8%) 11 9	33, 63, 114, 165	0
1	M	461/471 (97%)	0.24	34 (7%) 14 12	33, 64, 114, 166	0
2	B	18/26 (69%)	-0.00	2 (11%) 5 4	43, 63, 155, 214	0
2	F	18/26 (69%)	0.16	2 (11%) 5 4	42, 63, 154, 214	0
2	J	16/26 (61%)	-0.30	1 (6%) 20 19	45, 75, 149, 160	0
2	N	16/26 (61%)	-0.17	1 (6%) 20 19	45, 76, 150, 160	0
3	C	15/15 (100%)	0.05	1 (6%) 17 16	46, 63, 121, 129	0
3	G	15/15 (100%)	0.13	2 (13%) 3 2	46, 63, 122, 131	0
3	K	15/15 (100%)	-0.37	0 100 100	51, 69, 140, 148	0
3	O	15/15 (100%)	-0.20	1 (6%) 17 16	52, 70, 142, 150	0
4	D	3/9 (33%)	1.16	1 (33%) 0 0	142, 142, 147, 150	0
4	H	3/9 (33%)	1.40	1 (33%) 0 0	142, 142, 147, 150	0
4	L	4/9 (44%)	1.45	1 (25%) 0 0	124, 130, 130, 159	0
4	P	4/9 (44%)	1.39	2 (50%) 0 0	125, 130, 130, 159	0
All	All	1986/2084 (95%)	0.19	149 (7%) 14 12	32, 60, 113, 214	0

The worst 5 of 149 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	M	342	LEU	11.1
1	A	46	ASN	10.7
1	I	46	ASN	9.8
1	I	344	ALA	9.3
1	I	342	LEU	8.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
9	ZN	A	2001	1/1	0.28	0.40	114,114,114,114	1
7	IPA	N	6020	4/4	0.47	0.26	92,101,106,108	0
9	ZN	E	2002	1/1	0.47	0.22	111,111,111,111	1
7	IPA	A	6011	4/4	0.62	0.37	72,85,88,93	0
10	GOL	I	8002	6/6	0.65	0.26	82,122,129,131	0
7	IPA	A	6025	4/4	0.69	0.18	88,91,105,115	0
10	GOL	I	8003	6/6	0.70	0.32	94,114,116,117	0
10	GOL	E	8008	6/6	0.74	0.15	81,97,104,105	0
7	IPA	I	6023	4/4	0.74	0.50	75,81,84,91	0
7	IPA	E	6015	4/4	0.74	0.31	71,72,73,78	0
7	IPA	I	6012	4/4	0.77	0.30	89,102,107,110	0
7	IPA	E	6013	4/4	0.77	0.22	100,104,112,117	0
10	GOL	M	8009	6/6	0.78	0.15	75,82,101,108	0
7	IPA	O	6021	4/4	0.81	0.22	75,82,87,99	0
8	PEG	N	7004	7/7	0.81	0.21	79,88,98,104	0
10	GOL	I	8004	6/6	0.82	0.19	77,106,109,117	0
5	MG	E	3002	1/1	0.83	0.28	83,83,83,83	0
10	GOL	I	8006	6/6	0.83	0.13	94,98,108,120	0
7	IPA	M	6016	4/4	0.83	0.20	63,84,90,94	0
7	IPA	E	6010	4/4	0.84	0.24	60,71,76,77	0
8	PEG	J	7003	7/7	0.85	0.21	71,79,103,103	0
8	PEG	E	7002	7/7	0.85	0.18	61,67,96,99	0
7	IPA	A	6008	4/4	0.86	0.31	78,82,93,99	0
5	MG	I	3004	1/1	0.86	0.12	83,83,83,83	0
7	IPA	A	6009	4/4	0.89	0.32	52,72,73,78	0
7	IPA	M	6019	4/4	0.89	0.16	68,94,98,105	0
6	POP	A	5004	9/9	0.89	0.19	46,76,111,112	9

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
8	PEG	A	7001	7/7	0.90	0.13	58,73,84,89	0
6	POP	E	5003	9/9	0.90	0.21	36,57,112,122	9
7	IPA	M	6007	4/4	0.90	0.42	86,96,100,101	0
6	POP	M	5001	9/9	0.90	0.23	65,99,115,133	9
7	IPA	E	6014	4/4	0.91	0.24	57,61,67,76	0
6	POP	K	5002	9/9	0.91	0.14	62,94,108,117	9
5	MG	A	3006	1/1	0.91	0.34	81,81,81,81	0
10	GOL	N	8010	6/6	0.91	0.20	56,64,66,87	0
5	MG	M	3001	1/1	0.92	0.08	74,74,74,74	0
10	GOL	J	8005	6/6	0.94	0.39	52,55,71,73	0
9	ZN	M	2004	1/1	0.95	0.04	76,76,76,76	1
5	MG	A	3005	1/1	0.97	0.15	60,60,60,60	0
5	MG	E	3003	1/1	0.97	0.17	59,59,59,59	0
9	ZN	I	2003	1/1	0.98	0.11	78,78,78,78	1

6.5 Other polymers [i](#)

There are no such residues in this entry.