

wwPDB X-ray Structure Validation Summary Report (i)

May 15, 2020 - 03:35 am BST

PDB ID : 5OLH

Title: Structure of the A2A-StaR2-bRIL562-Vipadenant complex at 2.6A obtained

from in meso soaking experiments.

Authors: Rucktooa, P.; Cheng, R.K.Y.; Segala, E.; Geng, T.; Errey, J.C.; Brown, G.A.;

Cooke, R.; Marshall, F.H.; Dore, A.S.

Deposited on : 2017-07-27

Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.11

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \; resolution} \\ (\#{\rm Entries, \; resolution \; range(\AA)}) \end{array}$
R_{free}	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
			5%		
1	A	434	84%	6%	10%



2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 3559 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Adenosine receptor A2a, Soluble cytochrome b562, Adenosine receptor A2a.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	390	Total 3047	C 1986	N 515	O 524	S 22	0	3	0

There are 34 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-9	ALA	-	expression tag	UNP P29274
A	-8	ASP	_	expression tag	UNP P29274
A	-7	TYR	-	expression tag	UNP P29274
A	-6	LYS	-	expression tag	UNP P29274
A	-5	ASP	-	expression tag	UNP P29274
A	-4	ASP	=	expression tag	UNP P29274
A	-3	ASP	=	expression tag	UNP P29274
A	-2	ASP	=	expression tag	UNP P29274
A	-1	GLY	=	expression tag	UNP P29274
A	0	ALA	=	expression tag	UNP P29274
A	1	PRO	_	expression tag	UNP P29274
A	54	LEU	ALA	engineered mutation	UNP P29274
A	88	ALA	THR	engineered mutation	UNP P29274
A	107	ALA	ARG	engineered mutation	UNP P29274
A	122	ALA	LYS	engineered mutation	UNP P29274
A	154	ALA	ASN	engineered mutation	UNP P29274
A	202	ALA	LEU	engineered mutation	UNP P29274
A	1007	TRP	MET	engineered mutation	UNP P0ABE7
A	1102	ILE	HIS	engineered mutation	UNP P0ABE7
A	1106	LEU	ARG	engineered mutation	UNP P0ABE7
A	235	ALA	LEU	engineered mutation	UNP P29274
A	239	ALA	VAL	engineered mutation	UNP P29274
A	277	ALA	SER	engineered mutation	UNP P29274
A	318	ALA	=	expression tag	UNP P29274
A	319	HIS	-	expression tag	UNP P29274
A	320	HIS	ı	expression tag	UNP P29274


Continued on next page...



~ · · 1	e		
Continued	trom	previous	paae

Chain	Residue	Modelled	Actual	Comment	Reference
A	321	HIS	-	expression tag	UNP P29274
A	322	HIS	-	expression tag	UNP P29274
A	323	HIS	-	expression tag	UNP P29274
A	324	HIS	-	expression tag	UNP P29274
A	325	HIS	-	expression tag	UNP P29274
A	326	HIS	-	expression tag	UNP P29274
A	327	HIS	-	expression tag	UNP P29274
A	328	HIS	-	expression tag	UNP P29274


 $\bullet \ \, \text{Molecule 2 is 3-[(4-azanyl-3-methyl-phenyl)methyl]-7-(furan-2-yl)-[1,2,3]triazolo[4,5-d]pyrim idin-5-amine (three-letter code: 9XT) (formula: $C_{16}H_{15}N_{7}O$). }$



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	Δ	1	Total	С	N	О	0	0
	11	1	24	16	7	1	0	

 \bullet Molecule 3 is OLEIC ACID (three-letter code: OLA) (formula: $\mathrm{C_{18}H_{34}O_{2}}).$





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 20 18 2	0	0
3	A	1	Total C O 9 7 2	0	0
3	A	1	Total C O 18 16 2	0	0
3	A	1	Total C O 20 18 2	0	0
3	A	1	Total C O 20 18 2	0	0
3	A	1	Total C O 12 10 2	0	0
3	A	1	Total C O 8 6 2	0	0
3	A	1	Total C 9 9	0	0
3	A	1	Total C O 19 17 2	0	0
3	A	1	Total C O 14 12 2	0	0
3	A	1	Total C 10 10	0	0
3	A	1	Total C 13 13	0	0
3	A	1	Total C 12 12	0	0
3	A	1	Total C 13 13	0	0

Continued on next page...




Continued from previous page...

M	ol	Chain	Residues	At	oms		ZeroOcc	AltConf
ę	3	A	1	Total 20	C 18	O 2	0	0

• Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Na 1 1	0	0


• Molecule 5 is CHOLESTEROL (three-letter code: CLR) (formula: $C_{27}H_{46}O$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 28 27 1	0	0
5	A	1	Total C O 28 27 1	0	0
5	A	1	Total C O 28 27 1	0	0

• Molecule 6 is (2R)-2,3-dihydroxypropyl (9Z)-octadec-9-enoate (three-letter code: OLC) (formula: $C_{21}H_{40}O_4$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 16 12 4	0	0
6	A	1	Total C O 17 13 4	0	0
6	A	1	Total C O 17 15 2	0	0
6	A	1	Total C O 20 18 2	0	0
6	A	1	Total C O 24 20 4	0	0
6	A	1	Total C O 22 18 4	0	0
6	A	1	Total C O 20 18 2	0	0
6	A	1	Total C O 20 18 2	0	0

• Molecule 7 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	29	Total O 30 30	0	1



3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Adenosine receptor A2a, Soluble cytochrome b562, Adenosine receptor A2a





4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants	39.40Å 179.33Å 141.15Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.82 - 2.60	Depositor
resolution (11)	29.82 - 2.60	EDS
% Data completeness	97.9 (29.82-2.60)	Depositor
(in resolution range)	99.3 (29.82-2.60)	EDS
R_{merge}	0.25	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.78 (at 2.61Å)	Xtriage
Refinement program	PHENIX (1.12RC2_2821)	Depositor
R, R_{free}	0.200 , 0.249	Depositor
10, 10 free	0.194 , 0.243	DCC
R_{free} test set	815 reflections (5.15%)	wwPDB-VP
Wilson B-factor (Å ²)	31.8	Xtriage
Anisotropy	0.374	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.34 , 71.2	EDS
L-test for twinning ²	$ < L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3559	wwPDB-VP
Average B, all atoms $(Å^2)$	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 6.50% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: OLA, OLC, CLR, 9XT, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal		Chain	Bond lengths		Bond angles	
	MIGI	Chain	RMSZ	# Z >5	RMSZ	# Z > 5
	1	Α	0.26	0/3114	0.38	0/4235

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1106	LEU	Mainchain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	Α	3047	0	3112	17	0
2	A	24	0	0	0	0

Continued on next page...



Continued	trom	nromanne	naae
-	110116	picolous	puyc

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	217	0	329	6	0
4	A	1	0	0	0	0
5	A	84	0	138	1	0
6	A	156	0	232	4	0
7	A	30	0	0	0	0
All	All	3559	0	3811	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 22 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{array}{l} ext{Interatomic} \ ext{distance} \ (ext{Å}) \end{array}$	$egin{array}{c} ext{Clash} \ ext{overlap } (ext{Å}) \end{array}$
1:A:25:VAL:HG22	3:A:1214:OLA:H132	1.75	0.68
6:A:1223:OLC:H17A	6:A:1227:OLC:H17	1.81	0.62
1:A:65:THR:HG23	6:A:1227:OLC:H4A	1.83	0.61
1:A:174:MET:HG3	1:A:257:PHE:HB2	1.85	0.58
1:A:29[A]:TRP:HZ3	3:A:1214:OLA:H62	1.74	0.52

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percen	tiles
1	A	389/434 (90%)	386 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	317/353 (90%)	316 (100%)	1 (0%)	92 98		

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1106	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

Of 28 ligands modelled in this entry, 1 is monoatomic - leaving 27 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mal	TD	Chain	Dag	T ! 1-	Во	ond leng	ths	В	ond ang	;les
Mol	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	OLA	A	1208	_	14,17,19	0.75	1 (7%)	13,17,19	0.60	0
5	CLR	A	1204	-	31,31,31	0.66	0	48,48,48	0.91	1 (2%)
3	OLA	A	1212	-	4,7,19	0.24	0	3,7,19	0.52	0
3	OLA	A	1217	-	12,12,19	0.83	1 (8%)	11,11,19	0.51	0
3	OLA	A	1210	-	16,19,19	0.72	1 (6%)	15,19,19	0.58	0
3	OLA	A	1213	-	8,8,19	0.34	0	7,7,19	0.75	0
6	OLC	A	1226	-	16,19,24	0.22	0	15,19,25	0.58	0
3	OLA	A	1207	_	5,8,19	0.23	0	4,8,19	0.54	0
3	OLA	A	1202	_	16,19,19	0.71	1 (6%)	15,19,19	0.64	0
3	OLA	A	1211	-	8,11,19	1.00	1 (12%)	7,11,19	0.82	0
5	CLR	A	1205	-	31,31,31	0.68	0	48,48,48	0.88	0
3	OLA	A	1215	-	10,13,19	0.93	1 (10%)	8,13,19	0.54	0
6	OLC	A	1224	-	23,23,24	0.96	1 (4%)	24,24,25	0.95	1 (4%)
3	OLA	A	1216	-	9,9,19	0.95	1 (11%)	7,8,19	0.50	0
6	OLC	A	1222	_	13,16,24	0.25	0	12,16,25	0.50	0
6	OLC	A	1220	-	15,15,24	1.15	1 (6%)	16,16,25	1.10	2 (12%)
6	OLC	A	1225	-	21,21,24	0.98	1 (4%)	22,22,25	0.90	1 (4%)
3	OLA	A	1209	-	16,19,19	0.71	1 (6%)	15,19,19	0.58	0
6	OLC	A	1221	-	16,16,24	1.13	1 (6%)	17,17,25	1.19	2 (11%)
3	OLA	A	1219	-	12,12,19	0.82	1 (8%)	11,11,19	0.57	0
5	CLR	A	1206	-	31,31,31	0.63	0	48,48,48	0.87	1 (2%)
3	OLA	A	1228	-	16,19,19	0.72	1 (6%)	15,19,19	0.58	0
3	OLA	A	1214	-	15,18,19	0.73	1 (6%)	14,18,19	0.64	0
3	OLA	A	1218	-	11,11,19	0.86	1 (9%)	9,10,19	0.54	0
6	OLC	A	1223	-	16,19,24	0.22	0	15,19,25	0.62	0
2	9XT	A	1201	_	21,27,27	1.22	3 (14%)	22,39,39	2.05	5 (22%)
6	OLC	A	1227	_	16,19,24	0.20	0	15,19,25	0.69	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	OLA	A	1208	-	-	4/13/15/17	1
5	CLR	A	1204	-	-	0/10/68/68	0/4/4/4
3	OLA	A	1212	-	-	2/3/5/17	-
3	OLA	A	1217	-	-	1/10/10/17	-

Continued on next page...



 $Continued\ from\ previous\ page...$

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	OLA	A	1210	-	-	8/15/17/17	-
3	OLA	A	1213	-	-	1/6/6/17	-
6	OLC	A	1226	-	-	4/15/17/24	-
3	OLA	A	1207	-	-	3/4/6/17	-
3	OLA	A	1202	-	-	9/15/17/17	-
3	OLA	A	1211	-	-	4/7/9/17	-
5	CLR	A	1205	-	-	1/10/68/68	0/4/4/4
3	OLA	A	1215	-	-	7/9/11/17	-
6	OLC	A	1224	-	-	3/23/23/24	-
3	OLA	A	1216	-	-	7/7/7/17	-
6	OLC	A	1222	_	-	3/12/14/24	-
6	OLC	A	1220	-	-	0/15/15/24	-
6	OLC	A	1225	-	-	2/21/21/24	-
3	OLA	A	1209	-	-	7/15/17/17	-
6	OLC	A	1221	-	-	2/16/16/24	-
3	OLA	A	1219	-	-	5/10/10/17	-
5	CLR	A	1206	-	-	2/10/68/68	0/4/4/4
3	OLA	A	1228	-	-	10/15/17/17	-
3	OLA	A	1214	-	-	7/14/16/17	-
3	OLA	A	1218	-	-	6/9/9/17	-
6	OLC	A	1223		-	2/15/17/24	-
2	9XT	A	1201	-	-	0/4/8/8	0/4/4/4
6	OLC	A	1227		-	6/15/17/24	-

The worst 5 of 19 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	$\operatorname{Ideal}(ext{\AA})$
6	A	1224	OLC	O20-C1	4.34	1.46	1.33
6	A	1221	OLC	O20-C1	4.29	1.45	1.33
6	A	1225	OLC	O20-C1	4.27	1.45	1.33
6	A	1220	OLC	O20-C1	4.24	1.45	1.33
2	A	1201	9XT	N3-N2	3.02	1.40	1.34

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	${f Z}$	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
2	A	1201	9XT	C11-N6-C12	5.20	121.29	115.36
2	A	1201	9XT	C8-N2-N3	3.93	126.69	117.11

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\mathbf{Ideal}(^o)$
2	A	1201	9XT	C10-N5-C11	3.91	120.98	117.31
2	A	1201	9XT	N6-C11-N5	-3.65	122.36	127.22
6	A	1221	OLC	O20-C1-C2	3.16	121.84	111.91

There are no chirality outliers.

5 of 106 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1212	OLA	C1-C2-C3-C4
3	A	1210	OLA	C1-C2-C3-C4
3	A	1207	OLA	C1-C2-C3-C4
3	A	1211	OLA	C1-C2-C3-C4
3	A	1215	OLA	C1-C2-C3-C4


There are no ring outliers.


11 monomers are involved in 10 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1208	OLA	1	0
3	A	1217	OLA	1	0
3	A	1207	OLA	1	0
3	A	1202	OLA	1	0
6	A	1224	OLC	1	0
6	A	1220	OLC	1	0
6	A	1221	OLC	1	0
5	A	1206	CLR	1	0
3	A	1214	OLA	3	0
6	A	1223	OLC	1	0
6	A	1227	OLC	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














































































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$ $#$ RSRZ $>$ 2		$OWAB(A^2)$	Q < 0.9	
1	A	390/434 (89%)	-0.06	22 (5%) 24	19	14, 36, 80, 139	0

The worst 5 of 22 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1057	GLU	7.4
1	A	1056	PRO	5.6
1	A	1061	PHE	5.3
1	A	1059	LYS	4.9
1	A	1060	ASP	4.8

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	${f Res}$	Atoms	RSCC	RSR	$oxed{f B-factors({ m \AA}^2)}$	Q<0.9
3	OLA	A	1215	14/20	0.60	0.34	49,69,73,74	0
3	OLA	A	1209	20/20	0.65	0.33	36,62,76,77	0

Continued on next page...



Continued from previous page...


Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B\text{-factors}}({f \AA}^2)$	Q<0.9
3	OLA	A	1213	9/20	0.75	0.19	55,56,58,60	0
3	OLA	A	1217	13/20	0.78	0.27	54,59,68,69	0
3	OLA	A	1208	18/20	0.79	0.24	40,59,71,74	0
3	OLA	A	1210	20/20	0.80	0.33	46,61,68,69	0
3	OLA	A	1212	8/20	0.80	0.26	56,69,75,76	0
6	OLC	A	1227	20/25	0.80	0.26	38,58,74,78	0
3	OLA	A	1216	10/20	0.81	0.20	54,59,61,61	0
3	OLA	A	1211	12/20	0.81	0.46	37,55,69,70	0
3	OLA	A	1219	13/20	0.81	0.27	43,47,60,61	0
3	OLA	A	1214	19/20	0.81	0.25	58,62,67,68	0
3	OLA	A	1218	12/20	0.81	0.25	51,57,62,62	0
6	OLC	A	1225	22/25	0.81	0.30	52,64,73,80	0
6	OLC	A	1221	17/25	0.82	0.26	55,63,70,78	0
6	OLC	A	1224	24/25	0.83	0.26	34,56,67,72	0
3	OLA	A	1202	20/20	0.86	0.19	36,48,50,53	0
6	OLC	A	1222	17/25	0.86	0.19	34,45,62,62	0
6	OLC	A	1226	20/25	0.86	0.24	25,54,78,85	0
3	OLA	A	1228	20/20	0.87	0.19	62,63,64,65	20
6	OLC	A	1220	16/25	0.90	0.18	38,48,57,71	0
6	OLC	A	1223	20/25	0.92	0.20	30,41,68,69	0
3	OLA	A	1207	9/20	0.92	0.20	36,44,61,62	0
4	NA	A	1203	1/1	0.93	0.35	54,54,54,54	0
5	CLR	A	1205	28/28	0.94	0.15	27,33,36,41	0
5	CLR	A	1206	28/28	0.95	0.14	20,32,49,52	0
5	CLR	A	1204	28/28	0.95	0.15	28,32,62,67	0
2	9XT	A	1201	24/24	0.98	0.16	12,22,26,32	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around OLA A 1215:


 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around OLA A 1209:












Electron density around OLA A 1213:


 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around OLA A 1217:












Electron density around OLA A 1208:


 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around OLA A 1210:












Electron density around OLA A 1212:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around OLC A 1227:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around OLA A 1216:


 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around OLA A 1211:












Electron density around OLA A 1219:


 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around OLA A 1214:












Electron density around OLA A 1218:


 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around OLC A 1225:












Electron density around OLC A 1221:


 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around OLC A 1224:












Electron density around OLA A 1202:


 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around OLC A 1222:












Electron density around OLC A 1226:


 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around OLA A 1228:











Electron density around OLC A 1220:


 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around OLC A 1223:












Electron density around OLA A 1207:


 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around CLR A 1205:












Electron density around CLR A 1206:


 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around CLR A 1204:











6.5 Other polymers (i)

There are no such residues in this entry.

