



# Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 01:05 pm BST

PDB ID : 6OR9  
Title : Structure of L-lactate dehydrogenase from *Trichoplusia ni*  
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)  
Deposited on : 2019-04-29  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	364	 79% 16%
1	B	364	 79% 6% 15%
1	C	364	 79% 5% 16%
1	D	364	 74% 7% 18%

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 10401 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called L-lactate dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	306	2330	1471	408	438	13	0	12	0
1	B	308	2398	1519	414	451	14	0	19	0
1	C	305	2306	1461	397	435	13	0	9	0
1	D	299	2241	1420	380	428	13	0	7	0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	287	Total 293	O 293	0	6
2	B	322	Total 334	O 334	0	13
2	C	276	Total 284	O 284	0	8
2	D	212	Total 215	O 215	0	4





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.39Å 76.48Å 79.52Å 113.06° 109.84° 88.70°	Depositor
Resolution (Å)	48.91 – 1.80 48.91 – 1.80	Depositor EDS
% Data completeness (in resolution range)	97.0 (48.91-1.80) 97.0 (48.91-1.80)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.73 (at 1.79Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.158 , 0.195 0.158 , 0.195	Depositor DCC
$R_{free}$ test set	1971 reflections (1.71%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.1	Xtrriage
Anisotropy	0.364	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 51.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	10401	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.78% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.39	0/2399	0.55	0/3254
1	B	0.40	0/2487	0.56	0/3373
1	C	0.38	0/2367	0.56	0/3215
1	D	0.34	0/2295	0.54	0/3121
All	All	0.38	0/9548	0.55	0/12963

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2330	0	2385	15	0
1	B	2398	0	2465	16	0
1	C	2306	0	2348	13	0
1	D	2241	0	2252	14	0
2	A	293	0	0	4	0
2	B	334	0	0	7	0
2	C	284	0	0	5	0
2	D	215	0	0	1	0
All	All	10401	0	9450	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:224:GLU:OE1	2:B:401:HOH:O	2.02	0.77
1:B:122[A]:LYS:NZ	2:B:403[A]:HOH:O	2.22	0.72
1:C:350:GLN:OE1	2:C:401:HOH:O	2.08	0.70
1:A:201[B]:ARG:NH2	1:A:266:VAL:O	2.25	0.70
1:A:317:ASP:OD1	2:A:403:HOH:O	2.12	0.66
1:A:364:PHE:OXT	2:A:402:HOH:O	2.12	0.66
1:D:268:LYS:O	1:D:272[A]:GLU:HG2	1.98	0.62
1:B:138:ARG:NH2	2:B:408:HOH:O	2.33	0.61
1:B:353[B]:ASP:OD1	1:B:357:LYS:NZ	2.34	0.61
1:A:68:VAL:HG13	1:A:80[A]:ILE:HD13	1.83	0.60
1:C:309:LEU:HD21	1:C:315[B]:ILE:HD13	1.84	0.59
1:C:298:ASN:HB2	1:C:329:ARG:HG2	1.85	0.58
1:A:170:ASN:ND2	2:A:407:HOH:O	2.36	0.57
1:B:174[A]:ILE:HD11	1:B:362:ILE:HD11	1.89	0.55
1:D:84:ASP:H	1:D:90:LEU:HD11	1.73	0.54
1:C:237:ASN:OD1	2:C:402:HOH:O	2.19	0.53
1:D:246:LEU:HD22	1:D:248:GLN:HG2	1.90	0.52
1:B:298:ASN:HB2	1:B:329:ARG:HG2	1.91	0.51
1:B:122[B]:LYS:HE2	1:B:329:ARG:NE	2.25	0.51
1:C:104:MET:O	2:C:403:HOH:O	2.19	0.50
1:B:62[A]:GLN:NE2	1:B:131:ARG:HH2	2.09	0.50
1:C:90:LEU:HD22	1:C:111:ALA:HB1	1.95	0.49
1:A:201[A]:ARG:HG2	1:A:266:VAL:HG2	1.95	0.48
1:C:345:ARG:HD3	2:C:489:HOH:O	2.13	0.48
1:A:210:LEU:HD21	1:A:259:TRP:HH2	1.78	0.48
1:D:298:ASN:HB2	1:D:329:ARG:HG2	1.95	0.48
1:B:170:ASN:ND2	2:B:402:HOH:O	2.13	0.47
1:C:313:HIS:HB2	1:C:315[B]:ILE:HD12	1.96	0.47
1:B:146:THR:HG22	1:B:150:LYS:HE2	1.97	0.47
1:B:147:ASP:HA	1:B:150:LYS:HE3	1.95	0.47
1:B:214[B]:THR:HG21	2:B:426:HOH:O	2.15	0.47
1:D:305:VAL:O	1:D:322:SER:HA	2.15	0.46
1:D:68:VAL:HG13	1:D:80:ILE:HD12	1.97	0.46
1:C:115:TYR:OH	1:C:152:ILE:HG12	2.15	0.46
1:A:170:ASN:ND2	2:A:401:HOH:O	2.07	0.46
1:A:146:THR:HG22	1:A:150:LYS:HE3	1.98	0.45
1:A:174[A]:ILE:HD11	1:A:362:ILE:HD11	1.99	0.45
1:C:101:SER:HA	1:C:104:MET:CE	2.45	0.45
1:B:170:ASN:ND2	2:B:416[A]:HOH:O	2.47	0.44
1:B:74:GLN:O	1:B:293:ARG:NH2	2.50	0.44

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:101:SER:HA	1:C:104:MET:HE3	2.00	0.44
1:D:90:LEU:HD22	1:D:111:ALA:HB1	1.99	0.44
1:C:84:ASP:H	1:C:90:LEU:HD11	1.83	0.43
1:D:212[A]:ILE:HD11	1:D:217:CYS:SG	2.58	0.43
1:A:174[B]:ILE:HD11	1:A:355:MET:HG3	2.01	0.43
1:B:310:LYS:NZ	2:B:419:HOH:O	2.50	0.43
1:A:214[B]:THR:HG21	2:D:406:HOH:O	2.18	0.43
1:D:84:ASP:OD2	1:D:85:VAL:N	2.46	0.42
1:D:181:LYS:HD3	1:D:364:PHE:CZ	2.54	0.42
1:D:309:LEU:HD21	1:D:315[B]:ILE:HD13	2.01	0.42
1:A:201[B]:ARG:HG3	1:A:201[B]:ARG:HH11	1.85	0.42
1:D:313:HIS:HB2	1:D:315[B]:ILE:HD12	2.03	0.41
1:C:122:LYS:HD2	2:C:411:HOH:O	2.20	0.41
1:B:305:VAL:O	1:B:322:SER:HA	2.20	0.41
1:A:62[A]:GLN:HG2	1:A:279:TYR:CE1	2.56	0.41
1:D:94:LEU:HB2	1:D:111:ALA:HB3	2.03	0.41
1:A:201[B]:ARG:NH2	1:A:269:SER:OG	2.54	0.40
1:D:85:VAL:HG23	1:D:86:MET:HG3	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	314/364 (86%)	309 (98%)	5 (2%)	0	100	100
1	B	323/364 (89%)	318 (98%)	5 (2%)	0	100	100
1	C	310/364 (85%)	302 (97%)	7 (2%)	1 (0%)	41	27
1	D	300/364 (82%)	294 (98%)	5 (2%)	1 (0%)	41	27
All	All	1247/1456 (86%)	1223 (98%)	22 (2%)	2 (0%)	47	33

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	106	ASN
1	C	106	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	260/312 (83%)	257 (99%)	3 (1%)	71 65
1	B	271/312 (87%)	271 (100%)	0	100 100
1	C	256/312 (82%)	254 (99%)	2 (1%)	81 78
1	D	247/312 (79%)	239 (97%)	8 (3%)	39 25
All	All	1034/1248 (83%)	1021 (99%)	13 (1%)	73 62

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	201[A]	ARG
1	A	201[B]	ARG
1	A	355	MET
1	C	207	SER
1	C	271	TYR
1	D	80	ILE
1	D	169	SER
1	D	207	SER
1	D	212[A]	ILE
1	D	212[B]	ILE
1	D	267	VAL
1	D	271	TYR
1	D	355	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	218	HIS
1	C	99	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	306/364 (84%)	-0.47	2 (0%) 87 86	14, 23, 51, 74	0
1	B	308/364 (84%)	-0.41	1 (0%) 94 92	13, 22, 41, 80	0
1	C	305/364 (83%)	-0.40	2 (0%) 87 86	14, 25, 49, 79	0
1	D	299/364 (82%)	-0.23	5 (1%) 70 66	16, 30, 57, 82	0
All	All	1218/1456 (83%)	-0.38	10 (0%) 86 84	13, 24, 52, 82	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	259	TRP	5.4
1	A	259	TRP	4.2
1	D	90	LEU	3.4
1	D	112	SER	2.9
1	D	99	HIS	2.7
1	B	259	TRP	2.6
1	D	259	TRP	2.5
1	C	258	ASN	2.4
1	A	262	THR	2.2
1	D	246	LEU	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.