



Full wwPDB X-ray Structure Validation Report

Jun 29, 2021 – 11:29 AM BST

PDB ID : 7OTJ
Title : Crystal structure of Pif1 helicase from *Candida albicans*
Authors : Rety, S.; Xi, X.G.
Deposited on : 2021-06-10
Resolution : 2.58 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.20
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.20


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.58 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3676 (2.60-2.56)
Clashscore	141614	4049 (2.60-2.56)
Ramachandran outliers	138981	3979 (2.60-2.56)
Sidechain outliers	138945	3979 (2.60-2.56)
RSRZ outliers	127900	3614 (2.60-2.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	518	 5% (red), 73% (green), 11% (yellow), 15% (grey)
1	B	518	 6% (red), 74% (green), 10% (yellow), 16% (grey)
2	C	6	 67% (green), 33% (yellow)
2	D	6	 83% (green), 17% (yellow)

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	PO4	B	1007	-	-	-	X

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 7347 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP-dependent DNA helicase PIF1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	438	Total 3472	C 2218	N 606	O 635	S 13	0	0	0
1	B	436	Total 3447	C 2196	N 604	O 636	S 11	0	0	0


There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	366	MET	-	initiating methionine	UNP Q59RQ0
A	565	ASP	ASN	conflict	UNP Q59RQ0
A	744	ASP	GLU	conflict	UNP Q59RQ0
B	366	MET	-	initiating methionine	UNP Q59RQ0
B	565	ASP	ASN	conflict	UNP Q59RQ0
B	744	ASP	GLU	conflict	UNP Q59RQ0

- Molecule 2 is a DNA chain called DNA (5'-D(P*TP*TP*TP*TP*TP*T)-3').


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	C	6	Total 105	C 50	N 10	O 39	P 6	0	0	0
2	D	6	Total 105	C 50	N 10	O 39	P 6	0	0	0

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
3	A	1	27	10	5	10	2	0	0
3	B	1	27	10	5	10	2	0	0

- Molecule 4 is TETRAFLUOROALUMINATE ION (three-letter code: ALF) (formula: AlF_4) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Al F		
4	A	1	5	1 4	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	Al	F		
4	B	1	5	1	4	0	0


- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
5	A	1	1	1	0	0
5	B	1	1	1	0	0

- Molecule 6 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	K		
6	A	7	7	7	0	0
6	B	3	3	3	0	0

- Molecule 7 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
7	A	1	5	4	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	O	P	0	0
			5	4	1		
7	A	1	Total	O	P	0	0
			5	4	1		
7	A	1	Total	O	P	0	0
			5	4	1		
7	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	59	Total	O	0	0
			59	59		
8	B	50	Total	O	0	0
			50	50		
8	C	3	Total	O	0	0
			3	3		
8	D	5	Total	O	0	0
			5	5		

Chain C:  67% 33%



- Molecule 2: DNA (5'-D(P*TP*TP*TP*TP*TP*T)-3')

Chain D:  83% 17%



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	198.97Å 79.96Å 94.11Å 90.00° 117.28° 90.00°	Depositor
Resolution (Å)	82.51 – 2.58 82.51 – 2.58	Depositor EDS
% Data completeness (in resolution range)	96.5 (82.51-2.58) 96.7 (82.51-2.58)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.94 (at 2.58Å)	Xtrriage
Refinement program	PHENIX 1.19.2_4158, PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.206 , 0.242 0.203 , 0.238	Depositor DCC
R_{free} test set	2072 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å ²)	51.8	Xtrriage
Anisotropy	0.616	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 43.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	0.008 for -h-2*1,-k,l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7347	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, ALF, MG, K, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.28	0/3529	0.52	0/4760
1	B	0.28	0/3503	0.51	0/4727
2	C	1.17	1/114 (0.9%)	1.29	0/173
2	D	1.17	1/114 (0.9%)	1.27	0/173
All	All	0.34	2/7260 (0.0%)	0.56	0/9833

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1	DT	OP3-P	-10.64	1.48	1.61
2	D	1	DT	OP3-P	-10.56	1.48	1.61

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3472	0	3584	35	0
1	B	3447	0	3536	33	0
2	C	105	0	60	1	0
2	D	105	0	60	0	0
3	A	27	0	12	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	27	0	12	1	0
4	A	5	0	0	0	0
4	B	5	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	7	0	0	0	0
6	B	3	0	0	0	0
7	A	20	0	0	0	0
7	B	5	0	0	0	0
8	A	59	0	0	1	0
8	B	50	0	0	1	0
8	C	3	0	0	0	0
8	D	5	0	0	0	0
All	All	7347	0	7264	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:812:ALA:O	1:A:814:SER:N	2.20	0.74
1:B:614:ARG:NH2	8:B:1101:HOH:O	2.22	0.71
1:B:708:ASN:HB2	1:B:785:VAL:HB	1.77	0.67
1:A:603:ALA:HB2	1:A:835:ILE:HD12	1.77	0.66
1:B:631:THR:HG1	1:B:803:SER:HG	1.42	0.63
1:A:832:MET:HA	1:A:835:ILE:HD13	1.82	0.62
1:B:418:THR:HG21	1:B:433:LEU:HD13	1.82	0.62
1:B:600:GLU:OE1	1:B:615:LYS:NZ	2.31	0.61
1:B:634:LEU:HD13	1:B:801:LEU:HB3	1.85	0.59
1:B:648:ALA:O	1:B:804:ARG:NH1	2.38	0.56
1:A:575:VAL:HB	1:A:861:ARG:HG2	1.88	0.55
1:B:634:LEU:HB3	1:B:639:LYS:HB2	1.89	0.55
1:A:872:GLU:HA	1:A:875:LYS:HG2	1.88	0.54
1:B:642:VAL:HG11	1:B:802:VAL:HG11	1.88	0.54
1:A:772:LYS:HA	1:A:783:THR:HA	1.91	0.52
1:A:761:LYS:HE3	1:A:805:ILE:HD11	1.91	0.52
1:A:439:PHE:CE1	1:A:488:ILE:HD13	2.46	0.51
1:A:619:ILE:O	1:A:655:LYS:NZ	2.36	0.50
1:B:453:ILE:HD12	1:B:488:ILE:HD13	1.94	0.49
1:A:623:VAL:HB	1:A:655:LYS:HE3	1.95	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:398:VAL:HG21	3:B:1001:ADP:H2'	1.93	0.48
1:A:398:VAL:HG21	3:A:1001:ADP:H2'	1.94	0.48
1:B:401:ARG:NH1	1:B:427:ASN:O	2.47	0.48
1:A:667:ASP:HB3	1:A:670:LEU:H	1.78	0.48
1:A:387:PHE:O	1:A:548:THR:HA	2.14	0.47
1:B:384:VAL:HG23	1:B:386:LEU:HD23	1.96	0.47
1:A:571:ARG:O	1:A:866:SER:OG	2.19	0.47
1:A:418:THR:HA	1:A:431:ILE:O	2.14	0.47
1:B:786:VAL:HG12	1:B:807:PHE:HD2	1.80	0.46
1:B:418:THR:HG22	1:B:471:ILE:HA	1.98	0.46
1:B:387:PHE:O	1:B:548:THR:HA	2.16	0.46
1:B:399:LEU:O	1:B:403:ILE:HG12	2.16	0.45
1:A:678:VAL:HG21	1:A:681:PHE:CZ	2.52	0.45
1:B:498:PRO:HG3	1:B:543:GLU:HG2	1.99	0.45
1:B:554:ILE:HD12	1:B:564:ILE:HG23	1.98	0.45
1:B:707:LEU:CD2	1:B:787:GLU:H	2.30	0.45
1:B:706:GLY:O	1:B:707:LEU:HD23	2.17	0.45
1:B:707:LEU:HD22	1:B:707:LEU:HA	1.93	0.44
1:B:775:LEU:HB3	1:B:776:PRO:HD2	2.00	0.44
1:A:669:GLN:NE2	1:A:775:LEU:HD13	2.33	0.44
1:A:516:VAL:HG13	2:C:2:DT:C2	2.53	0.43
1:B:387:PHE:CZ	1:B:389:THR:HB	2.53	0.43
1:A:387:PHE:CZ	1:A:389:THR:HB	2.52	0.43
1:A:679:ILE:HD11	1:A:772:LYS:HB2	1.99	0.43
1:B:680:ASP:OD1	1:B:681:PHE:N	2.50	0.43
1:A:376:VAL:HB	1:A:399:LEU:HD21	2.00	0.43
1:A:758:ASP:HA	1:A:761:LYS:HD3	2.00	0.43
1:A:797:ASP:OD2	1:B:772:LYS:HD2	2.18	0.43
1:B:418:THR:HA	1:B:431:ILE:O	2.19	0.43
1:A:574:ASN:O	8:A:1101:HOH:O	2.21	0.43
1:A:794:GLU:HG2	1:A:800:VAL:HA	2.01	0.42
1:B:770:LEU:HD13	1:B:785:VAL:HG22	2.00	0.42
1:A:764:LYS:HG3	1:A:768:TYR:OH	2.19	0.42
1:A:669:GLN:HB3	1:A:782:ARG:HH21	1.84	0.42
1:A:375:TYR:O	1:A:379:GLN:HG2	2.19	0.42
1:B:560:ASP:OD2	1:B:563:PHE:HB2	2.20	0.42
1:B:418:THR:HB	1:B:470:ILE:O	2.20	0.41
1:A:588:ARG:NH2	1:A:854:LEU:O	2.53	0.41
1:A:665:ASN:HA	1:A:671:VAL:HA	2.01	0.41
1:A:768:TYR:CE2	1:A:787:GLU:HG2	2.55	0.41
1:A:484:LYS:O	1:A:488:ILE:HD12	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:669:GLN:HG2	1:A:775:LEU:HD22	2.02	0.41
1:A:827:LYS:HA	1:A:853:GLY:O	2.21	0.41
1:B:703:GLU:HB3	1:B:704:VAL:H	1.62	0.41
1:B:766:LYS:HG2	1:B:767:LYS:H	1.86	0.41
1:B:707:LEU:HD22	1:B:787:GLU:H	1.86	0.40
1:A:795:ASP:OD1	1:A:799:THR:N	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	432/518 (83%)	426 (99%)	4 (1%)	2 (0%)	29	50
1	B	428/518 (83%)	428 (100%)	0	0	100	100
All	All	860/1036 (83%)	854 (99%)	4 (0%)	2 (0%)	47	69

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	813	TRP
1	A	765	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	387/461 (84%)	384 (99%)	3 (1%)	81	92
1	B	384/461 (83%)	379 (99%)	5 (1%)	69	85
All	All	771/922 (84%)	763 (99%)	8 (1%)	76	89

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	488	ILE
1	A	588	ARG
1	A	813	TRP
1	B	418	THR
1	B	588	ARG
1	B	670	LEU
1	B	704	VAL
1	B	707	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 21 ligands modelled in this entry, 12 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ADP	A	1001	5	24,29,29	0.95	1 (4%)	29,45,45	1.34	4 (13%)
7	PO4	A	1012	-	4,4,4	0.91	0	6,6,6	0.40	0
4	ALF	B	1002	-	0,4,4	0.00	-	-	-	-
7	PO4	A	1014	-	4,4,4	0.91	0	6,6,6	0.52	0
4	ALF	A	1002	-	0,4,4	0.00	-	-	-	-
3	ADP	B	1001	5	24,29,29	0.96	1 (4%)	29,45,45	1.31	3 (10%)
7	PO4	A	1013	-	4,4,4	0.94	0	6,6,6	0.40	0
7	PO4	A	1011	-	4,4,4	0.89	0	6,6,6	0.59	0
7	PO4	B	1007	-	4,4,4	0.92	0	6,6,6	0.65	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ADP	B	1001	5	-	0/12/32/32	0/3/3/3
3	ADP	A	1001	5	-	0/12/32/32	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1001	ADP	C5-C4	2.46	1.47	1.40
3	B	1001	ADP	C5-C4	2.39	1.47	1.40

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1001	ADP	N3-C2-N1	-3.17	123.72	128.68
3	B	1001	ADP	N3-C2-N1	-3.07	123.87	128.68
3	B	1001	ADP	C4-C5-N7	-2.77	106.51	109.40
3	A	1001	ADP	C4-C5-N7	-2.65	106.64	109.40
3	B	1001	ADP	PA-O3A-PB	-2.65	123.73	132.83
3	A	1001	ADP	PA-O3A-PB	-2.54	124.11	132.83
3	A	1001	ADP	C3'-C2'-C1'	2.07	104.09	100.98

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.


2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1001	ADP	1	0
3	B	1001	ADP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	438/518 (84%)	0.52	27 (6%) 20 17	39, 58, 114, 164	0
1	B	436/518 (84%)	0.55	31 (7%) 16 13	39, 57, 126, 177	0
2	C	6/6 (100%)	0.04	0 100 100	53, 58, 84, 105	0
2	D	6/6 (100%)	-0.05	0 100 100	58, 62, 82, 103	0
All	All	886/1048 (84%)	0.52	58 (6%) 18 16	39, 57, 122, 177	0

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	633	ILE	10.9
1	B	703	GLU	8.9
1	B	632	GLY	8.2
1	B	636	GLU	6.4
1	A	688	MET	6.1
1	A	763	TYR	5.8
1	B	763	TYR	5.8
1	B	637	PRO	5.8
1	A	755	LEU	5.1
1	B	802	VAL	5.1
1	B	647	LEU	5.0
1	A	686	THR	5.0
1	B	799	THR	4.5
1	B	635	PRO	4.4
1	B	801	LEU	4.4
1	A	685	ASP	4.3
1	A	756	LYS	4.2
1	A	684	ARG	4.2
1	B	705	SER	4.1
1	A	594	GLU	4.1
1	A	681	PHE	4.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	762	ASP	4.0
1	A	812	ALA	4.0
1	A	687	TYR	3.7
1	B	455	ARG	3.6
1	B	439	PHE	3.4
1	A	765	ASN	3.3
1	B	800	VAL	3.2
1	B	634	LEU	3.2
1	B	794	GLU	3.2
1	B	666	PHE	3.2
1	B	797	ASP	3.1
1	B	631	THR	3.1
1	A	670	LEU	3.0
1	B	708	ASN	2.9
1	A	766	LYS	2.9
1	A	439	PHE	2.8
1	A	366	MET	2.8
1	B	798	GLY	2.7
1	B	460	PHE	2.5
1	B	638	GLN	2.5
1	B	645	ASN	2.5
1	A	754	LYS	2.5
1	A	781	PHE	2.5
1	A	595	GLY	2.5
1	A	680	ASP	2.4
1	A	622	ASP	2.4
1	A	682	VAL	2.4
1	B	459	ALA	2.3
1	A	779	ILE	2.3
1	B	670	LEU	2.1
1	A	683	ASP	2.1
1	A	656	VAL	2.1
1	B	707	LEU	2.1
1	B	456	ASN	2.1
1	A	760	MET	2.1
1	B	643	LEU	2.0
1	A	530	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	PO4	A	1011	5/5	0.63	0.22	93,100,130,158	0
7	PO4	B	1007	5/5	0.64	0.56	101,117,138,149	0
6	K	A	1007	1/1	0.80	0.24	93,93,93,93	0
6	K	A	1008	1/1	0.84	0.27	98,98,98,98	0
6	K	A	1009	1/1	0.86	0.20	87,87,87,87	0
6	K	A	1010	1/1	0.86	0.13	88,88,88,88	0
5	MG	A	1003	1/1	0.87	0.16	38,38,38,38	0
6	K	B	1006	1/1	0.88	0.21	107,107,107,107	0
7	PO4	A	1012	5/5	0.89	0.26	67,75,100,118	0
6	K	B	1004	1/1	0.90	0.16	74,74,74,74	0
6	K	A	1004	1/1	0.91	0.10	82,82,82,82	0
7	PO4	A	1014	5/5	0.92	0.23	83,90,115,120	0
7	PO4	A	1013	5/5	0.93	0.16	71,74,86,97	0
5	MG	B	1003	1/1	0.95	0.08	37,37,37,37	0
6	K	A	1005	1/1	0.96	0.20	93,93,93,93	0
3	ADP	B	1001	27/27	0.97	0.16	37,49,66,67	0
6	K	B	1005	1/1	0.97	0.20	81,81,81,81	0
3	ADP	A	1001	27/27	0.98	0.18	37,50,64,66	0
4	ALF	A	1002	5/5	0.98	0.17	39,43,47,48	0
6	K	A	1006	1/1	0.98	0.14	79,79,79,79	0
4	ALF	B	1002	5/5	0.99	0.21	37,37,47,50	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.












Electron density around ADP A 1001:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ALF A 1002:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.