



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 9, 2023 – 10:10 am BST

PDB ID : 8OXW
Title : Transglutaminase 3 in complex with DH patient-derived Fab DH63-B02
Authors : Heggelund, J.E.; Sollid, L.M.
Deposited on : 2023-05-02
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION




The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	693	
2	B	225	
3	C	216	

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 18576 atoms, of which 8640 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein-glutamine gamma-glutamyltransferase E 27 kDa non-catalytic chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	679	10687	3386	5314	933	1028	26	139	10	0


- Molecule 2 is a protein called Antibody fab fragment heavy chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	B	225	3412	1079	1687	295	342	9	73	4	0

- Molecule 3 is a protein called Antibody fab fragment light chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
3	C	216	3208	1013	1573	274	342	6	74	1	0

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	1	0
			10	2	6	2		
4	A	1	Total	C	H	O	1	0
			10	2	6	2		
4	A	1	Total	C	H	O	1	0
			10	2	6	2		
4	A	1	Total	C	H	O	1	0
			10	2	6	2		
4	A	1	Total	C	H	O	1	0
			10	2	6	2		
4	A	1	Total	C	H	O	1	0
			10	2	6	2		
4	A	1	Total	C	H	O	1	0
			10	2	6	2		
4	A	1	Total	C	H	O	1	0
			10	2	6	2		
4	C	1	Total	C	H	O	1	0
			10	2	6	2		

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	3	Total	Ca	0	0
			3	3		

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Cl	0	0
			1	1		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	631	Total	O	0	1
			632	632		
7	B	230	Total	O	0	1
			231	231		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	C	290	Total 292	O 292	0	2


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Protein-glutamine gamma-glutamyltransferase E 27 kDa non-catalytic chain



- Molecule 2: Antibody fab fragment heavy chain



- Molecule 3: Antibody fab fragment light chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	81.50Å 93.49Å 90.64Å 90.00° 92.63° 90.00°	Depositor
Resolution (Å)	45.31 – 1.70 45.27 – 1.70	Depositor EDS
% Data completeness (in resolution range)	98.4 (45.31-1.70) 97.1 (45.27-1.70)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.17 (at 1.70Å)	Xtrriage
Refinement program	REFMAC 5.8.0405	Depositor
R, R_{free}	0.172 , 0.210 0.176 , 0.211	Depositor DCC
R_{free} test set	7361 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	26.2	Xtrriage
Anisotropy	0.516	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 44.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.016 for -h,-l,-k 0.000 for -h,l,k 0.025 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	18576	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, CL, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.58	0/5490	0.91	5/7443 (0.1%)
2	B	0.58	0/1768	0.85	0/2403
3	C	0.69	3/1674 (0.2%)	0.92	2/2284 (0.1%)
All	All	0.60	3/8932 (0.0%)	0.90	7/12130 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
2	B	0	1
3	C	0	1
All	All	0	4

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	84	GLU	CD-OE1	9.67	1.36	1.25
3	C	127	GLU	CD-OE1	5.60	1.31	1.25
3	C	202	GLU	CD-OE2	5.42	1.31	1.25

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	248	ARG	NE-CZ-NH2	-7.06	116.77	120.30
1	A	269	ARG	NE-CZ-NH2	-6.93	116.83	120.30
1	A	248	ARG	NE-CZ-NH1	6.38	123.49	120.30
1	A	340	ARG	NE-CZ-NH1	6.14	123.37	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	40	ARG	NE-CZ-NH1	-5.73	117.44	120.30
1	A	340	ARG	NE-CZ-NH2	-5.61	117.50	120.30
3	C	118	SER	CB-CA-C	5.09	119.78	110.10

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	17	ARG	Sidechain
1	A	201[A]	ARG	Sidechain
2	B	219	ARG	Sidechain
3	C	55	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5373	5314	5298	15	0
2	B	1725	1687	1682	6	0
3	C	1635	1573	1568	9	0
4	A	40	60	60	0	0
4	C	4	6	6	0	0
5	A	3	0	0	0	0
6	A	1	0	0	0	0
7	A	632	0	0	1	0
7	B	231	0	0	0	0
7	C	292	0	0	1	0
All	All	9936	8640	8614	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:201[A]:ARG:NH1	7:A:801:HOH:O	1.98	0.95

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:54:LEU:CD2	1:A:117:ILE:HG12	2.35	0.56
3:C:95:ASP:HB2	3:C:96:PRO:CD	2.36	0.56
3:C:106[A]:LYS:NZ	7:C:404:HOH:O	2.38	0.55
1:A:665[B]:ARG:HB2	1:A:669:LYS:HE3	1.90	0.53
1:A:638:LEU:HD23	1:A:638:LEU:N	2.24	0.53
1:A:386:MET:N	1:A:387:PRO:CD	2.73	0.52
3:C:163:VAL:HA	3:C:181:TYR:O	2.10	0.51
1:A:622:ASP:C	1:A:651:PRO:HB3	2.31	0.51
2:B:83[B]:MET:HB3	2:B:86:LEU:HD21	1.91	0.51
1:A:650:GLY:N	1:A:653:GLU:OE1	2.33	0.49
1:A:305:ARG:HH21	1:A:305:ARG:HG2	1.79	0.47
3:C:95:ASP:HB2	3:C:96:PRO:HD2	1.96	0.47
1:A:2:ALA:HB3	1:A:45:ASN:ND2	2.31	0.46
2:B:218:LYS:HE3	2:B:219:ARG:O	2.17	0.44
1:A:63:TYR:N	1:A:64:PRO:HD3	2.33	0.44
2:B:83[A]:MET:HB3	2:B:86:LEU:HD21	2.01	0.43
2:B:99:ASP:HA	2:B:108:GLY:O	2.19	0.43
2:B:187:LEU:HD12	2:B:187:LEU:C	2.39	0.43
1:A:192:ILE:HG23	1:A:223:MET:HB3	2.01	0.42
2:B:47:TRP:CG	3:C:99:VAL:HB	2.54	0.42
3:C:153:LYS:HA	3:C:157:SER:O	2.20	0.41
1:A:383:ASN:HB3	1:A:384:PHE:CD1	2.55	0.41
1:A:485:GLY:HA2	1:A:504:LEU:O	2.21	0.41
1:A:598:THR:O	1:A:615:MET:HA	2.21	0.41
1:A:383:ASN:HA	1:A:384:PHE:HA	1.91	0.40
3:C:55:ARG:HG2	3:C:59:VAL:HB	2.03	0.40
3:C:130:GLN:O	3:C:130:GLN:HG3	2.20	0.40
3:C:10:VAL:HG23	3:C:107:LEU:HD13	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	685/693 (99%)	662 (97%)	23 (3%)	0	100	100
2	B	227/225 (101%)	221 (97%)	6 (3%)	0	100	100
3	C	215/216 (100%)	206 (96%)	9 (4%)	0	100	100
All	All	1127/1134 (99%)	1089 (97%)	38 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	594/595 (100%)	585 (98%)	9 (2%)	65	51
2	B	195/191 (102%)	193 (99%)	2 (1%)	76	67
3	C	190/189 (100%)	187 (98%)	3 (2%)	62	48
All	All	979/975 (100%)	965 (99%)	14 (1%)	65	53

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	VAL
1	A	65	SER
1	A	77	SER
1	A	125	SER
1	A	211	ASN
1	A	611	VAL
1	A	615	MET
1	A	623	GLU
1	A	641	ASN
2	B	109	MET
2	B	165	SER
3	C	57	SER
3	C	118	SER
3	C	215	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 4 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	EDO	A	702	-	3,3,3	0.24	0	2,2,2	0.64	0
4	EDO	A	703	-	3,3,3	0.29	0	2,2,2	0.03	0
4	EDO	A	709	-	3,3,3	0.24	0	2,2,2	0.49	0
4	EDO	A	705	-	3,3,3	0.06	0	2,2,2	0.06	0
4	EDO	A	704	-	3,3,3	0.43	0	2,2,2	0.75	0
4	EDO	A	710	-	3,3,3	0.66	0	2,2,2	0.76	0
4	EDO	A	706	-	3,3,3	0.24	0	2,2,2	0.52	0
4	EDO	A	707	-	3,3,3	0.49	0	2,2,2	0.78	0
4	EDO	C	301	-	3,3,3	0.32	0	2,2,2	0.36	0
4	EDO	A	708	-	3,3,3	0.48	0	2,2,2	0.41	0
4	EDO	A	701	-	3,3,3	0.23	0	2,2,2	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	702	-	-	0/1/1/1	-
4	EDO	A	703	-	-	0/1/1/1	-
4	EDO	A	709	-	-	1/1/1/1	-
4	EDO	A	705	-	-	1/1/1/1	-
4	EDO	A	704	-	-	1/1/1/1	-
4	EDO	A	710	-	-	1/1/1/1	-
4	EDO	A	706	-	-	0/1/1/1	-
4	EDO	A	707	-	-	1/1/1/1	-
4	EDO	C	301	-	-	1/1/1/1	-
4	EDO	A	708	-	-	0/1/1/1	-
4	EDO	A	701	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	701	EDO	O1-C1-C2-O2
4	A	704	EDO	O1-C1-C2-O2
4	A	705	EDO	O1-C1-C2-O2
4	A	710	EDO	O1-C1-C2-O2
4	A	709	EDO	O1-C1-C2-O2
4	A	707	EDO	O1-C1-C2-O2
4	C	301	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	679/693 (97%)	0.38	63 (9%) 8 9	18, 34, 66, 108	0
2	B	225/225 (100%)	0.27	13 (5%) 23 25	22, 35, 63, 128	0
3	C	216/216 (100%)	-0.00	4 (1%) 66 70	19, 29, 51, 173	0
All	All	1120/1134 (98%)	0.28	80 (7%) 16 18	18, 33, 63, 173	0

All (80) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	215	CYS	20.6
3	C	216	SER	9.0
1	A	2	ALA	7.0
2	B	140	THR	6.7
2	B	141	SER	6.6
2	B	139	SER	6.4
2	B	142	GLY	6.3
2	B	224	SER	5.9
2	B	225	CYS	5.9
1	A	78	ASN	5.3
1	A	3	ALA	5.0
1	A	82	GLY	4.8
1	A	474	LEU	4.8
1	A	118	PHE	4.7
1	A	510	ARG	4.4
1	A	508	LEU	4.3
1	A	81	SER	4.2
1	A	479	GLN	4.1
1	A	478	GLU	4.1
1	A	256	ILE	4.0
1	A	492[A]	MET	3.8
2	B	138	LYS	3.7
1	A	275	VAL	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	484	ILE	3.7
1	A	515	VAL	3.6
1	A	273	CYS	3.6
1	A	51	ASN	3.6
1	A	584	VAL	3.6
1	A	276	PHE	3.5
2	B	200	THR	3.5
1	A	4	LEU	3.4
1	A	504	LEU	3.4
1	A	232	VAL	3.4
1	A	233	LEU	3.4
2	B	137	SER	3.4
1	A	260	TRP	3.3
1	A	268	VAL	3.3
1	A	121	GLY	3.3
1	A	552	PRO	3.3
1	A	274	TRP	3.1
1	A	502	VAL	3.1
1	A	280	LEU	3.1
1	A	120	GLN	3.0
1	A	123	ILE	3.0
2	B	66	GLY	3.0
1	A	481	PRO	2.9
1	A	122	GLY	2.9
1	A	53	ARG	2.8
2	B	136	SER	2.8
1	A	278	GLY	2.8
1	A	48	LEU	2.8
1	A	250	TRP	2.8
1	A	486	LYS	2.8
1	A	543	LEU	2.8
1	A	332	VAL	2.7
1	A	224	ILE	2.6
1	A	257	LEU	2.6
1	A	189	CYS	2.6
3	C	213	THR	2.5
1	A	545	PRO	2.5
3	C	33	TYR	2.5
2	B	167	ALA	2.5
1	A	119	SER	2.4
1	A	460	LEU	2.4
1	A	279	THR	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	277	ALA	2.4
1	A	624	PRO	2.3
1	A	231	GLY	2.3
1	A	485	GLY	2.3
1	A	410	TRP	2.2
1	A	505	LEU	2.2
2	B	16	ARG	2.2
1	A	192	ILE	2.1
1	A	483	ILE	2.1
1	A	549	ALA	2.1
1	A	654	GLY	2.1
1	A	188	ILE	2.0
1	A	49	GLY	2.0
1	A	254	VAL	2.0
1	A	489	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	EDO	A	709	4/4	0.69	0.28	65,73,74,75	1
4	EDO	A	701	4/4	0.82	0.10	43,54,55,55	1
4	EDO	A	710	4/4	0.83	0.21	44,58,59,65	1
4	EDO	A	706	4/4	0.84	0.14	57,60,63,65	1
4	EDO	A	707	4/4	0.87	0.21	47,56,59,61	1
4	EDO	A	705	4/4	0.92	0.11	60,63,64,64	1
4	EDO	A	703	4/4	0.93	0.11	48,48,59,59	1
4	EDO	A	708	4/4	0.94	0.08	45,48,55,55	1

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å²)	Q<0.9
4	EDO	C	301	4/4	0.96	0.15	43,46,55,55	1
5	CA	A	713	1/1	0.96	0.06	37,37,37,37	0
4	EDO	A	702	4/4	0.97	0.15	31,35,35,36	1
4	EDO	A	704	4/4	0.97	0.15	29,44,72,74	1
5	CA	A	712	1/1	0.99	0.07	35,35,35,35	0
5	CA	A	711	1/1	0.99	0.14	22,22,22,22	0
6	CL	A	714	1/1	0.99	0.06	46,46,46,46	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around CA A 713:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around CA A 712:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.