



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 10:58 PM UTC

PDB ID : 8OXY / pdb\_00008oxy  
Title : Transglutaminase 3 without calcium in complex with DH patient-derived Fab DH63-B02  
Authors : Heggelund, J.E.; Sollid, L.M.  
Deposited on : 2023-05-02  
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	693	<div> <div>%</div> <div> <div></div> <div>84%</div> <div>13%</div> <div>..</div> </div> </div>
2	B	225	<div> <div>2%</div> <div> <div></div> <div>92%</div> <div>7%</div> </div> </div>
3	C	216	<div> <div>%</div> <div> <div></div> <div>87%</div> <div>12%</div> <div>.</div> </div> </div>

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 18049 atoms, of which 8531 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein-glutamine gamma-glutamyltransferase E 27 kDa non-catalytic chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	678	Total	C	H	N	O	S	136	3	0
			10558	3350	5244	921	1019	24			


- Molecule 2 is a protein called Antibody fab fragment heavy chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	224	Total	C	H	N	O	S	74	2	0
			3361	1064	1661	291	338	7			

- Molecule 3 is a protein called Antibody fab fragment light chain.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	216	Total	C	H	N	O	S	74	0	0
			3186	1007	1560	272	341	6			

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C H O 10 2 6 2	1	0
4	A	1	Total C H O 10 2 6 2	1	0
4	A	1	Total C H O 10 2 6 2	1	0
4	A	1	Total C H O 10 2 6 2	1	0
4	A	1	Total C H O 10 2 6 2	1	0
4	A	1	Total C H O 10 2 6 2	1	0
4	C	1	Total C H O 10 2 6 2	1	0
4	C	1	Total C H O 10 2 6 2	1	0
4	C	1	Total C H O 10 2 6 2	1	0
4	C	1	Total C H O 10 2 6 2	1	0
4	C	1	Total C H O 10 2 6 2	1	0

- Molecule 5 is PHOSPHATE ION (CCD ID: PO4) (formula:  $O_4P$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total O P 5 4 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	486	Total 486	O 486	0	0
6	B	169	Total 169	O 169	0	0
6	C	174	Total 174	O 174	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Protein-glutamine gamma-glutamyltransferase E 27 kDa non-catalytic chain



- Molecule 2: Antibody fab fragment heavy chain



- Molecule 3: Antibody fab fragment light chain



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	80.02Å 92.28Å 90.36Å 90.00° 92.31° 90.00°	Depositor
Resolution (Å)	58.77 – 2.00 58.77 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (58.77-2.00) 97.3 (58.77-2.00)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.23 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.8.0405	Depositor
R, $R_{free}$	0.176 , 0.226 0.174 , 0.226	Depositor DCC
$R_{free}$ test set	4530 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.7	Xtriage
Anisotropy	0.623	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 50.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.016 for -h,-l,-k 0.000 for -h,l,k 0.023 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	18049	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.99% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.71	0/5425	1.09	4/7358 (0.1%)
2	B	0.71	0/1740	1.08	1/2368 (0.0%)
3	C	0.73	0/1665	1.15	2/2273 (0.1%)
All	All	0.72	0/8830	1.10	7/11999 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
3	C	0	2
All	All	0	5

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	55	ARG	CB-CA-C	7.00	117.83	109.85
2	B	202	THR	CA-CB-OG1	-6.43	99.96	109.60
1	A	365	PHE	CA-CB-CG	-6.18	107.61	113.80
1	A	159	ASP	CA-CB-CG	5.47	118.07	112.60
1	A	18	GLN	CB-CG-CD	-5.43	103.37	112.60

There are no chirality outliers.

All (5) planarity outliers are listed below:



Mol	Chain	Res	Type	Group
1	A	17	ARG	Sidechain
1	A	285	ARG	Sidechain
1	A	53	ARG	Sidechain
3	C	55	ARG	Sidechain
3	C	62	ARG	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5314	5244	5231	58	0
2	B	1700	1661	1655	12	0
3	C	1626	1560	1556	25	0
4	A	24	36	36	0	0
4	C	20	30	30	1	0
5	B	5	0	0	0	0
6	A	486	0	0	7	1
6	B	169	0	0	1	0
6	C	174	0	0	3	1
All	All	9518	8531	8508	90	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 90 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:224:SER:O	3:C:215:CYS:SG	2.37	0.82
2:B:224:SER:C	3:C:215:CYS:SG	2.72	0.72
2:B:219:ARG:HD3	2:B:221:GLU:OE2	1.91	0.71
1:A:110[A]:ARG:HG3	1:A:110[A]:ARG:HH11	1.57	0.70
1:A:110[B]:ARG:NH1	1:A:110[B]:ARG:HG2	2.08	0.69

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:873:HOH:O	6:C:536:HOH:O[2_546]	2.18	0.02

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	677/693 (98%)	659 (97%)	18 (3%)	0	100	100
2	B	224/225 (100%)	218 (97%)	5 (2%)	1 (0%)	30	27
3	C	214/216 (99%)	207 (97%)	7 (3%)	0	100	100
All	All	1115/1134 (98%)	1084 (97%)	30 (3%)	1 (0%)	48	46

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	92	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	586/595 (98%)	575 (98%)	11 (2%)	50	56
2	B	192/191 (100%)	190 (99%)	2 (1%)	68	75
3	C	189/189 (100%)	184 (97%)	5 (3%)	40	44
All	All	967/975 (99%)	949 (98%)	18 (2%)	50	56

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	57	SER
3	C	215	CYS
3	C	193	ARG
1	A	536	LYS
3	C	13	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	329	ASN
1	A	358	GLN
2	B	213	ASN
1	A	225	ASN
1	A	177	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	EDO	C	305	-	3,3,3	0.56	0	2,2,2	0.62	0
4	EDO	C	301	-	3,3,3	0.24	0	2,2,2	0.14	0
4	EDO	C	304	-	3,3,3	0.15	0	2,2,2	0.35	0
5	PO4	B	301	-	4,4,4	2.05	1 (25%)	6,6,6	0.68	0
4	EDO	A	702	-	3,3,3	0.27	0	2,2,2	0.05	0
4	EDO	A	703	-	3,3,3	0.40	0	2,2,2	0.15	0
4	EDO	A	704	-	3,3,3	0.14	0	2,2,2	0.24	0
4	EDO	C	302	-	3,3,3	0.31	0	2,2,2	0.12	0
4	EDO	A	706	-	3,3,3	0.43	0	2,2,2	0.68	0
4	EDO	A	701	-	3,3,3	0.17	0	2,2,2	0.27	0
4	EDO	A	705	-	3,3,3	0.31	0	2,2,2	0.39	0
4	EDO	C	303	-	3,3,3	0.15	0	2,2,2	0.39	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	C	305	-	-	0/1/1/1	-
4	EDO	C	301	-	-	1/1/1/1	-
4	EDO	C	304	-	-	1/1/1/1	-
4	EDO	A	702	-	-	0/1/1/1	-
4	EDO	A	703	-	-	0/1/1/1	-
4	EDO	A	704	-	-	0/1/1/1	-
4	EDO	C	302	-	-	0/1/1/1	-
4	EDO	A	706	-	-	0/1/1/1	-
4	EDO	A	701	-	-	1/1/1/1	-
4	EDO	A	705	-	-	1/1/1/1	-
4	EDO	C	303	-	-	0/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	301	PO4	P-O1	3.90	1.59	1.50

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	C	301	EDO	O1-C1-C2-O2
4	C	304	EDO	O1-C1-C2-O2
4	A	701	EDO	O1-C1-C2-O2
4	A	705	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	302	EDO	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	678/693 (97%)	-0.28	7 (1%) 79 79	12, 33, 66, 96	3 (0%)
2	B	224/225 (99%)	-0.23	4 (1%) 67 67	14, 32, 62, 122	2 (0%)
3	C	216/216 (100%)	-0.38	2 (0%) 81 80	20, 31, 54, 87	0
All	All	1118/1134 (98%)	-0.29	13 (1%) 76 76	12, 32, 64, 122	5 (0%)

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	ALA	4.3
1	A	82	GLY	3.9
3	C	96	PRO	3.0
3	C	215	CYS	3.0
1	A	476	THR	2.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	EDO	A	701	4/4	0.86	0.17	44,57,69,72	1
5	PO4	B	301	5/5	0.86	0.13	43,55,65,87	0
4	EDO	C	302	4/4	0.93	0.09	34,36,38,38	1
4	EDO	C	301	4/4	0.93	0.11	40,43,49,51	1
4	EDO	A	702	4/4	0.95	0.08	41,41,43,45	1
4	EDO	A	703	4/4	0.95	0.09	33,37,39,41	1
4	EDO	C	304	4/4	0.95	0.14	36,44,48,51	1
4	EDO	C	305	4/4	0.95	0.09	41,42,46,46	1
4	EDO	A	706	4/4	0.95	0.08	34,35,39,39	1
4	EDO	C	303	4/4	0.96	0.07	37,40,49,49	1
4	EDO	A	705	4/4	0.97	0.06	33,37,43,43	1
4	EDO	A	704	4/4	0.97	0.06	35,37,39,40	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.