



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 4, 2024 – 08:13 AM EST

PDB ID : 1P1X
Title : Comparison of class I aldolase binding site architecture based on the crystal structure of 2-deoxyribose-5-phosphate aldolase determined at 0.99 Angstrom resolution
Authors : Heine, A.; Luz, J.G.; Wong, C.H.; Wilson, I.A.
Deposited on : 2003-04-14
Resolution : 0.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 0.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1050 (1.06-0.94)
Clashscore	141614	1117 (1.06-0.94)
Ramachandran outliers	138981	1043 (1.06-0.94)
Sidechain outliers	138945	1045 (1.06-0.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	260	 89% 7%
1	B	260	 90% 6%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4455 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Deoxyribose-phosphate aldolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	250	1904	1200	332	362	10	0	3	0
1	B	251	1907	1202	332	363	10	0	4	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	HIS	-	expression tag	UNP P0A6L0
B	1000	HIS	-	expression tag	UNP P0A6L0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	326	Total 326	O 326	0	0
2	B	318	Total 318	O 318	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Deoxyribose-phosphate aldolase

Chain A:  89% 7%



- Molecule 1: Deoxyribose-phosphate aldolase

Chain B:  90% 6%



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	48.73Å 42.01Å 145.41Å 90.00° 98.53° 90.00°	Depositor
Resolution (Å)	8.00 – 0.99 37.54 – 0.99	Depositor EDS
% Data completeness (in resolution range)	92.5 (8.00-0.99) 90.7 (37.54-0.99)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.10 (at 0.99Å)	Xtrriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	(Not available) , (Not available) 0.142 , 0.159	Depositor DCC
R_{free} test set	15816 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	12.3	Xtrriage
Anisotropy	0.617	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 54.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.017 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	4455	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.75	1/1943 (0.1%)	1.33	19/2627 (0.7%)
1	B	0.73	0/1951	1.29	19/2639 (0.7%)
All	All	0.74	1/3894 (0.0%)	1.31	38/5266 (0.7%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	89	GLU	CD-OE1	-5.48	1.19	1.25

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1152	ARG	NE-CZ-NH1	15.66	128.13	120.30
1	A	108	ARG	NE-CZ-NH2	-14.86	112.87	120.30
1	B	1091	ARG	CD-NE-CZ	14.67	144.14	123.60
1	A	91[A]	ARG	NE-CZ-NH1	13.13	126.87	120.30
1	A	91[B]	ARG	NE-CZ-NH1	13.13	126.87	120.30
1	A	91[A]	ARG	CD-NE-CZ	11.94	140.32	123.60
1	A	91[B]	ARG	CD-NE-CZ	11.94	140.32	123.60
1	B	1091	ARG	NE-CZ-NH1	11.32	125.96	120.30
1	A	69	ARG	NE-CZ-NH2	10.54	125.57	120.30
1	A	57	ARG	NE-CZ-NH1	9.59	125.10	120.30
1	B	1108[A]	ARG	NE-CZ-NH1	8.01	124.31	120.30
1	B	1108[B]	ARG	NE-CZ-NH1	8.01	124.31	120.30
1	A	22	ASP	CB-CG-OD2	7.89	125.40	118.30
1	B	1023	ASP	CB-CG-OD1	7.59	125.13	118.30
1	B	1207	ARG	CD-NE-CZ	7.33	133.86	123.60
1	A	10[A]	ARG	NE-CZ-NH1	7.08	123.84	120.30
1	A	10[B]	ARG	NE-CZ-NH1	7.08	123.84	120.30
1	A	186	MET	CG-SD-CE	-6.89	89.17	100.20
1	A	165	PHE	CB-CG-CD2	-6.61	116.17	120.80
1	A	147	ASP	CB-CG-OD2	-6.58	112.38	118.30
1	A	91[A]	ARG	NE-CZ-NH2	-6.57	117.01	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	91[B]	ARG	NE-CZ-NH2	-6.57	117.01	120.30
1	A	207	ARG	NE-CZ-NH1	6.43	123.52	120.30
1	B	1010	ARG	NE-CZ-NH2	6.29	123.45	120.30
1	A	27	GLU	O-C-N	-6.13	112.89	122.70
1	B	1233	TYR	CB-CG-CD2	-6.06	117.36	121.00
1	B	1152	ARG	NE-CZ-NH2	-5.91	117.35	120.30
1	B	1171	GLY	C-N-CA	5.90	136.45	121.70
1	B	1165	PHE	CG-CD2-CE2	5.55	126.90	120.80
1	B	1171	GLY	O-C-N	-5.46	113.97	122.70
1	A	210	GLU	OE1-CD-OE2	-5.46	116.75	123.30
1	B	1108[A]	ARG	NE-CZ-NH2	-5.36	117.62	120.30
1	B	1108[B]	ARG	NE-CZ-NH2	-5.36	117.62	120.30
1	B	1091	ARG	NH1-CZ-NH2	-5.33	113.54	119.40
1	B	1210	GLU	OE1-CD-OE2	5.24	129.58	123.30
1	A	152	ARG	NE-CZ-NH2	-5.17	117.72	120.30
1	B	1102	ASP	CB-CG-OD1	5.13	122.92	118.30
1	B	1215	TYR	CB-CG-CD2	5.04	124.03	121.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1904	0	1946	3	0
1	B	1907	0	1938	3	0
2	A	326	0	0	2	0
2	B	318	0	0	2	0
All	All	4455	0	3884	6	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (6) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:214:LYS:HE2	2:A:2558:HOH:O	1.97	0.65
1:A:146:LYS:HE3	2:A:2306:HOH:O	2.03	0.58
1:A:202:PRO:HD2	1:A:234:ARG:O	2.15	0.46
1:B:1202:PRO:HD2	1:B:1234:ARG:O	2.15	0.46
1:B:1186:MET:HE1	2:B:2595:HOH:O	2.18	0.43
1:B:1146:LYS:HE3	2:B:2504:HOH:O	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	251/260 (96%)	245 (98%)	6 (2%)	0	100	100
1	B	253/260 (97%)	247 (98%)	6 (2%)	0	100	100
All	All	504/520 (97%)	492 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	198/202 (98%)	198 (100%)	0	100	100
1	B	199/202 (98%)	199 (100%)	0	100	100
All	All	397/404 (98%)	397 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	35	GLN
1	A	42	ASN
1	A	250	HIS
1	B	1035	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.