



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 12, 2024 – 10:39 AM EDT

PDB ID : 2P1J
Title : Crystal structure of a polC-type DNA polymerase III exonuclease domain from *Thermotoga maritima*
Authors : Bonanno, J.B.; Rutter, M.; Bain, K.T.; Izuka, M.; Sridhar, V.; Smith, D.; Wasserman, S.; Sauder, J.M.; Burley, S.K.; Almo, S.C.; New York SGX Research Center for Structural Genomics (NYSGXRC)
Deposited on : 2007-03-05
Resolution : 2.50 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.20.1
EDS : 2.36.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	186	 6% 58% 22% 8% • 12%
1	B	186	 5% 53% 27% 11% • 8%

2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 2809 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase III polC-type.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	164	1331	857	224	246	4	0	0	0
1	B	171	1392	894	239	255	4	0	0	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	345	MET	-	cloning artifact	UNP Q9ZHF6
A	346	SER	-	cloning artifact	UNP Q9ZHF6
A	523	GLU	-	cloning artifact	UNP Q9ZHF6
A	524	GLY	-	cloning artifact	UNP Q9ZHF6
A	525	HIS	-	cloning artifact	UNP Q9ZHF6
A	526	HIS	-	cloning artifact	UNP Q9ZHF6
A	527	HIS	-	cloning artifact	UNP Q9ZHF6
A	528	HIS	-	cloning artifact	UNP Q9ZHF6
A	529	HIS	-	cloning artifact	UNP Q9ZHF6
A	530	HIS	-	cloning artifact	UNP Q9ZHF6
B	345	MET	-	cloning artifact	UNP Q9ZHF6
B	346	SER	-	cloning artifact	UNP Q9ZHF6
B	523	GLU	-	cloning artifact	UNP Q9ZHF6
B	524	GLY	-	cloning artifact	UNP Q9ZHF6
B	525	HIS	-	cloning artifact	UNP Q9ZHF6
B	526	HIS	-	cloning artifact	UNP Q9ZHF6
B	527	HIS	-	cloning artifact	UNP Q9ZHF6
B	528	HIS	-	cloning artifact	UNP Q9ZHF6
B	529	HIS	-	cloning artifact	UNP Q9ZHF6
B	530	HIS	-	cloning artifact	UNP Q9ZHF6


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	45	Total 45	O 45	0	0
2	B	41	Total 41	O 41	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA polymerase III polC-type



- Molecule 1: DNA polymerase III polC-type



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	55.70Å 124.94Å 143.06Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.50 19.99 – 2.50	Depositor EDS
% Data completeness (in resolution range)	96.0 (20.00-2.50) 96.0 (19.99-2.50)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.42 (at 2.50Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.229 , 0.298 0.224 , 0.288	Depositor DCC
R_{free} test set	848 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	62.2	Xtrriage
Anisotropy	0.036	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 55.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2809	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.81	0/1356	0.96	2/1831 (0.1%)
1	B	0.89	1/1420 (0.1%)	0.95	2/1915 (0.1%)
All	All	0.85	1/2776 (0.0%)	0.95	4/3746 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	377	GLU	CG-CD	5.53	1.60	1.51

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	453	LEU	CA-CB-CG	8.93	135.83	115.30
1	A	361	LEU	CA-CB-CG	6.31	129.81	115.30
1	B	478	LEU	N-CA-C	5.72	126.44	111.00
1	B	501	ARG	CG-CD-NE	-5.16	100.97	111.80

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	436	ASP	Peptide
1	B	495	GLY	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1331	0	1351	47	0
1	B	1392	0	1407	57	0
2	A	45	0	0	3	0
2	B	41	0	0	10	0
All	All	2809	0	2758	99	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All (99) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:480:LEU:HD22	1:B:482:SER:O	1.40	1.21
1:A:423:ILE:H	1:A:423:ILE:CD1	1.46	1.21
1:B:366:THR:CG2	1:B:373:ASP:HB2	1.75	1.16
1:A:423:ILE:H	1:A:423:ILE:HD13	0.93	1.08
1:A:423:ILE:HD13	1:A:423:ILE:N	1.72	1.03
1:B:366:THR:HG22	1:B:373:ASP:HB2	1.03	1.02
1:B:366:THR:HG22	1:B:373:ASP:CB	1.92	0.99
1:A:434:LEU:O	1:A:435:GLU:HB2	1.65	0.97
1:A:419:ASN:HD21	1:A:420:LYS:HE3	1.30	0.95
1:B:480:LEU:CD2	1:B:482:SER:O	2.14	0.94
1:B:364:GLU:HB2	1:B:377:GLU:HB3	1.49	0.91
1:B:481:ARG:HB3	2:B:66:HOH:O	1.71	0.89
1:A:368:LEU:HD22	2:A:65:HOH:O	1.75	0.84
1:A:423:ILE:CD1	1:A:423:ILE:N	2.30	0.83
1:A:435:GLU:HA	1:A:465:ARG:NH2	1.93	0.83
1:A:475:LYS:HE3	2:A:31:HOH:O	1.81	0.80
1:A:361:LEU:HD13	1:A:439:ILE:HD12	1.66	0.77
1:B:465:ARG:HH11	1:B:465:ARG:HB2	1.49	0.77
1:B:505:ASP:O	1:B:509:THR:HG23	1.88	0.72
1:A:361:LEU:CD1	1:A:439:ILE:HD12	2.20	0.71
1:A:419:ASN:ND2	1:A:420:LYS:HE3	2.04	0.71
1:B:435:GLU:HB2	2:B:63:HOH:O	1.89	0.70
1:B:495:GLY:HA3	2:B:45:HOH:O	1.91	0.69

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:478:LEU:O	1:B:479:LYS:HB2	1.92	0.69
1:B:394:THR:HG21	1:B:421:ARG:NH1	2.07	0.69
1:A:434:LEU:O	1:A:435:GLU:CB	2.39	0.68
1:B:523:GLU:HA	2:B:78:HOH:O	1.93	0.67
1:A:519:MET:CB	1:A:520:MET:HA	2.24	0.66
1:A:519:MET:HB3	1:A:520:MET:HA	1.78	0.64
1:A:419:ASN:HD22	1:A:419:ASN:H	1.46	0.63
1:B:409:ILE:HD13	2:B:64:HOH:O	1.98	0.63
1:B:402:ILE:HD11	1:B:406:SER:HB3	1.81	0.62
1:B:486:ASP:O	1:B:490:GLU:HG3	1.98	0.62
1:A:501:ARG:HD3	2:B:16:HOH:O	1.99	0.62
1:B:427:LEU:O	1:B:431:LEU:HD23	2.01	0.60
1:A:478:LEU:O	1:A:479:LYS:HB3	2.01	0.60
1:A:486:ASP:O	1:A:490:GLU:HG3	2.02	0.59
1:B:357:THR:N	2:B:81:HOH:O	2.36	0.58
1:A:394:THR:HG21	1:A:421:ARG:NH1	2.19	0.57
1:A:391:GLU:CG	1:B:503:LEU:HD11	2.35	0.56
1:A:442:HIS:HD2	1:A:470:THR:OG1	1.87	0.56
1:A:361:LEU:HD11	1:A:439:ILE:CD1	2.36	0.56
1:B:365:THR:HG23	1:B:367:GLY:H	1.70	0.56
1:A:435:GLU:O	1:A:436:ASP:CB	2.54	0.55
1:B:508:VAL:HA	1:B:511:GLN:HE21	1.71	0.55
1:B:394:THR:HG21	1:B:421:ARG:HH12	1.72	0.55
1:B:402:ILE:CD1	1:B:406:SER:HB3	2.36	0.55
1:B:360:VAL:HB	1:B:381:VAL:HG22	1.89	0.54
1:A:435:GLU:HA	1:A:465:ARG:HH21	1.73	0.53
1:A:477:LEU:O	1:A:478:LEU:HD23	2.08	0.53
1:A:361:LEU:CD1	1:A:439:ILE:CD1	2.86	0.53
1:B:366:THR:HG21	1:B:373:ASP:HB2	1.83	0.52
1:B:421:ARG:HD3	2:B:17:HOH:O	2.10	0.52
1:A:453:LEU:HD12	1:A:454:TRP:CD1	2.45	0.52
1:A:382:LYS:NZ	1:A:433:PHE:O	2.31	0.51
1:A:419:ASN:HD22	1:A:419:ASN:N	2.07	0.51
1:B:504:ASP:O	1:B:508:VAL:HG13	2.12	0.50
1:A:368:LEU:HD12	1:A:446:PHE:CE1	2.47	0.49
1:B:402:ILE:HG12	1:B:406:SER:CB	2.43	0.49
1:B:516:PHE:O	1:B:520:MET:HG3	2.12	0.49
1:A:510:ALA:HB2	1:B:381:VAL:HG21	1.94	0.49
1:B:480:LEU:HD21	1:B:487:SER:HB3	1.94	0.48
1:B:421:ARG:NH2	1:B:429:GLU:OE1	2.47	0.47
1:B:364:GLU:HB2	1:B:377:GLU:CB	2.34	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:423:ILE:HG12	2:B:18:HOH:O	2.14	0.47
1:B:508:VAL:HA	1:B:511:GLN:NE2	2.30	0.46
1:A:377:GLU:OE2	1:B:501:ARG:HB2	2.16	0.46
1:A:421:ARG:HH22	1:A:429:GLU:CD	2.19	0.46
1:B:360:VAL:HA	1:B:440:VAL:O	2.14	0.46
1:A:382:LYS:HG2	1:A:389:VAL:CG2	2.45	0.46
1:B:442:HIS:O	1:B:443:ASN:HB3	2.16	0.45
1:B:419:ASN:CG	1:B:419:ASN:O	2.55	0.45
1:B:480:LEU:HD21	1:B:487:SER:CB	2.46	0.45
1:B:431:LEU:N	1:B:431:LEU:CD2	2.79	0.45
1:B:451:LEU:O	1:B:455:ILE:HG13	2.17	0.45
1:A:456:LYS:HE2	1:A:462:ASP:OD1	2.18	0.44
1:A:425:GLU:O	1:A:428:PRO:HD2	2.17	0.44
1:A:480:LEU:HD21	1:A:491:LYS:HD3	1.99	0.44
1:A:388:ILE:HD13	2:A:32:HOH:O	2.17	0.44
1:B:358:PHE:CD2	1:B:438:ILE:HB	2.53	0.44
1:A:361:LEU:HD11	1:A:439:ILE:HD13	1.99	0.44
1:B:437:SER:O	1:B:465:ARG:HD3	2.18	0.44
1:A:391:GLU:HG2	1:B:503:LEU:HD11	1.99	0.44
1:B:369:ASP:HA	1:B:370:PRO:HD3	1.89	0.44
1:B:431:LEU:HD11	1:B:455:ILE:HD13	2.01	0.43
1:A:446:PHE:HD2	1:A:447:ASP:OD1	2.01	0.43
1:A:435:GLU:O	1:A:436:ASP:HB2	2.18	0.42
1:B:467:TYR:CD2	1:B:467:TYR:C	2.93	0.42
1:A:442:HIS:CD2	1:A:470:THR:OG1	2.70	0.42
1:B:431:LEU:N	1:B:431:LEU:HD22	2.33	0.42
1:B:365:THR:CG2	1:B:367:GLY:O	2.68	0.41
1:A:391:GLU:HG3	1:B:503:LEU:HD11	2.02	0.41
1:B:402:ILE:HG12	1:B:406:SER:HB2	2.02	0.41
1:A:381:VAL:CG2	1:A:388:ILE:HG13	2.50	0.41
1:A:452:ARG:HH11	1:A:452:ARG:HD3	1.76	0.41
1:B:430:PHE:O	1:B:433:PHE:HB3	2.20	0.41
1:B:358:PHE:CD1	1:B:358:PHE:N	2.89	0.40
1:B:465:ARG:HG3	1:B:466:PRO:HD2	2.03	0.40
1:B:382:LYS:HE3	2:B:68:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	162/186 (87%)	149 (92%)	10 (6%)	3 (2%)	8	13
1	B	169/186 (91%)	150 (89%)	15 (9%)	4 (2%)	6	9
All	All	331/372 (89%)	299 (90%)	25 (8%)	7 (2%)	7	11

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	435	GLU
1	B	496	PRO
1	B	484	SER
1	A	436	ASP
1	B	421	ARG
1	B	456	LYS
1	A	372	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	147/166 (89%)	121 (82%)	26 (18%)	2	3
1	B	153/166 (92%)	122 (80%)	31 (20%)	1	2
All	All	300/332 (90%)	243 (81%)	57 (19%)	1	2

All (57) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	360	VAL
1	A	361	LEU
1	A	368	LEU
1	A	381	VAL
1	A	391	GLU
1	A	394	THR
1	A	400	ARG
1	A	415	GLU
1	A	419	ASN
1	A	423	ILE
1	A	424	GLU
1	A	425	GLU
1	A	431	LEU
1	A	437	SER
1	A	439	ILE
1	A	450	PHE
1	A	451	LEU
1	A	453	LEU
1	A	456	LYS
1	A	458	VAL
1	A	461	LEU
1	A	486	ASP
1	A	487	SER
1	A	501	ARG
1	A	514	LEU
1	A	519	MET
1	B	365	THR
1	B	366	THR
1	B	371	GLN
1	B	384	GLN
1	B	394	THR
1	B	402	ILE
1	B	404	ARG
1	B	419	ASN
1	B	422	SER
1	B	423	ILE
1	B	434	LEU
1	B	438	ILE
1	B	445	ASN
1	B	451	LEU
1	B	453	LEU
1	B	458	VAL
1	B	462	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	465	ARG
1	B	470	THR
1	B	479	LYS
1	B	480	LEU
1	B	482	SER
1	B	484	SER
1	B	487	SER
1	B	488	VAL
1	B	490	GLU
1	B	496	PRO
1	B	501	ARG
1	B	509	THR
1	B	514	LEU
1	B	518	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	419	ASN
1	A	442	HIS
1	A	511	GLN
1	B	445	ASN
1	B	511	GLN
1	B	526	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	164/186 (88%)	0.41	12 (7%) 15 15	45, 64, 82, 92	2 (1%)
1	B	171/186 (91%)	0.47	10 (5%) 23 24	48, 64, 79, 85	1 (0%)
All	All	335/372 (90%)	0.44	22 (6%) 18 19	45, 64, 81, 92	3 (0%)

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	520	MET	5.2
1	B	435	GLU	4.2
1	B	372	VAL	4.1
1	A	375	ILE	3.9
1	B	368	LEU	3.6
1	A	436	ASP	3.4
1	B	521	LYS	3.4
1	A	376	ILE	2.7
1	B	375	ILE	2.6
1	B	376	ILE	2.6
1	A	460	GLY	2.5
1	A	418	GLU	2.5
1	A	451	LEU	2.4
1	A	462	ASP	2.4
1	B	446	PHE	2.3
1	A	415	GLU	2.2
1	A	396	ILE	2.2
1	B	522	LYS	2.2
1	B	451	LEU	2.1
1	A	518	GLU	2.1
1	A	481	ARG	2.1
1	B	481	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.