



# wwPDB X-ray Structure Validation Summary Report

Jan 8, 2024 – 10:58 pm GMT

PDB ID : 8P4L  
Title : Beta-N-acetylgalactosaminidase from Niabella aurantiaca  
Authors : Fjermedal, S.; Wilkens, C.  
Deposited on : 2023-05-22  
Resolution : 2.79 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	562	69% 24% • 5%
1	B	562	67% 26% • 5%
1	C	562	70% 23% • 5%
1	D	562	66% 27% •• 5%
1	E	562	67% 25% • 5%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	562	<p>% 70% 24% • 5%</p>
1	G	562	<p>% 67% 26% • 5%</p>
1	H	562	<p>% 67% 25% • 5%</p>
1	I	562	<p>2% 71% 22% • 5%</p>
1	J	562	<p>4% 66% 27% • 5%</p>
1	K	562	<p>2% 67% 26% • 5%</p>
1	L	562	<p>6% 63% 29% • 5%</p>

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 53723 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-N-acetylgalactosaminidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	536	4455	2896	753	788	18	0	0	0
1	B	536	4466	2906	754	788	18	0	1	0
1	C	536	4455	2896	753	788	18	0	0	0
1	D	536	4466	2906	754	788	18	0	1	0
1	E	536	4455	2896	753	788	18	0	0	0
1	F	536	4455	2896	753	788	18	0	0	0
1	G	536	4455	2896	753	788	18	0	0	0
1	H	536	4455	2896	753	788	18	0	0	0
1	I	536	4455	2896	753	788	18	0	0	0
1	J	536	4455	2896	753	788	18	0	0	0
1	K	536	4455	2896	753	788	18	0	0	0
1	L	536	4455	2896	753	788	18	0	0	0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	40	Total	O	0	0
			40	40		
2	B	18	Total	O	0	0
			18	18		

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>	<b>ZeroOcc</b>	<b>AltConf</b>
2	C	39	Total O 39 39	0	0
2	D	28	Total O 28 28	0	0
2	E	30	Total O 30 30	0	0
2	F	26	Total O 26 26	0	0
2	G	14	Total O 14 14	0	0
2	H	14	Total O 14 14	0	0
2	I	7	Total O 7 7	0	0
2	J	13	Total O 13 13	0	0
2	K	7	Total O 7 7	0	0
2	L	5	Total O 5 5	0	0


### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Beta-N-acetylgalactosaminidase




- Molecule 1: Beta-N-acetylgalactosaminidase






● Molecule 1: Beta-N-acetylgalactosaminidase




● Molecule 1: Beta-N-acetylgalactosaminidase




- Molecule 1: Beta-N-acetylgalactosaminidase

Chain E:  67% 25% 5%




- Molecule 1: Beta-N-acetylgalactosaminidase

Chain F:  70% 24% 5%




- Molecule 1: Beta-N-acetylgalactosaminidase


Chain G:  67% 26% 5%








• Molecule 1: Beta-N-acetylgalactosaminidase




• Molecule 1: Beta-N-acetylgalactosaminidase






● Molecule 1: Beta-N-acetylgalactosaminidase




● Molecule 1: Beta-N-acetylgalactosaminidase





• Molecule 1: Beta-N-acetylgalactosaminidase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	135.46Å 117.62Å 282.44Å 90.00° 97.74° 90.00°	Depositor
Resolution (Å)	47.02 – 2.79 47.02 – 2.79	Depositor EDS
% Data completeness (in resolution range)	98.4 (47.02-2.79) 98.7 (47.02-2.79)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.40 (at 2.81Å)	Xtrriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, $R_{free}$	0.193 , 0.259 0.193 , 0.258	Depositor DCC
$R_{free}$ test set	2263 reflections (1.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	60.3	Xtrriage
Anisotropy	0.687	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 34.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	53723	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	76.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.51% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.48	0/4608	0.62	1/6269 (0.0%)
1	B	0.45	0/4624	0.62	1/6292 (0.0%)
1	C	0.52	0/4608	0.65	0/6269
1	D	0.48	0/4624	0.66	2/6292 (0.0%)
1	E	0.49	1/4608 (0.0%)	0.63	0/6269
1	F	0.47	0/4608	0.62	0/6269
1	G	0.43	0/4608	0.59	0/6269
1	H	0.45	0/4608	0.62	0/6269
1	I	0.42	0/4608	0.59	0/6269
1	J	0.41	0/4608	0.61	0/6269
1	K	0.39	0/4608	0.60	1/6269 (0.0%)
1	L	0.38	0/4608	0.58	0/6269
All	All	0.45	1/55328 (0.0%)	0.62	5/75274 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	205	CYS	CB-SG	-6.34	1.71	1.82

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	368	LEU	CA-CB-CG	8.12	133.97	115.30
1	A	115	ALA	C-N-CA	6.05	136.83	121.70
1	K	7	ASP	C-N-CA	5.19	134.67	121.70
1	B	93	LEU	CA-CB-CG	5.12	127.09	115.30
1	D	518	LEU	CA-CB-CG	5.12	127.07	115.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4455	0	4312	92	0
1	B	4466	0	4322	86	0
1	C	4455	0	4312	92	0
1	D	4466	0	4322	107	0
1	E	4455	0	4311	94	0
1	F	4455	0	4312	83	0
1	G	4455	0	4312	93	0
1	H	4455	0	4312	98	0
1	I	4455	0	4312	76	0
1	J	4455	0	4312	110	0
1	K	4455	0	4312	102	0
1	L	4455	0	4312	117	0
2	A	40	0	0	2	0
2	B	18	0	0	0	0
2	C	39	0	0	5	0
2	D	28	0	0	0	0
2	E	30	0	0	0	0
2	F	26	0	0	2	0
2	G	14	0	0	4	0
2	H	14	0	0	2	0
2	I	7	0	0	1	0
2	J	13	0	0	1	0
2	K	7	0	0	0	0
2	L	5	0	0	0	0
All	All	53723	0	51763	1134	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 1134 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:237:ILE:HB	1:C:249:ASP:HB3	1.51	0.92
1:I:237:ILE:HB	1:I:249:ASP:HB3	1.50	0.92
1:F:232:PRO:HD2	1:F:293:TYR:HD2	1.33	0.91

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:143:ARG:HH21	1:A:156:LEU:HD21	1.35	0.90
1:H:237:ILE:HB	1:H:249:ASP:HB3	1.55	0.89

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	534/562 (95%)	507 (95%)	23 (4%)	4 (1%)	22	53
1	B	535/562 (95%)	504 (94%)	27 (5%)	4 (1%)	22	53
1	C	534/562 (95%)	505 (95%)	24 (4%)	5 (1%)	17	46
1	D	535/562 (95%)	509 (95%)	20 (4%)	6 (1%)	14	41
1	E	534/562 (95%)	502 (94%)	25 (5%)	7 (1%)	12	36
1	F	534/562 (95%)	506 (95%)	22 (4%)	6 (1%)	14	41
1	G	534/562 (95%)	507 (95%)	22 (4%)	5 (1%)	17	46
1	H	534/562 (95%)	509 (95%)	21 (4%)	4 (1%)	22	53
1	I	534/562 (95%)	505 (95%)	24 (4%)	5 (1%)	17	46
1	J	534/562 (95%)	507 (95%)	23 (4%)	4 (1%)	22	53
1	K	534/562 (95%)	494 (92%)	32 (6%)	8 (2%)	10	33
1	L	534/562 (95%)	500 (94%)	30 (6%)	4 (1%)	22	53
All	All	6410/6744 (95%)	6055 (94%)	293 (5%)	62 (1%)	15	44

5 of 62 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	372	ARG
1	C	339	GLU
1	C	405	THR

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	F	405	THR
1	H	30	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	474/494 (96%)	444 (94%)	30 (6%)	18 46
1	B	475/494 (96%)	432 (91%)	43 (9%)	9 27
1	C	474/494 (96%)	449 (95%)	25 (5%)	22 54
1	D	475/494 (96%)	440 (93%)	35 (7%)	13 37
1	E	474/494 (96%)	437 (92%)	37 (8%)	12 35
1	F	474/494 (96%)	445 (94%)	29 (6%)	18 48
1	G	474/494 (96%)	442 (93%)	32 (7%)	16 42
1	H	474/494 (96%)	439 (93%)	35 (7%)	13 37
1	I	474/494 (96%)	445 (94%)	29 (6%)	18 48
1	J	474/494 (96%)	443 (94%)	31 (6%)	17 44
1	K	474/494 (96%)	449 (95%)	25 (5%)	22 54
1	L	474/494 (96%)	438 (92%)	36 (8%)	13 36
All	All	5690/5928 (96%)	5303 (93%)	387 (7%)	16 42

5 of 387 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	16	SER
1	I	366	ASP
1	H	251	SER
1	H	494	SER
1	J	114	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:



Mol	Chain	Res	Type
1	L	28	GLN
1	K	234	GLN
1	J	498	GLN
1	C	234	GLN
1	K	46	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	536/562 (95%)	-0.06	4 (0%) 87 84	42, 61, 89, 120	0
1	B	536/562 (95%)	0.19	9 (1%) 70 63	49, 70, 100, 197	0
1	C	536/562 (95%)	0.01	4 (0%) 87 84	38, 56, 88, 133	0
1	D	536/562 (95%)	0.00	6 (1%) 80 75	41, 61, 93, 114	0
1	E	536/562 (95%)	0.02	2 (0%) 92 91	44, 63, 93, 154	0
1	F	536/562 (95%)	0.05	6 (1%) 80 75	48, 65, 93, 131	0
1	G	536/562 (95%)	0.13	7 (1%) 77 72	53, 74, 112, 151	0
1	H	536/562 (95%)	0.13	8 (1%) 73 68	50, 74, 115, 147	0
1	I	536/562 (95%)	0.26	11 (2%) 63 54	56, 77, 112, 170	0
1	J	536/562 (95%)	0.40	21 (3%) 39 29	58, 87, 119, 159	0
1	K	536/562 (95%)	0.37	14 (2%) 56 46	64, 93, 127, 166	0
1	L	536/562 (95%)	0.50	33 (6%) 20 13	72, 101, 132, 197	0
All	All	6432/6744 (95%)	0.17	125 (1%) 66 59	38, 72, 115, 197	0

The worst 5 of 125 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	373	TYR	11.2
1	L	373	TYR	10.5
1	I	373	TYR	8.9
1	J	373	TYR	7.3
1	C	373	TYR	7.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.