



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 12, 2024 – 11:13 AM EDT

PDB ID : 2P5Z  
Title : The E. coli c3393 protein is a component of the type VI secretion system and exhibits structural similarity to T4 bacteriophage tail proteins gp27 and gp5  
Authors : Ramagopal, U.A.; Bonanno, J.B.; Sridhar, V.; Lau, C.; Toro, R.; Gheyi, T.; Maletic, M.; Freeman, J.C.; Sauder, J.M.; Burley, S.K.; Almo, S.C.; New York SGX Research Center for Structural Genomics (NYSGXRC)  
Deposited on : 2007-03-16  
Resolution : 2.60 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.20.1  
EDS : 2.36.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	X	491	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2962 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Type VI secretion system component.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	X	361	2900	1842	508	544	6	0	1	0

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
X	-1	MET	-	cloning artifact	UNP Q8FED2
X	0	SER	-	cloning artifact	UNP Q8FED2
X	1	LEU	-	cloning artifact	UNP Q8FED2
X	482	GLU	-	cloning artifact	UNP Q8FED2
X	483	GLY	-	cloning artifact	UNP Q8FED2
X	484	HIS	-	cloning artifact	UNP Q8FED2
X	485	HIS	-	cloning artifact	UNP Q8FED2
X	486	HIS	-	cloning artifact	UNP Q8FED2
X	487	HIS	-	cloning artifact	UNP Q8FED2
X	488	HIS	-	cloning artifact	UNP Q8FED2
X	489	HIS	-	cloning artifact	UNP Q8FED2


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	X	62	Total	O	0	0
			62	62		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Type VI secretion system component



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	116.35Å 116.35Å 80.58Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.19 – 2.60 47.17 – 2.61	Depositor EDS
% Data completeness (in resolution range)	99.9 (47.19-2.60) 99.9 (47.17-2.61)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.31 (at 2.61Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.217 , 0.264 0.215 , 0.260	Depositor DCC
$R_{free}$ test set	983 reflections (5.15%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	61.4	Xtrriage
Anisotropy	0.172	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 50.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.046 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	2962	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	60.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.61% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	X	0.52	0/2958	0.85	10/3998 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	X	0	4

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	X	212	LEU	C-N-CD	-12.72	92.62	120.60
1	X	212	LEU	C-N-CA	8.23	156.58	122.00
1	X	358	LEU	CA-CB-CG	7.71	133.04	115.30
1	X	411	LEU	CA-CB-CG	6.80	130.95	115.30
1	X	115	LEU	CA-CB-CG	5.97	129.02	115.30
1	X	212	LEU	CA-CB-CG	-5.83	101.88	115.30
1	X	375	LEU	CA-CB-CG	5.67	128.33	115.30
1	X	445	LEU	CA-CB-CG	5.40	127.72	115.30
1	X	447	GLY	N-CA-C	-5.06	100.46	113.10
1	X	213	PRO	CA-N-CD	-5.04	104.44	111.50

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	X	212	LEU	Peptide
1	X	213	PRO	Peptide
1	X	294	ALA	Peptide

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Group
1	X	355	SER	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	X	2900	0	2858	27	0
2	X	62	0	0	1	0
All	All	2962	0	2858	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:135:LYS:NZ	1:X:422:TYR:OH	2.23	0.69
1:X:293:GLY:HA3	1:X:294:ALA:HB3	1.87	0.56
1:X:293:GLY:CA	1:X:294:ALA:HB3	2.39	0.52
1:X:181:GLY:O	1:X:372:ARG:NH1	2.45	0.49
1:X:213:PRO:O	1:X:214:LEU:HB2	2.11	0.49
1:X:445:LEU:H	1:X:446:ALA:HB2	1.78	0.49
1:X:446:ALA:HA	1:X:448:THR:HG22	1.94	0.48
1:X:405:GLY:N	1:X:406:ARG:HB2	2.28	0.48
1:X:59:VAL:HG13	1:X:89:ILE:HG21	1.96	0.47
1:X:293:GLY:HA3	1:X:295:PHE:HB3	1.97	0.47
1:X:203:GLN:H	1:X:203:GLN:HG2	1.20	0.47
1:X:146:GLN:NE2	2:X:517:HOH:O	2.46	0.47
1:X:203:GLN:NE2	1:X:366:SER:O	2.48	0.47
1:X:143:MET:HE1	1:X:197:ILE:HD12	1.97	0.46
1:X:293:GLY:CA	1:X:294:ALA:CB	2.93	0.46
1:X:297:ALA:HA	1:X:300:ARG:HG2	1.99	0.45
1:X:31:PHE:HE2	1:X:348:THR:CG2	2.30	0.45
1:X:396:ASP:HA	1:X:397:ILE:HA	1.82	0.44
1:X:156:TYR:HA	1:X:157:PRO:HD3	1.85	0.43
1:X:31:PHE:HE2	1:X:348:THR:HG22	1.84	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:320:LEU:HD11	1:X:326:ILE:HD13	2.02	0.41
1:X:419:LYS:HA	1:X:420:PRO:HD3	1.96	0.41
1:X:455:GLU:HB3	1:X:461:ARG:HD3	2.03	0.41
1:X:115:LEU:HD22	1:X:141:HIS:CD2	2.55	0.41
1:X:118:SER:OG	1:X:169:ASP:OD2	2.38	0.41
1:X:456:GLU:OE2	1:X:461:ARG:NH1	2.54	0.41
1:X:127:GLN:HB2	1:X:132:ILE:HG13	2.04	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	X	348/491 (71%)	320 (92%)	22 (6%)	6 (2%)	<b>9</b>   <b>18</b>

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	X	213	PRO
1	X	214	LEU
1	X	294	ALA
1	X	399	ALA
1	X	295	PHE
1	X	466	GLY

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was



analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	X	307/417 (74%)	276 (90%)	31 (10%)	7 14

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	X	17	LEU
1	X	39	THR
1	X	50	SER
1	X	67	SER
1	X	68	LEU
1	X	82	LEU
1	X	84	THR
1	X	85	LEU
1	X	88	VAL
1	X	115	LEU
1	X	180	VAL
1	X	203	GLN
1	X	213	PRO
1	X	228	VAL
1	X	300	ARG
1	X	305	LEU
1	X	325	GLU
1	X	327	LYS
1	X	331	ASP
1	X	336	VAL
1	X	348	THR
1	X	349	SER
1	X	355	SER
1	X	358	LEU
1	X	368	ARG
1	X	384	THR
1	X	409	VAL
1	X	411	LEU
1	X	424	SER
1	X	425	LEU
1	X	448	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	X	308	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	X	361/491 (73%)	0.26	10 (2%) 53 46	34, 55, 95, 112	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	X	333	ALA	4.2
1	X	330	GLY	3.1
1	X	400	HIS	2.6
1	X	332	ASP	2.6
1	X	428	ARG	2.5
1	X	401	ILE	2.4
1	X	405	GLY	2.3
1	X	402	ASP	2.2
1	X	406	ARG	2.1
1	X	369	TYR	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.