

# wwPDB X-ray Structure Validation Summary Report (i)

#### May 15, 2020 - 10:55 am BST

PDB ID	:	6P8X
$\operatorname{Title}$	:	Crystal structure of human KRAS G12C covalently bound to an acryloylaze-
		tidine acetamide inhibitor.
Authors	:	Mohr, C.
Deposited on	:	2019-06-08
Resolution	:	2.11 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.


The following versions of software and data (see references (1)) were used in the production of this report:

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.11 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries},{ m resolution\ range}({ m \AA}))$
$R_{free}$	130704	6241 (2.14-2.10)
Clashscore	141614	6778 (2.14-2.10)
Ramachandran outliers	138981	6705(2.14-2.10)
Sidechain outliers	138945	6706 (2.14-2.10)
RSRZ outliers	127900	6112 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain			
1	А	183	79%	11%	•	9%
1	В	183	4% 69%	20%	•	9%
1	С	183	3% 77%	14%		9%
1	D	183	83%	8%	·	9%



# 2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 5781 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	166	Total	С	Ν	Ο	S	0	0 0	
	A	100	1328	833	228	263	4	0	0	0
1	В	166	Total	С	Ν	Ο	S	0	1	0
	D	100	1334	837	229	264	4	0	1	0
1	С	167	Total	С	Ν	Ο	S	0	0	0
	U	107	1336	838	229	264	5	0	0	0
1	п	167	Total	С	Ν	Ο	S	0	0	0
		107	1336	838	229	264	5			U

• Molecule 1 is a protein called GTPase KRas.

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	-13	MET	-	expression tag	UNP P01116
A	-12	LYS	-	expression tag	UNP P01116
A	-11	HIS	-	expression tag	UNP P01116
А	-10	HIS	-	expression tag	UNP P01116
A	-9	HIS	-	expression tag	UNP P01116
A	-8	HIS	-	expression tag	UNP P01116
А	-7	HIS	-	expression tag	UNP P01116
A	-6	HIS	-	expression tag	UNP P01116
А	-5	HIS	-	expression tag	UNP P01116
A	-4	ASP	P - expression		UNP P01116
А	-3	-3 GLU - expres		expression tag	UNP P01116
А	-2	VAL	-	expression tag	UNP P01116
А	-1	ASP	-	expression tag	UNP P01116
А	0	GLY	-	expression tag	UNP P01116
A	12	CYS	GLY	variant	UNP P01116
A	51	SER	CYS	engineered mutation	UNP P01116
A	80	LEU	CYS	engineered mutation	UNP P01116
А	118 SER CYS		engineered mutation	UNP P01116	
В	B -13 MET -		expression tag	UNP P01116	
В	-12	LYS	-	expression tag	UNP P01116
В	-11	HIS	-	expression tag	UNP P01116



6I	28X
· · ·	· · ·

В -10 Н	lelled Actual	Comment	Reference
	TO		
B -9 H	IS –	expression tag	UNP P01116
	IS -	expression tag	UNP P01116
В -8 Н	IS -	expression tag	UNP P01116
В -7 Н	IS -	expression tag	UNP P01116
В -6 Н	IS -	expression tag	UNP P01116
В -5 Н	IS -	expression tag	UNP P01116
B -4 A	SP -	expression tag	UNP P01116
B -3 G	LU -	expression tag	UNP P01116
B -2 V.	AL -	expression tag	UNP P01116
В -1 А	SP -	expression tag	UNP P01116
B 0 G	LY -	expression tag	UNP P01116
B 12 C	YS GLY	variant	UNP P01116
B 51 SI	ER CYS	engineered mutation	UNP P01116
	EU CYS	engineered mutation	UNP P01116
B 118 S	ER CYS	engineered mutation	UNP P01116
С -13 М	ET -	expression tag	UNP P01116
C -12 L	YS -	expression tag	UNP P01116
С -11 Н	IS -	expression tag	UNP P01116
С -10 Н	IS -	expression tag	UNP P01116
С -9 Н	IS -	expression tag	UNP P01116
С -8 Н	IS -	expression tag	UNP P01116
С -7 Н	IS -	expression tag	UNP P01116
С -6 Н	IS -	expression tag	UNP P01116
С -5 Н	IS -	expression tag	UNP P01116
C -4 A	SP -	expression tag	UNP P01116
C -3 G	LU -	expression tag	UNP P01116
C -2 V.	AL -	expression tag	UNP P01116
С -1 А	SP -	expression tag	UNP P01116
C 0 G	LY -	expression tag	UNP P01116
C 12 C	YS GLY	variant	UNP P01116
C 51 SI	ER CYS	engineered mutation	UNP P01116
C 80 L	EU CYS	engineered mutation	UNP P01116
C 118 SI	ER CYS	engineered mutation	UNP P01116
D -13 M	ET -	expression tag	UNP P01116
D -12 L	YS -	expression tag	UNP P01116
D -11 H	IS -	expression tag	UNP P01116
D -10 H	IS -	expression tag	UNP P01116
D -9 H	IS -	expression tag	UNP P01116
D -8 H	IS -	expression tag	UNP P01116
D -7 H	IS -	expression tag	UNP P01116
D -6 H	IS -	expression tag	UNP P01116
D -5 H	IS -	expression tag	UNP P01116

Continued from previous page...



6P8X	
------	--


Chain	Residue	Modelled	Actual	Comment	Reference
D	-4	ASP	-	expression tag	UNP P01116
D	-3	GLU	-	expression tag	UNP P01116
D	-2	VAL	-	expression tag	UNP P01116
D	-1	ASP	-	expression tag	UNP P01116
D	0	GLY	-	expression tag	UNP P01116
D	12	CYS	GLY	variant	UNP P01116
D	51	SER	CYS	engineered mutation	UNP P01116
D	80	LEU	CYS	engineered mutation	UNP P01116
D	118	SER	CYS	engineered mutation	UNP P01116

Continued from previous page...

• Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	В	3	Total Ca 3 3	0	0
2	А	3	Total Ca 3 3	0	0
2	D	2	Total Ca 2 2	0	0
2	С	2	Total Ca 2 2	0	0


• Molecule 3 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula:  $C_{10}H_{15}N_5O_{11}P_2$ ).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	Λ	1	Total	С	Ν	Ο	Р	0	Ο
J	Л	T	28	10	5	11	2	0	0
3	B	1	Total	С	Ν	Ο	Р	0	Ο
J	D	T	28	10	5	11	2	0	0
3	С	1	Total	С	Ν	Ο	Р	0	0
0	U	I	28	10	5	11	2	0	0
3	Л	1	Total	С	Ν	Ο	Р	0	Ο
	D		28	10	5	11	2	0	0

• Molecule 4 is 2-[4-bromo-2-(3-phenyl-2,5-dihydro-1H-pyrrole-1-carbonyl)phenoxy]-N-(1-pro panoylazetidin-3-yl)acetamide (three-letter code: O5V) (formula: C<sub>25</sub>H<sub>26</sub>BrN<sub>3</sub>O<sub>4</sub>) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
4	Δ	1	Total	Br	С	Ν	Ο	0	0
4	А	1	33	1	25	3	4	0	0
4	В	1	Total	Br	С	Ν	Ο	0	0
4	D	I	33	1	25	3	4	0	0
4	С	1	Total	Br	С	Ν	Ο	0	0
4	U	1	33	1	25	3	4	0	0
4	Л	1	Total	Br	С	Ν	0	0	0
4	D	1	33	1	25	3	4	0	0

• Molecule 5 is water.

$\begin{bmatrix} 5 & A & 53 & Total & O & 0 \\ 53 & 53 & 53 & 0 & 0 \end{bmatrix}$	Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
	5	А	53	Total O 53 53	0	0




Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	В	42	$\begin{array}{cc} \text{Total} & \text{O} \\ 42 & 42 \end{array}$	0	0
5	С	40	Total         O           40         40	0	0
5	D	58	Total         O           58         58	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: GTPase KRas

## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants	33.33Å 66.15Å 78.63Å	Deperitor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$66.48^{\circ}$ $78.75^{\circ}$ $89.47^{\circ}$	Depositor
Resolution (Å)	29.86 - 2.11	Depositor
Resolution (A)	29.86 - 2.11	EDS
% Data completeness	96.7 (29.86-2.11)	Depositor
(in resolution range)	96.8(29.86-2.11)	EDS
R <sub>merge</sub>	(Not available)	Depositor
R <sub>sym</sub>	0.15	Depositor
$< I/\sigma(I) > 1$	$1.86 (at 2.10 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
D D	0.265 , $0.313$	Depositor
$R, R_{free}$	0.269 , $0.312$	DCC
$R_{free}$ test set	1594 reflections $(4.73\%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	29.8	Xtriage
Anisotropy	0.050	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.33, $32.6$	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.45, < L^2 > = 0.28$	Xtriage
	0.000 for h,-k,h-l	
Estimated twinning fraction	0.000 for -h,k,k-l	Xtriage
	0.000 for -h,-k,-h-k+l	
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5781	wwPDB-VP
Average B, all atoms $(Å^2)$	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 89.55 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.2658e-08. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, CA,  $\rm O5V$ 

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles	
	Cham	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	А	0.50	0/1349	0.72	0/1821
1	В	0.52	0/1358	0.74	0/1833
1	С	0.50	0/1357	0.71	0/1831
1	D	0.51	0/1357	0.70	0/1831
All	All	0.51	0/5421	0.72	0/7316

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1328	0	1309	26	0
1	В	1334	0	1317	38	0
1	С	1336	0	1321	22	0
1	D	1336	0	1321	8	0
2	А	3	0	0	0	0
2	В	3	0	0	0	0
2	С	2	0	0	0	0
2	D	2	0	0	0	0
3	А	28	0	12	0	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	В	28	0	12	0	0
3	С	28	0	12	5	0
3	D	28	0	12	0	0
4	А	33	0	0	0	0
4	В	33	0	0	1	0
4	С	33	0	0	3	0
4	D	33	0	0	0	0
5	А	53	0	0	2	0
5	В	42	0	0	4	0
5	С	40	0	0	2	0
5	D	58	0	0	0	0
All	All	5781	0	5316	90	0

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 90 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:48:GLY:O	1:B:49:GLU:HG2	1.46	1.15
1:B:41:ARG:HG3	1:B:54:ASP:OD2	1.64	0.97
1:B:41:ARG:CG	1:B:54:ASP:OD2	2.20	0.90
1:C:139:ILE:HG22	5:C:404:HOH:O	1.74	0.87
1:B:41:ARG:HG3	1:B:54:ASP:CG	1.97	0.84

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	164/183~(90%)	154~(94%)	10~(6%)	0	100 100



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	$\mathbf{ntiles}$
1	В	165/183~(90%)	154~(93%)	11 (7%)	0	100	100
1	С	165/183~(90%)	158 (96%)	7 (4%)	0	100	100
1	D	165/183~(90%)	157~(95%)	8 (5%)	0	100	100
All	All	659/732~(90%)	623 (94%)	36 (6%)	0	100	100

Continued from previous page...

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	147/163~(90%)	143~(97%)	4(3%)	44 47
1	В	148/163~(91%)	140~(95%)	8 (5%)	22 19
1	С	148/163~(91%)	139~(94%)	9 (6%)	18 15
1	D	148/163~(91%)	142~(96%)	6 (4%)	30 30
All	All	591/652~(91%)	564~(95%)	27~(5%)	27 25

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	С	12	CYS
1	С	38	ASP
1	D	123	ARG
1	С	31	GLU
1	В	12	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type	
1	D	150	GLN	



#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

#### 5.6 Ligand geometry (i)

Of 18 ligands modelled in this entry, 10 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Res	Link	B	ond leng	gths	B	ond ang	les		
	Type	Chain	nes	nes	nes		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GDP	С	303	2	$24,\!30,\!30$	1.18	2 (8%)	31,47,47	1.91	<mark>6 (19%)</mark>		
3	GDP	В	304	2	$24,\!30,\!30$	1.12	2 (8%)	31,47,47	1.89	7 (22%)		
3	GDP	А	304	2	24,30,30	1.14	2 (8%)	31,47,47	1.88	7 (22%)		
3	GDP	D	303	2	24,30,30	1.10	2 (8%)	31,47,47	1.89	7 (22%)		
4	O5V	А	305	1	34,36,36	1.93	10 (29%)	$39,\!50,\!50$	1.85	7 (17%)		
4	O5V	D	304	1	34,36,36	2.09	11 (32%)	$39,\!50,\!50$	1.95	7 (17%)		
4	O5V	С	304	1	34,36,36	1.92	7 (20%)	$39,\!50,\!50$	1.32	<mark>5 (12%)</mark>		
4	O5V	В	305	1	34,36,36	2.13	9 (26%)	$39,\!50,\!50$	1.37	2(5%)		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GDP	С	303	2	-	1/12/32/32	0/3/3/3
3	GDP	В	304	2	-	1/12/32/32	0/3/3/3
3	GDP	А	304	2	-	5/12/32/32	0/3/3/3
3	GDP	D	303	2	-	1/12/32/32	0/3/3/3
4	O5V	А	305	1	-	4/25/44/44	0/4/4/4
4	O5V	D	304	1	-	4/25/44/44	0/4/4/4
4	O5V	С	304	1	-	2/25/44/44	0/4/4/4
4	O5V	В	305	1	-	4/25/44/44	0/4/4/4

The worst 5 of 45 bond length outliers are listed below:

Mol	Chain	$\mathbf{Res}$	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	В	305	O5V	C10-N09	6.46	1.47	1.34
4	В	305	O5V	C21-N23	5.41	1.46	1.34
4	С	304	O5V	C21-N23	5.33	1.46	1.34
4	D	304	O5V	C03-N05	5.31	1.46	1.35
4	В	305	O5V	C03-N05	5.21	1.45	1.35

The worst 5 of 48 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
4	D	304	O5V	C24-C25-C26	-7.10	107.09	112.33
4	А	305	O5V	C24-C25-C26	-6.88	107.25	112.33
4	В	305	O5V	C24-C25-C26	-5.88	107.99	112.33
4	D	304	O5V	C25-C24-N23	5.29	105.90	102.03
3	С	303	GDP	C2-N3-C4	5.22	121.32	115.36

There are no chirality outliers.

5 of 22 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	С	303	GDP	PA-O3A-PB-O2B
3	А	304	GDP	C5'-O5'-PA-O3A
4	С	304	O5V	C01-C02-C03-O04
4	С	304	O5V	C01-C02-C03-N05
3	D	303	GDP	PA-O3A-PB-O2B


There are no ring outliers.

3 monomers are involved in 9 short contacts:




Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	С	303	GDP	5	0
4	С	304	O5V	3	0
4	В	305	O5V	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

























## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\# RSRZ {>}2$	$OWAB(Å^2)$	Q<0.9
1	А	166/183~(90%)	0.44	9 (5%) 25 31	19, 31, 51, 62	0
1	В	166/183~(90%)	0.42	8 (4%) 30 35	20, 32, 50, 62	0
1	С	167/183~(91%)	0.41	6 (3%) 42 49	18, 30, 44, 57	0
1	D	167/183~(91%)	0.38	5 (2%) 50 56	19, 30, 45, 56	0
All	All	666/732~(90%)	0.41	28 (4%) 36 42	18, 31, 49, 62	0

The worst 5 of 28 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	50	THR	5.6
1	С	1	MET	5.5
1	В	121	PRO	5.0
1	А	122	SER	3.4
1	А	56	LEU	3.2

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




6P8X
------


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} extsf{-}\mathbf{factors}(\mathbf{A}^2)$	Q<0.9
2	CA	А	303	1/1	0.75	0.10	$40,\!40,\!40,\!40$	0
4	O5V	D	304	33/33	0.85	0.16	21,31,37,46	0
4	O5V	С	304	33/33	0.88	0.17	$31,\!36,\!47,\!48$	0
4	O5V	В	305	33/33	0.90	0.15	$30,\!36,\!41,\!43$	0
2	CA	D	302	1/1	0.91	0.08	$20,\!20,\!20,\!20$	0
4	O5V	А	305	33/33	0.92	0.14	22,35,41,41	0
2	CA	В	303	1/1	0.93	0.17	$37,\!37,\!37,\!37$	0
3	GDP	С	303	28/28	0.94	0.11	$18,\!22,\!26,\!29$	0
2	CA	В	302	1/1	0.94	0.08	$26,\!26,\!26,\!26$	0
3	GDP	D	303	28/28	0.95	0.09	$21,\!23,\!25,\!29$	0
2	CA	С	302	1/1	0.95	0.07	21,21,21,21	0
3	GDP	В	304	28/28	0.95	0.11	22,27,32,34	0
3	GDP	А	304	28/28	0.96	0.09	$23,\!26,\!28,\!29$	0
2	CA	А	302	1/1	0.97	0.09	$30,\!30,\!30,\!30$	0
2	CA	С	301	1/1	0.98	0.06	$26,\!26,\!26,\!26$	0
2	CA	D	301	1/1	0.99	0.04	21,21,21,21	0
2	CA	В	301	1/1	0.99	0.07	23,23,23,23	0
2	CA	А	301	1/1	0.99	0.03	22,22,22,22	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.































## 6.5 Other polymers (i)

There are no such residues in this entry.

