



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 22, 2022 – 12:42 am GMT

PDB ID : 7P8H
Title : chicken GRIFIN bound to blood group tetrasaccharide B (type 1)
Authors : Ruiz, F.M.; Romero, A.
Deposited on : 2021-07-22
Resolution : 1.13 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.26
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.26


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.13 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1168 (1.14-1.10)
Clashscore	141614	1205 (1.14-1.10)
Ramachandran outliers	138981	1168 (1.14-1.10)
Sidechain outliers	138945	1165 (1.14-1.10)
RSRZ outliers	127900	1146 (1.14-1.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	139	 83% 11% 6%
1	C	139	 81% 93% 6%
2	E	4	 50% 50%
2	F	4	 50% 50%

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 5339 atoms, of which 2413 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Galectin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	138	2331	760	1160	194	212	5	0	9	0
1	C	138	2342	763	1163	194	217	5	0	10	0

- Molecule 2 is an oligosaccharide called alpha-L-fucopyranose-(1-2)-[alpha-D-galactopyranose-(1-3)]beta-D-galactopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
2	E	4	92	26	45	1	20	0	0	0
2	F	4	92	26	45	1	20	0	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	242	Total	O	0	0
			242	242		
3	C	240	Total	O	0	0
			240	240		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Galectin



- Molecule 1: Galectin



- Molecule 2: alpha-L-fucopyranose-(1-2)-[alpha-D-galactopyranose-(1-3)]beta-D-galactopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: alpha-L-fucopyranose-(1-2)-[alpha-D-galactopyranose-(1-3)]beta-D-galactopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  50% 50%

MAN1
GAL2
FUC3
GLA4

4 Data and refinement statistics

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, α , β , γ	39.10Å 70.55Å 87.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.85 – 1.13 43.85 – 1.13	Depositor EDS
% Data completeness (in resolution range)	99.1 (43.85-1.13) 99.1 (43.85-1.13)	Depositor EDS
R_{merge}	0.01	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.63 (at 1.13Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.153 , 0.176 0.154 , 0.176	Depositor DCC
R_{free} test set	4580 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	10.0	Xtriage
Anisotropy	0.369	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5339	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, GAL, FUC, GLA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.41	0/1235	0.68	0/1665
1	C	0.38	0/1247	0.67	0/1681
All	All	0.39	0/2482	0.68	0/3346

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	95	TYR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1171	1160	1156	11	0
1	C	1179	1163	1156	7	0
2	E	47	45	42	1	0
2	F	47	45	42	1	0
3	A	242	0	0	7	3
3	C	240	0	0	3	2
All	All	2926	2413	2396	20	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:28:SER:OG	3:A:201:HOH:O	1.99	0.79
1:A:83:GLU:OE1	3:A:202:HOH:O	2.02	0.76
1:A:80:GLU:OE1	3:A:203:HOH:O	2.11	0.69
1:C:128[A]:GLU:OE2	3:C:201:HOH:O	2.15	0.65
1:A:136:LYS:O	3:A:204:HOH:O	2.17	0.60
2:F:1:NAG:H82	2:F:3:FUC:H4	1.90	0.53
2:E:1:NAG:H82	2:E:3:FUC:H4	1.89	0.53
1:A:80:GLU:OE2	3:A:205:HOH:O	2.18	0.52
1:A:29:ASN:HA	1:A:81:ALA:HB1	1.91	0.51
1:C:78[A]:PRO:HG2	1:C:100:ILE:HD12	1.95	0.49
1:A:113:LYS:NZ	3:A:211:HOH:O	2.45	0.49
1:A:97:HIS:C	1:A:98:ILE:HD12	2.37	0.45
1:A:78:PRO:HA	3:A:412:HOH:O	2.17	0.44
1:C:122:GLN:NE2	3:C:203:HOH:O	2.28	0.44
1:C:97:HIS:C	1:C:98:ILE:HD12	2.38	0.44
1:C:40:HIS:O	1:C:117:SER:HB2	2.20	0.42
1:A:48:HIS:O	1:A:60:CYS:HA	2.21	0.40
1:C:78[A]:PRO:HG2	1:C:100:ILE:CD1	2.50	0.40
1:C:128[B]:GLU:OE1	3:C:202:HOH:O	2.22	0.40
1:A:29:ASN:OD1	1:A:82:LYS:HE3	2.21	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:373:HOH:O	3:A:373:HOH:O[2_565]	1.91	0.29
3:A:390:HOH:O	3:A:390:HOH:O[2_565]	1.94	0.26

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:387:HOH:O	3:C:429:HOH:O[4_654]	1.96	0.24
3:C:360:HOH:O	3:C:360:HOH:O[2_565]	2.06	0.14

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	145/139 (104%)	138 (95%)	6 (4%)	1 (1%)	22	4
1	C	146/139 (105%)	141 (97%)	4 (3%)	1 (1%)	22	4
All	All	291/278 (105%)	279 (96%)	10 (3%)	2 (1%)	22	4

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	125	ASN
1	C	125	ASN

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	134/127 (106%)	134 (100%)	0	100	100
1	C	136/127 (107%)	136 (100%)	0	100	100
All	All	270/254 (106%)	270 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	29	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	E	1	2	15,15,15	1.17	0	21,21,21	1.50	4 (19%)
2	GAL	E	2	2	11,11,12	1.31	1 (9%)	15,15,17	1.00	0
2	FUC	E	3	2	10,10,11	1.63	2 (20%)	14,14,16	0.98	0
2	GLA	E	4	2	11,11,12	1.21	0	15,15,17	0.96	1 (6%)
2	NAG	F	1	2	15,15,15	1.05	2 (13%)	21,21,21	1.60	6 (28%)
2	GAL	F	2	2	11,11,12	0.94	0	15,15,17	0.95	0
2	FUC	F	3	2	10,10,11	1.80	2 (20%)	14,14,16	1.05	0
2	GLA	F	4	2	11,11,12	0.95	0	15,15,17	0.80	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '·' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	E	1	2	-	2/6/26/26	0/1/1/1
2	GAL	E	2	2	-	0/2/19/22	0/1/1/1
2	FUC	E	3	2	-	-	0/1/1/1
2	GLA	E	4	2	-	0/2/19/22	0/1/1/1
2	NAG	F	1	2	-	2/6/26/26	0/1/1/1
2	GAL	F	2	2	-	0/2/19/22	0/1/1/1
2	FUC	F	3	2	-	-	0/1/1/1
2	GLA	F	4	2	-	0/2/19/22	0/1/1/1

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	3	FUC	C2-C3	4.07	1.58	1.52
2	E	3	FUC	C2-C3	3.46	1.57	1.52
2	F	3	FUC	O2-C2	-2.42	1.38	1.43
2	F	1	NAG	C4-C5	2.36	1.58	1.53
2	E	3	FUC	O2-C2	-2.23	1.38	1.43
2	F	1	NAG	C1-C2	2.15	1.55	1.52
2	E	2	GAL	O5-C1	-2.09	1.40	1.43

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	1	NAG	C4-C3-C2	-3.38	105.39	110.34
2	F	1	NAG	C1-C2-C3	-2.85	106.66	110.54
2	F	1	NAG	C1-C2-N2	-2.78	107.51	110.73
2	E	1	NAG	C1-C2-N2	-2.75	107.54	110.73
2	E	1	NAG	C4-C3-C2	-2.66	106.45	110.34
2	F	1	NAG	C1-O5-C5	-2.34	109.25	113.66
2	E	1	NAG	C1-C2-C3	-2.24	107.49	110.54
2	E	4	GLA	C1-O5-C5	2.18	115.15	112.19
2	F	1	NAG	C8-C7-N2	2.14	119.73	116.10
2	F	1	NAG	O7-C7-C8	-2.13	118.09	122.06
2	E	1	NAG	C8-C7-N2	2.01	119.50	116.10

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	E	1	NAG	C8-C7-N2-C2
2	E	1	NAG	O7-C7-N2-C2
2	F	1	NAG	C8-C7-N2-C2

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	F	1	NAG	O7-C7-N2-C2


There are no ring outliers.

4 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	3	FUC	1	0
2	E	3	FUC	1	0
2	F	1	NAG	1	0
2	E	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	138/139 (99%)	3.20	115 (83%) 0 0	8, 11, 21, 34	0
1	C	138/139 (99%)	3.02	112 (81%) 0 0	8, 11, 17, 35	0
All	All	276/278 (99%)	3.11	227 (82%) 0 0	8, 11, 20, 35	0

All (227) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	140	TYR	10.6
1	A	76	THR	7.6
1	A	137	ARG	7.6
1	A	140	TYR	6.9
1	C	139	LEU	6.4
1	A	139	LEU	6.4
1	C	99	PHE	6.3
1	C	138	GLY	6.0
1	A	26	THR	5.8
1	A	138	GLY	5.6
1	C	134	ILE	5.5
1	A	105	ILE	5.4
1	A	90	TYR	5.2
1	C	124	LEU	5.1
1	C	38	LEU	5.1
1	A	129	ILE	5.1
1	A	58	ILE	4.9
1	A	18	TRP	4.8
1	A	21	VAL	4.8
1	C	127	ILE	4.8
1	C	18	TRP	4.7
1	A	77	PHE	4.7
1	C	105	ILE	4.7
1	A	42	GLY	4.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	38	LEU	4.6
1	A	118	ILE	4.6
1	C	4	LEU	4.5
1	A	68	TRP	4.5
1	C	54	ALA	4.5
1	C	60	CYS	4.5
1	A	53	PHE	4.5
1	A	89[A]	ILE	4.5
1	A	78	PRO	4.5
1	C	9	LEU	4.4
1	C	123	ILE	4.4
1	A	75	LYS	4.4
1	A	9	LEU	4.4
1	A	124[A]	LEU	4.3
1	C	129	ILE	4.3
1	C	118	ILE	4.3
1	A	121	LEU	4.1
1	A	37	PHE	4.1
1	A	45	ILE	4.1
1	A	3	ALA	4.1
1	A	6[A]	PHE	4.1
1	A	35	ILE	4.1
1	C	58	ILE	4.1
1	C	42	GLY	4.1
1	A	95	TYR	4.1
1	C	57	ARG	4.1
1	C	90	TYR	4.0
1	C	59	VAL	4.0
1	A	134	ILE	4.0
1	C	121	LEU	4.0
1	C	6[A]	PHE	4.0
1	C	37	PHE	4.0
1	C	14[A]	MET	4.0
1	A	20	VAL	3.9
1	C	43	ASP	3.9
1	C	22	VAL	3.9
1	C	115	LEU	3.9
1	C	15	CYS	3.9
1	A	63	PHE	3.9
1	A	17[A]	GLY	3.9
1	A	59	VAL	3.9
1	A	14[A]	MET	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	35	ILE	3.8
1	A	85	PHE	3.8
1	A	82	LYS	3.8
1	A	51	PRO	3.8
1	A	60	CYS	3.8
1	C	108	TYR	3.8
1	C	20	VAL	3.8
1	A	54	ALA	3.8
1	A	22	VAL	3.7
1	A	132	VAL	3.7
1	C	68	TRP	3.7
1	C	96	PHE	3.7
1	A	133	GLU	3.7
1	C	77	PHE	3.7
1	A	28	SER	3.7
1	C	55	SER	3.7
1	A	100	ILE	3.7
1	A	123	ILE	3.7
1	C	28	SER	3.7
1	A	79	PHE	3.6
1	A	27	SER	3.6
1	C	132	VAL	3.6
1	A	127	ILE	3.6
1	C	100	ILE	3.6
1	A	115	LEU	3.6
1	A	49	PHE	3.5
1	C	47	PHE	3.5
1	A	30	THR	3.5
1	A	119	THR	3.5
1	A	46	ALA	3.5
1	A	108	TYR	3.5
1	C	120	LYS	3.5
1	A	43	ASP	3.4
1	C	21	VAL	3.4
1	C	41	PRO	3.4
1	C	10	TYR	3.4
1	C	53	PHE	3.4
1	A	64	LEU	3.4
1	C	40	HIS	3.4
1	A	15	CYS	3.4
1	C	64	LEU	3.4
1	C	102	GLU	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	98	ILE	3.3
1	A	33	PHE	3.3
1	A	87	VAL	3.3
1	A	40	HIS	3.3
1	A	81	ALA	3.3
1	A	73	VAL	3.3
1	A	47	PHE	3.2
1	C	33	PHE	3.2
1	C	106	LEU	3.2
1	A	41	PRO	3.2
1	C	63	PHE	3.2
1	A	136	LYS	3.2
1	A	4	LEU	3.2
1	C	8	ALA	3.2
1	C	98	ILE	3.2
1	C	85	PHE	3.1
1	C	80[A]	GLU	3.1
1	C	46	ALA	3.1
1	A	5[A]	ARG	3.1
1	C	87	VAL	3.1
1	A	10	TYR	3.0
1	C	89	ILE	3.0
1	C	119	THR	3.0
1	C	79	PHE	3.0
1	A	130	SER	3.0
1	A	29	ASN	3.0
1	A	109	LYS	3.0
1	C	122	GLN	3.0
1	A	16	PRO	3.0
1	A	61	ASN	3.0
1	A	13	GLY	3.0
1	C	49	PHE	3.0
1	A	8	ALA	2.9
1	C	29	ASN	2.9
1	C	61	ASN	2.9
1	A	70	LYS	2.9
1	A	74	ASN	2.9
1	C	84	PRO	2.9
1	A	80	GLU	2.9
1	A	117[A]	SER	2.9
1	C	26	THR	2.9
1	C	32	MET	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	45	ILE	2.8
1	C	131[A]	SER	2.8
1	A	84	PRO	2.8
1	C	13	GLY	2.8
1	C	24	GLY	2.8
1	A	96	PHE	2.8
1	C	95	TYR	2.8
1	A	99	PHE	2.7
1	C	16	PRO	2.7
1	C	51	PRO	2.7
1	A	135	THR	2.7
1	C	30	THR	2.7
1	C	5[A]	ARG	2.7
1	A	106	LEU	2.7
1	C	74	ASN	2.7
1	C	76	THR	2.7
1	C	81	ALA	2.7
1	C	128[A]	GLU	2.7
1	A	19[A]	SER	2.6
1	C	93	GLN	2.6
1	C	94	ASP	2.6
1	C	130	SER	2.6
1	A	24	GLY	2.6
1	C	112	GLN	2.6
1	A	122	GLN	2.5
1	C	3	ALA	2.5
1	A	56	SER	2.5
1	A	32	MET	2.5
1	A	116[A]	SER	2.5
1	C	65	ALA	2.5
1	A	11	PRO	2.5
1	A	62	SER	2.5
1	A	131	SER	2.5
1	C	135	THR	2.4
1	C	17	GLY	2.4
1	C	69	GLY	2.4
1	C	11	PRO	2.4
1	C	78[A]	PRO	2.4
1	A	110	HIS	2.4
1	C	136	LYS	2.4
1	C	73	VAL	2.4
1	C	48	HIS	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	19[A]	SER	2.4
1	C	36	ASN	2.3
1	C	70	LYS	2.3
1	C	109	LYS	2.3
1	C	110	HIS	2.3
1	A	112	GLN	2.3
1	C	133	GLU	2.3
1	C	56	SER	2.2
1	A	120	LYS	2.2
1	A	12	GLU	2.2
1	A	23	LYS	2.2
1	A	57	ARG	2.2
1	A	111	ARG	2.2
1	A	48	HIS	2.2
1	A	67	HIS	2.2
1	C	7	GLU	2.2
1	A	36	ASN	2.2
1	A	25	LYS	2.2
1	A	93	GLN	2.2
1	A	39	SER	2.2
1	A	86	GLN	2.2
1	A	50	ASN	2.1
1	A	7	GLU	2.1
1	C	62	SER	2.1
1	C	116[A]	SER	2.1
1	A	103	ASN	2.1
1	C	50	ASN	2.1
1	C	39	SER	2.1
1	C	117	SER	2.1
1	C	23	LYS	2.1
1	A	126	ASP	2.1
1	C	27[A]	SER	2.1
1	C	82	LYS	2.0
1	C	31	SER	2.0
1	A	107	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains


There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	E	1	15/15	0.78	0.20	12,19,27,27	0
2	FUC	E	3	10/11	0.85	0.16	12,15,18,19	0
2	GLA	E	4	11/12	0.85	0.15	13,17,20,22	0
2	NAG	F	1	15/15	0.87	0.18	10,17,24,24	0
2	FUC	F	3	10/11	0.87	0.19	11,14,17,17	0
2	GLA	F	4	11/12	0.89	0.15	11,15,18,19	0
2	GAL	E	2	11/12	0.93	0.15	10,12,14,14	0
2	GAL	F	2	11/12	0.94	0.16	8,10,11,12	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.