

# wwPDB X-ray Structure Validation Summary Report (i)

#### Nov 20, 2024 - 09:05 am GMT

PDB ID	:	8P8O
Title	:	M. tuberculosis dUTPase - Stl1-159 (StlNT) complex structure
Authors	:	Benedek, A.; Leveles, I.; Toth, Z.S.; Harmat, V.; Vertessy, B.G.
Deposited on	:	2023-06-02
Resolution	:	3.40  Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Motric	Whole archive	Similar resolution		
	$(\# { m Entries})$	(# Entries, resolution range(Å))		
R <sub>free</sub>	164625	1140 (3.46-3.34)		
Clashscore	180529	1172 (3.46-3.34)		
Ramachandran outliers	177936	1172 (3.46-3.34)		
Sidechain outliers	177891	1172 (3.46-3.34)		
RSRZ outliers	164620	1140 (3.46-3.34)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
1	А	174	% •	16%	26%			
		117	23%	10%	20%			
1	В	174	64%	10%	26%			
1	С	174	64%	13%	24%			
1	D	174	59%	15%	26%			
1	E	174	66%	10%	24%			



Mol	Chain	Length	Quality of chain						
1	G	174	63% 12%	25%					
2	F	165	76%	9% • 13%					
2	Н	165	78%	8% • 13%					
2	Ι	165	73%	13% • 13%					
2	J	165	72%	13% • 14%					
2	K	165	% 56% · ·	39%					
2	L	165	73%	13% 13%					



# 2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 11820 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	Δ	190	Total	С	Ν	0	S	0	0	0
1	A	129	941	595	173	172	1	0	0	
1	Р	198	Total	С	Ν	0	S	0	0	0
1	D	128	931	592	166	172	1	0		0
1	1 0	133	Total	С	Ν	0	S	0	0	0
1	U		955	604	171	179	1			
1	Л	129	Total	С	Ν	0	S	0	0	0
1	D		898	571	158	168	1			
1	F	122	Total	С	Ν	0	S	0	0	0
	155	948	598	171	178	1	0	0	0	
1 G	191	Total	С	Ν	0	S	0	0	0	
	131	914	580	159	174	1	U	U	U	

• Molecule 1 is a protein called Deoxyuridine 5'-triphosphate nucleotidohydrolase.

There are 120 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	-19	MET	-	initiating methionine	UNP A0A045IIQ9
А	-18	GLY	-	expression tag	UNP A0A045IIQ9
А	-17	SER	-	expression tag	UNP A0A045IIQ9
А	-16	SER	-	expression tag	UNP A0A045IIQ9
А	-15	HIS	-	expression tag	UNP A0A045IIQ9
А	-14	HIS	-	expression tag	UNP A0A045IIQ9
А	-13	HIS	-	expression tag	UNP A0A045IIQ9
А	-12	HIS	-	expression tag	UNP A0A045IIQ9
А	-11	HIS	-	expression tag	UNP A0A045IIQ9
А	-10	HIS	-	expression tag	UNP A0A045IIQ9
А	-9	SER	-	expression tag	UNP A0A045IIQ9
А	-8	SER	-	expression tag	UNP A0A045IIQ9
А	-7	GLY	-	expression tag	UNP A0A045IIQ9
А	-6	LEU	-	expression tag	UNP A0A045IIQ9
А	-5	VAL	-	expression tag	UNP A0A045IIQ9
А	-4	PRO	-	expression tag	UNP A0A045IIQ9
A	-3	ARG	-	expression tag	UNP A0A045IIQ9



Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	_	expression tag	UNP A0A045IIQ9
A	-1	SER	-	expression tag	UNP A0A045IIQ9
A	0	HIS	_	expression tag	UNP A0A045IIQ9
В	-19	MET	_	initiating methionine	UNP A0A045IIQ9
В	-18	GLY	-	expression tag	UNP A0A045IIQ9
В	-17	SER	-	expression tag	UNP A0A045IIQ9
В	-16	SER	-	expression tag	UNP A0A045IIQ9
В	-15	HIS	-	expression tag	UNP A0A045IIQ9
В	-14	HIS	-	expression tag	UNP A0A045IIQ9
В	-13	HIS	-	expression tag	UNP A0A045IIQ9
В	-12	HIS	-	expression tag	UNP A0A045IIQ9
В	-11	HIS	-	expression tag	UNP A0A045IIQ9
В	-10	HIS	-	expression tag	UNP A0A045IIQ9
В	-9	SER	-	expression tag	UNP A0A045IIQ9
В	-8	SER	-	expression tag	UNP A0A045IIQ9
В	-7	GLY	-	expression tag	UNP A0A045IIQ9
В	-6	LEU	-	expression tag	UNP A0A045IIQ9
В	-5	VAL	-	expression tag	UNP A0A045IIQ9
В	-4	PRO	-	expression tag	UNP A0A045IIQ9
В	-3	ARG	-	expression tag	UNP A0A045IIQ9
В	-2	GLY	-	expression tag	UNP A0A045IIQ9
В	-1	SER	-	expression tag	UNP A0A045IIQ9
В	0	HIS	-	expression tag	UNP A0A045IIQ9
С	-19	MET	-	initiating methionine	UNP A0A045IIQ9
С	-18	GLY	-	expression tag	UNP A0A045IIQ9
С	-17	SER	-	expression tag	UNP A0A045IIQ9
С	-16	SER	-	expression tag	UNP A0A045IIQ9
С	-15	HIS	-	expression tag	UNP A0A045IIQ9
С	-14	HIS	-	expression tag	UNP A0A045IIQ9
С	-13	HIS	-	expression tag	UNP A0A045IIQ9
С	-12	HIS	-	expression tag	UNP A0A045IIQ9
С	-11	HIS	-	expression tag	UNP A0A045IIQ9
С	-10	HIS	-	expression tag	UNP A0A045IIQ9
С	-9	SER	-	expression tag	UNP A0A045IIQ9
С	-8	SER	-	expression tag	UNP A0A045IIQ9
C	-7	GLY	-	expression tag	UNP A0A045IIQ9
C	-6	LEU	-	expression tag	UNP A0A045IIQ9
C	-5	VAL	-	expression tag	UNP A0A045IIQ9
C	-4	PRO	-	expression tag	UNP A0A045IIQ9
C	-3	ARG	-	expression tag	UNP A0A045IIQ9
C	-2	GLY	-	expression tag	UNP A0A045IIQ9
C C	-1	SER	-	expression tag	UNP A0A045IIQ9



Chain	Residue	Modelled	Actual	Comment	Reference
С	0	HIS	_	expression tag	UNP A0A045IIQ9
D	-19	MET	_	initiating methionine	UNP A0A045IIQ9
D	-18	GLY	_	expression tag	UNP A0A045IIQ9
D	-17	SER	_	expression tag	UNP A0A045IIQ9
D	-16	SER	_	expression tag	UNP A0A045IIQ9
D	-15	HIS	-	expression tag	UNP A0A045IIQ9
D	-14	HIS	-	expression tag	UNP A0A045IIQ9
D	-13	HIS	-	expression tag	UNP A0A045IIQ9
D	-12	HIS	-	expression tag	UNP A0A045IIQ9
D	-11	HIS	-	expression tag	UNP A0A045IIQ9
D	-10	HIS	-	expression tag	UNP A0A045IIQ9
D	-9	SER	-	expression tag	UNP A0A045IIQ9
D	-8	SER	-	expression tag	UNP A0A045IIQ9
D	-7	GLY	-	expression tag	UNP A0A045IIQ9
D	-6	LEU	-	expression tag	UNP A0A045IIQ9
D	-5	VAL	-	expression tag	UNP A0A045IIQ9
D	-4	PRO	-	expression tag	UNP A0A045IIQ9
D	-3	ARG	-	expression tag	UNP A0A045IIQ9
D	-2	GLY	-	expression tag	UNP A0A045IIQ9
D	-1	SER	-	expression tag	UNP A0A045IIQ9
D	0	HIS	-	expression tag	UNP A0A045IIQ9
Е	-19	MET	-	initiating methionine	UNP A0A045IIQ9
Е	-18	GLY	-	expression tag	UNP A0A045IIQ9
Е	-17	SER	-	expression tag	UNP A0A045IIQ9
Е	-16	SER	-	expression tag	UNP A0A045IIQ9
Е	-15	HIS	-	expression tag	UNP A0A045IIQ9
Е	-14	HIS	-	expression tag	UNP A0A045IIQ9
Е	-13	HIS	-	expression tag	UNP A0A045IIQ9
E	-12	HIS	-	expression tag	UNP A0A045IIQ9
Е	-11	HIS	-	expression tag	UNP A0A045IIQ9
Е	-10	HIS	-	expression tag	UNP A0A045IIQ9
E	-9	SER	-	expression tag	UNP A0A045IIQ9
Е	-8	SER	-	expression tag	UNP A0A045IIQ9
E	-7	GLY	-	expression tag	UNP A0A045IIQ9
Е	-6	LEU	-	expression tag	UNP A0A045IIQ9
Е	-5	VAL	-	expression tag	UNP A0A045IIQ9
Е	-4	PRO	-	expression tag	UNP A0A045IIQ9
Е	-3	ARG	-	expression tag	UNP A0A045IIQ9
Е	-2	GLY	-	expression tag	UNP A0A045IIQ9
E	-1	SER	-	expression tag	UNP A0A045IIQ9
Е	0	HIS	-	expression tag	UNP A0A045IIQ9
G	-19	MET	-	initiating methionine	UNP A0A045IIQ9



Chain	Residue	Modelled	Actual	Comment	Reference
G	-18	GLY	-	expression tag	UNP A0A045IIQ9
G	-17	SER	-	expression tag	UNP A0A045IIQ9
G	-16	SER	-	expression tag	UNP A0A045IIQ9
G	-15	HIS	-	expression tag	UNP A0A045IIQ9
G	-14	HIS	-	expression tag	UNP A0A045IIQ9
G	-13	HIS	-	expression tag	UNP A0A045IIQ9
G	-12	HIS	-	expression tag	UNP A0A045IIQ9
G	-11	HIS	-	expression tag	UNP A0A045IIQ9
G	-10	HIS	-	expression tag	UNP A0A045IIQ9
G	-9	SER	-	expression tag	UNP A0A045IIQ9
G	-8	SER	-	expression tag	UNP A0A045IIQ9
G	-7	GLY	-	expression tag	UNP A0A045IIQ9
G	-6	LEU	-	expression tag	UNP A0A045IIQ9
G	-5	VAL	-	expression tag	UNP A0A045IIQ9
G	-4	PRO	-	expression tag	UNP A0A045IIQ9
G	-3	ARG	-	expression tag	UNP A0A045IIQ9
G	-2	GLY	-	expression tag	UNP A0A045IIQ9
G	-1	SER	-	expression tag	UNP A0A045IIQ9
G	0	HIS	-	expression tag	UNP A0A045IIQ9

• Molecule 2 is a protein called Orf20.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
9	Ц	1/12	Total	С	Ν	Ο	S	0	0	0
	11	140	1108	710	173	223	2	0	0	0
0	т	1/12	Total	С	Ν	0	S	0	0	0
	1	140	1075	690	169	214	2		0	0
0	т	149	Total	С	Ν	0	S	0	0	0
	1	142	1151	740	186	223	2	0		0
0	Б	1/12	Total	С	Ν	0	S	0	0	0
	T,	140	1114	708	178	226	2	0	0	0
0	V	101	Total	С	Ν	0	S	0	0	0
	Γ	101	707	452	111	142	2	0	0	0
0	т	1/12	Total	С	Ν	0	S	0	0	0
	Г	140	1078	682	174	220	2			U

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual Comment		Reference
H	-5	GLY	-	expression tag	UNP Q9F0J8
Н	-4	SER	-	expression tag	UNP Q9F0J8
Н	-3	PRO	-	expression tag	UNP Q9F0J8



Chain	Residue	Modelled	Actual	Comment	Reference
Н	-2	GLU	-	expression tag	UNP Q9F0J8
Н	-1	PHE	-	expression tag	UNP Q9F0J8
Н	0	SER	-	expression tag	UNP Q9F0J8
Ι	-5	GLY	-	expression tag	UNP Q9F0J8
Ι	-4	SER	-	expression tag	UNP Q9F0J8
Ι	-3	PRO	-	expression tag	UNP Q9F0J8
Ι	-2	GLU	-	expression tag	UNP Q9F0J8
Ι	-1	PHE	-	expression tag	UNP Q9F0J8
Ι	0	SER	-	expression tag	UNP Q9F0J8
J	-5	GLY	-	expression tag	UNP Q9F0J8
J	-4	SER	-	expression tag	UNP Q9F0J8
J	-3	PRO	-	expression tag	UNP Q9F0J8
J	-2	GLU	-	expression tag	UNP Q9F0J8
J	-1	PHE	-	expression tag	UNP Q9F0J8
J	0	SER	-	expression tag	UNP Q9F0J8
F	-5	GLY	-	expression tag	UNP Q9F0J8
F	-4	SER	-	expression tag	UNP Q9F0J8
F	-3	PRO	-	expression tag	UNP Q9F0J8
F	-2	GLU	-	expression tag	UNP Q9F0J8
F	-1	PHE	-	expression tag	UNP Q9F0J8
F	0	SER	-	expression tag	UNP Q9F0J8
K	-5	GLY	-	expression tag	UNP Q9F0J8
K	-4	SER	-	expression tag	UNP Q9F0J8
K	-3	PRO	-	expression tag	UNP Q9F0J8
K	-2	GLU	-	expression tag	UNP Q9F0J8
K	-1	PHE	-	expression tag	UNP Q9F0J8
K	0	SER	-	expression tag	UNP Q9F0J8
L	-5	GLY	-	expression tag	UNP Q9F0J8
L	-4	SER	-	expression tag	UNP Q9F0J8
L	-3	PRO	-	expression tag	UNP Q9F0J8
L	-2	GLU	-	expression tag	UNP Q9F0J8
L	-1	PHE	-	expression tag	UNP Q9F0J8
L	0	SER	-	expression tag	UNP Q9F0J8



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Deoxyuridine 5'-triphosphate nucleotidohydrolase





## • Molecule 1: Deoxyuridine 5'-triphosphate nucleotidohydrolase Chain E: 66% 10% 24% MET GLY SER SER HIS HIS HIS HIS HIS HIS SER VAL PRO ARG GLY SEH SEH HIS MET ALA SER SER ARG GLY ASP ASP CLY CLY SER SER ALA ALA ALA ALA ALA LEU • Molecule 1: Deoxyuridine 5'-triphosphate nucleotidohydrolase Chain G: 63% 25% 12% MET GLY SERR SERR HIS HIS HIS HIS HIS HIS HIS HIS SER HIS SER KIS CUY VAL LEU VAL LEU VAL CEUN ALA GLY GLY ALEU ALEU SERR SERR ARG GLY HIS SER HHIS SER CLY GLY HIS SER HIS SER HIS SER ALA • Molecule 2: Orf20 Chain H: 78% 8% 13% GLY SER PRO GLU PHE SER MET GLV GLV GLV MET ALA ALA CLU LEU THH GLU LYS SEB • Molecule 2: Orf20 Chain I: 73% 13% 13% GLY SER PRO GLU GLU GLU GLY GLY GLY MET ALA ALA ALA ALA CLU ILEU GLU SER SER • Molecule 2: Orf20 Chain J: 72% 13% 14% GLY SER GLU GLU GLU MET MET ALA GLY GLN MET ALA ALA ALA ALA GLU LYS SER

• Molecule 2: Orf20







## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	104.67Å 123.82Å 170.23Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
Bosolution (Å)	46.38 - 3.40	Depositor
Resolution (A)	46.38 - 3.40	EDS
% Data completeness	96.9 (46.38-3.40)	Depositor
(in resolution range)	96.9(46.38-3.40)	EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.39 (at 3.40 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
P. P.	0.244 , $0.286$	Depositor
$n, n_{free}$	0.249 , $0.252$	DCC
$R_{free}$ test set	2051 reflections $(6.58%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	136.3	Xtriage
Anisotropy	0.275	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.32, 108.3	EDS
L-test for $twinning^2$	$ L  > = 0.48, < L^2 > = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	11820	wwPDB-VP
Average B, all atoms $(Å^2)$	145.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.91% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond	angles
IVIOI	Ullaill	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	А	0.26	0/956	0.57	0/1305
1	В	0.26	0/946	0.54	0/1294
1	С	0.25	0/970	0.54	0/1327
1	D	0.25	0/913	0.50	0/1253
1	Ε	0.25	0/962	0.54	0/1315
1	G	0.25	0/929	0.51	0/1277
2	F	0.24	0/1138	0.40	0/1545
2	Н	0.24	0/1132	0.40	0/1541
2	Ι	0.23	0/1099	0.39	0/1500
2	J	0.24	0/1176	0.41	0/1588
2	Κ	0.23	0/722	0.35	0/986
2	L	0.24	0/1102	0.40	0/1502
All	All	0.24	0/12045	0.46	0/16433

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	941	0	965	18	0
1	В	931	0	961	13	0
1	С	955	0	968	17	0
1	D	898	0	892	18	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	Ε	948	0	955	13	0
1	G	914	0	902	17	0
2	F	1114	0	995	13	0
2	Н	1108	0	996	9	0
2	Ι	1075	0	933	13	0
2	J	1151	0	1109	14	0
2	Κ	707	0	539	4	0
2	L	1078	0	913	13	0
All	All	11820	0	11128	137	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 137 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:53:VAL:HB	1:A:84:ALA:HA	1.61	0.83
2:L:30:SER:HA	2:L:40:GLN:HE21	1.49	0.78
1:E:53:VAL:HB	1:E:84:ALA:HA	1.67	0.76
1:D:53:VAL:HB	1:D:84:ALA:HA	1.70	0.73
1:D:94:LEU:HD11	1:D:111:ILE:HD13	1.72	0.71

There are no symmetry-related clashes.

## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	127/174~(73%)	123 (97%)	4 (3%)	0	100	100
1	В	126/174~(72%)	123 (98%)	3 (2%)	0	100	100
1	С	131/174~(75%)	125~(95%)	6 (5%)	0	100	100



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	D	127/174~(73%)	123~(97%)	4 (3%)	0	100	100
1	Ε	129/174~(74%)	125~(97%)	4 (3%)	0	100	100
1	G	129/174~(74%)	122 (95%)	7 (5%)	0	100	100
2	F	141/165~(86%)	135~(96%)	6 (4%)	0	100	100
2	Н	141/165~(86%)	136 (96%)	5 (4%)	0	100	100
2	Ι	141/165~(86%)	134 (95%)	7 (5%)	0	100	100
2	J	140/165~(85%)	135 (96%)	5 (4%)	0	100	100
2	Κ	97/165~(59%)	93~(96%)	4 (4%)	0	100	100
2	L	141/165~(86%)	135 (96%)	6 (4%)	0	100	100
All	All	1570/2034~(77%)	1509 (96%)	61 (4%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	98/134~(73%)	97~(99%)	1 (1%)	73 83
1	В	98/134~(73%)	98 (100%)	0	100 100
1	С	98/134~(73%)	97~(99%)	1 (1%)	73 83
1	D	90/134~(67%)	89~(99%)	1 (1%)	70 81
1	Е	97/134~(72%)	97~(100%)	0	100 100
1	G	92/134~(69%)	91 (99%)	1 (1%)	70 81
2	F	113/149~(76%)	111 (98%)	2(2%)	54 73
2	Н	112/149~(75%)	107 (96%)	5(4%)	23 50
2	Ι	102/149~(68%)	101 (99%)	1 (1%)	73 83
2	J	124/149~(83%)	123 (99%)	1 (1%)	79 87
2	K	55/149~(37%)	53 (96%)	2 (4%)	30 56
2	L	102/149~(68%)	100 (98%)	2 (2%)	50 70



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	1181/1698~(70%)	1164 (99%)	17 (1%)	62 77

5 of 17 residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
2	Κ	152	LEU
2	L	148	TYR
2	Н	102	ASN
2	Н	148	TYR
2	Ι	148	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

#### 5.6 Ligand geometry (i)

There are no ligands in this entry.

## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	#RSRZ>2	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q<0.9
1	А	129/174~(74%)	-0.60	1 (0%) 82 75	77, 119, 155, 205	0
1	В	128/174~(73%)	-0.79	0 100 100	90, 123, 161, 181	0
1	С	133/174~(76%)	-0.75	0 100 100	92, 131, 169, 224	0
1	D	129/174~(74%)	-0.72	0 100 100	110, 148, 185, 247	0
1	Е	133/174~(76%)	-0.72	0 100 100	114, 142, 179, 200	0
1	G	131/174~(75%)	-0.66	0 100 100	111, 147, 196, 387	0
2	F	143/165~(86%)	-0.87	0 100 100	113, 143, 176, 207	0
2	Н	143/165~(86%)	-0.90	0 100 100	106, 144, 181, 193	0
2	Ι	143/165~(86%)	-0.84	0 100 100	125, 159, 188, 202	0
2	J	142/165~(86%)	-0.84	0 100 100	75, 119, 157, 194	0
2	K	101/165~(61%)	-0.72	2 (1%) 64 55	122, 178, 221, 236	0
2	L	143/165~(86%)	-0.73	0 100 100	139, 170, 209, 223	0
All	All	1598/2034 (78%)	-0.76	3 (0%) 92 92	75, 144, 194, 387	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	Κ	17	ILE	3.1
2	Κ	18	ILE	2.5
1	А	111	ILE	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



## 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

## 6.4 Ligands (i)

There are no ligands in this entry.

## 6.5 Other polymers (i)

There are no such residues in this entry.

