



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 3, 2023 – 11:10 PM EDT

PDB ID : 6PAY  
Title : Structure of HsICDH1:Mg(II):ICT:NADPH(50%) complex reveals structural basis for observation of half-sites reactivity  
Authors : Silvaggi, N.R.; Melkonian, T.R.; Roman, J.V.; Moran, G.R.  
Deposited on : 2019-06-12  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	414	 7% 93% 5%
1	B	414	 6% 95%
1	C	414	 12% 92% 6%
1	D	414	 14% 91% 7%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	FMT	D	503	-	-	-	X

## 2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 25350 atoms, of which 12115 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Isocitrate dehydrogenase [NADP] cytoplasmic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	409	Total 6278	C 2044	H 3061	N 544	O 611	S 18	0	0	0
1	B	412	Total 6302	C 2057	H 3070	N 545	O 612	S 18	0	0	0
1	C	408	Total 6125	C 2011	H 2967	N 528	O 602	S 17	0	0	0
1	D	406	Total 6127	C 2005	H 2977	N 531	O 596	S 18	0	0	0

- Molecule 2 is ISOCITRIC ACID (three-letter code: ICT) (formula: C<sub>6</sub>H<sub>8</sub>O<sub>7</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
2	A	1	Total 17	C 6	H 4	O 7	0	0
2	B	1	Total 17	C 6	H 4	O 7	0	0

*Continued on next page...*





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	H	O	0	0
			4	1	1	2		
5	A	1	Total	C	H	O	0	0
			4	1	1	2		
5	A	1	Total	C	H	O	0	0
			4	1	1	2		
5	A	1	Total	C	H	O	0	0
			5	1	2	2		
5	B	1	Total	C	H	O	0	0
			4	1	1	2		
5	B	1	Total	C	H	O	0	0
			4	1	1	2		
5	D	1	Total	C	H	O	0	0
			4	1	1	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	113	Total	O	0	0
			113	113		
6	B	106	Total	O	0	0
			106	106		
6	C	76	Total	O	0	0
			76	76		
6	D	66	Total	O	0	0
			66	66		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Isocitrate dehydrogenase [NADP] cytoplasmic




- Molecule 1: Isocitrate dehydrogenase [NADP] cytoplasmic



- Molecule 1: Isocitrate dehydrogenase [NADP] cytoplasmic



- Molecule 1: Isocitrate dehydrogenase [NADP] cytoplasmic





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	94.60Å 74.60Å 142.90Å 90.00° 95.20° 90.00°	Depositor
Resolution (Å)	47.10 – 2.20 47.11 – 2.20	Depositor EDS
% Data completeness (in resolution range)	85.5 (47.10-2.20) 85.2 (47.11-2.20)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.11 (at 2.20Å)	Xtrriage
Refinement program	PHENIX (dev_3525: ???)	Depositor
R, $R_{free}$	0.224 , 0.242 0.224 , 0.242	Depositor DCC
$R_{free}$ test set	2000 reflections (2.21%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.4	Xtrriage
Anisotropy	0.107	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 53.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	25350	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 50.80 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.1451e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FMT, NDP, MG, ICT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.25	0/3285	0.46	0/4438
1	B	0.26	0/3300	0.46	0/4456
1	C	0.25	0/3226	0.46	0/4368
1	D	0.26	0/3217	0.47	0/4350
All	All	0.26	0/13028	0.46	0/17612

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3217	3061	3144	13	0
1	B	3232	3070	3164	6	0
1	C	3158	2967	3046	14	0
1	D	3150	2977	3053	19	0
2	A	13	4	4	0	0
2	B	13	4	4	0	0
2	C	13	4	5	1	0
2	D	13	4	4	0	0
3	A	1	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	40	16	19	1	0
5	A	12	5	4	1	0
5	B	6	2	2	0	0
5	D	3	1	1	0	0
6	A	113	0	0	0	0
6	B	106	0	0	0	0
6	C	76	0	0	0	0
6	D	66	0	0	2	0
All	All	13235	12115	12450	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:365:GLU:O	6:D:601:HOH:O	1.84	0.94
1:D:81:LYS:NZ	1:D:85:GLU:OE1	2.09	0.85
1:D:309:HIS:HE1	1:D:331:ALA:HB3	1.52	0.74
1:C:67:HIS:O	1:C:343:ARG:NH2	2.22	0.72
1:A:160:ASP:OD1	1:A:162:THR:OG1	2.11	0.69
1:C:6:SER:O	1:C:349:ASN:ND2	2.33	0.60
1:C:247:GLU:OE1	1:C:249:ARG:NH2	2.34	0.60
1:B:12:GLU:OE2	1:B:27:LYS:NZ	2.28	0.56
4:A:503:NDP:H8A	4:A:503:NDP:H51A	1.87	0.56
1:B:247:GLU:OE1	1:B:249:ARG:NH2	2.38	0.55
1:D:20:ARG:NH2	1:D:43:ASP:OD1	2.40	0.54
1:D:342:HIS:HB2	6:D:612:HOH:O	2.07	0.54
1:D:403:GLU:O	1:D:407:ILE:HD13	2.07	0.54
1:C:392:LEU:HB3	1:C:396:GLU:HG3	1.89	0.53
1:C:325:THR:HG23	1:C:394:THR:OG1	2.09	0.53
1:C:25:LEU:HD11	1:C:29:LYS:HD2	1.92	0.51
1:D:81:LYS:HD3	1:D:81:LYS:C	2.31	0.50
1:C:17:GLU:HB2	1:C:311:THR:HB	1.93	0.50
1:B:160:ASP:OD1	1:B:162:THR:OG1	2.27	0.50
1:A:247:GLU:OE1	1:A:249:ARG:NH2	2.44	0.50
1:C:80:GLU:O	1:C:83:VAL:HG22	2.11	0.50
1:C:20:ARG:NH2	1:C:43:ASP:OD1	2.46	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:14:GLN:NE2	1:B:24:GLU:OE2	2.46	0.49
1:D:309:HIS:CE1	1:D:331:ALA:HB3	2.41	0.49
1:A:167:TYR:HB3	1:D:142:THR:HG21	1.95	0.48
1:D:378:ALA:O	1:D:382:GLY:N	2.46	0.47
1:A:375:ASP:HB3	5:A:507:FMT:C	2.45	0.47
1:A:260:LYS:HA	1:D:120:LEU:HD13	1.97	0.47
1:D:397:PHE:CE2	1:D:401:LEU:HD11	2.50	0.47
1:A:397:PHE:CE2	1:A:401:LEU:HD11	2.50	0.46
1:A:316:TYR:O	1:A:320:GLN:HG2	2.15	0.46
1:D:140:ARG:HD3	1:D:183:TYR:OH	2.16	0.45
1:A:126:LYS:CE	1:A:262:GLU:HG3	2.46	0.44
1:A:247:GLU:OE1	1:A:249:ARG:NH1	2.49	0.44
1:D:316:TYR:O	1:D:320:GLN:HG2	2.16	0.44
1:D:22:ILE:HD11	1:D:327:THR:HB	2.00	0.43
1:C:361:GLU:OE1	1:C:408:LYS:NZ	2.52	0.43
1:C:94:SER:OG	2:C:501:ICT:O3	2.27	0.43
1:D:127:PRO:HG2	1:D:205:TRP:HH2	1.83	0.43
1:D:365:GLU:OE2	1:D:408:LYS:NZ	2.47	0.43
1:C:22:ILE:HD11	1:C:327:THR:HB	2.00	0.42
1:A:252:ASP:N	1:A:252:ASP:OD1	2.53	0.41
1:C:22:ILE:HD11	1:C:327:THR:CG2	2.50	0.41
1:A:123:GLY:O	1:A:262:GLU:HA	2.21	0.41
1:C:151:LYS:HE3	1:C:153:GLU:CD	2.42	0.41
1:D:309:HIS:HE1	1:D:331:ALA:CB	2.29	0.41
1:A:236:LYS:NZ	1:A:240:GLU:OE2	2.50	0.41
1:B:236:LYS:NZ	1:B:240:GLU:OE1	2.54	0.40
1:D:107:VAL:HG23	1:D:134:ALA:HB2	2.02	0.40
1:A:293:SER:HB2	1:A:308:ALA:HB2	2.04	0.40
1:B:24:GLU:O	1:B:28:GLU:HG2	2.20	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	407/414 (98%)	391 (96%)	15 (4%)	1 (0%)	47	55
1	B	410/414 (99%)	397 (97%)	12 (3%)	1 (0%)	47	55
1	C	406/414 (98%)	392 (97%)	14 (3%)	0	100	100
1	D	402/414 (97%)	386 (96%)	16 (4%)	0	100	100
All	All	1625/1656 (98%)	1566 (96%)	57 (4%)	2 (0%)	51	60

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	382	GLY
1	B	382	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	338/350 (97%)	332 (98%)	6 (2%)	59	72
1	B	338/350 (97%)	331 (98%)	7 (2%)	53	67
1	C	326/350 (93%)	321 (98%)	5 (2%)	65	78
1	D	326/350 (93%)	323 (99%)	3 (1%)	78	88
All	All	1328/1400 (95%)	1307 (98%)	21 (2%)	62	76

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	140	ARG
1	A	162	THR
1	A	180	MET
1	A	320	GLN
1	A	389	SER
1	A	398	MET
1	B	4	LYS
1	B	122	SER
1	B	155	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	162	THR
1	B	180	MET
1	B	315	HIS
1	B	389	SER
1	C	29	LYS
1	C	237	SER
1	C	309	HIS
1	C	325	THR
1	C	398	MET
1	D	195	SER
1	D	237	SER
1	D	320	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	40	HIS
1	A	194	HIS
1	B	194	HIS
1	B	309	HIS
1	B	315	HIS
1	C	170	HIS
1	C	194	HIS
1	D	40	HIS
1	D	67	HIS
1	D	194	HIS
1	D	309	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

Of 16 ligands modelled in this entry, 4 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	FMT	D	503	-	2,2,2	0.68	0	1,1,1	0.42	0
2	ICT	C	501	3	12,12,12	1.15	0	13,16,16	1.64	5 (38%)
2	ICT	D	501	3	12,12,12	1.11	0	13,16,16	1.62	5 (38%)
5	FMT	A	505	-	2,2,2	0.68	0	1,1,1	0.43	0
5	FMT	B	503	-	2,2,2	0.69	0	1,1,1	0.44	0
5	FMT	A	504	-	2,2,2	0.69	0	1,1,1	0.43	0
5	FMT	A	506	-	2,2,2	0.66	0	1,1,1	0.43	0
2	ICT	A	501	3	12,12,12	1.13	0	13,16,16	1.56	4 (30%)
5	FMT	B	504	-	2,2,2	0.70	0	1,1,1	0.45	0
4	NDP	A	503	-	36,43,52	2.55	6 (16%)	44,67,80	1.71	10 (22%)
5	FMT	A	507	-	2,2,2	0.67	0	1,1,1	0.22	0
2	ICT	B	501	3	12,12,12	1.21	1 (8%)	13,16,16	1.56	4 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ICT	C	501	3	-	6/16/16/16	-
2	ICT	D	501	3	-	2/16/16/16	-
2	ICT	A	501	3	-	2/16/16/16	-
4	NDP	A	503	-	-	6/23/59/77	0/4/4/5
2	ICT	B	501	3	-	2/16/16/16	-

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	503	NDP	P2B-O2B	12.07	1.82	1.59

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	503	NDP	PN-O5D	4.48	1.77	1.59
4	A	503	NDP	O4B-C1B	4.15	1.46	1.41
4	A	503	NDP	O2B-C2B	-2.61	1.34	1.44
4	A	503	NDP	C2A-N1A	2.42	1.38	1.33
2	B	501	ICT	C2-C1	-2.09	1.49	1.52
4	A	503	NDP	C4A-N3A	2.00	1.38	1.35

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	503	NDP	PN-O3-PA	-5.93	112.46	132.83
4	A	503	NDP	O2B-P2B-O1X	-3.38	96.35	109.39
4	A	503	NDP	PA-O5B-C5B	-3.04	103.85	121.68
4	A	503	NDP	O5D-PN-O1N	-2.72	98.42	109.07
4	A	503	NDP	O3X-P2B-O2X	2.63	117.69	107.64
2	C	501	ICT	O2-C1-C2	2.59	120.26	113.27
2	A	501	ICT	O2-C1-C2	2.49	120.01	113.27
2	D	501	ICT	O2-C1-C2	2.43	119.83	113.27
2	B	501	ICT	O6-C6-O5	-2.40	118.65	124.09
2	D	501	ICT	O6-C6-O5	-2.39	118.65	124.09
2	C	501	ICT	O6-C6-C3	2.39	120.61	114.03
2	B	501	ICT	O2-C1-C2	2.36	119.65	113.27
2	A	501	ICT	O2-C1-O1	-2.31	118.85	124.09
2	D	501	ICT	O6-C6-C3	2.29	120.32	114.03
4	A	503	NDP	C5D-C4D-C3D	-2.28	106.64	115.18
2	C	501	ICT	O6-C6-O5	-2.27	118.93	124.09
2	A	501	ICT	O6-C6-C3	2.22	120.15	114.03
2	D	501	ICT	O4-C5-C4	2.20	121.14	114.07
2	A	501	ICT	O6-C6-O5	-2.16	119.19	124.09
2	B	501	ICT	O6-C6-C3	2.16	119.96	114.03
4	A	503	NDP	C2A-N1A-C6A	-2.11	115.15	118.75
2	C	501	ICT	C3-C4-C5	-2.09	109.89	114.04
4	A	503	NDP	O2N-PN-O1N	2.07	122.50	112.24
2	C	501	ICT	O2-C1-O1	-2.04	119.45	124.09
4	A	503	NDP	PN-O5D-C5D	-2.03	109.80	121.68
2	B	501	ICT	O4-C5-C4	2.02	120.56	114.07
2	D	501	ICT	O2-C1-O1	-2.01	119.52	124.09
4	A	503	NDP	O4B-C1B-C2B	-2.00	103.11	106.59

There are no chirality outliers.

All (18) torsion outliers are listed below:




Mol	Chain	Res	Type	Atoms
4	A	503	NDP	C5D-O5D-PN-O2N
4	A	503	NDP	O4B-C4B-C5B-O5B
4	A	503	NDP	C3B-C4B-C5B-O5B
2	C	501	ICT	O2-C1-C2-C3
2	C	501	ICT	O1-C1-C2-C3
2	C	501	ICT	O2-C1-C2-O7
2	A	501	ICT	C4-C3-C6-O6
2	B	501	ICT	C4-C3-C6-O5
2	B	501	ICT	C4-C3-C6-O6
2	C	501	ICT	C4-C3-C6-O5
2	C	501	ICT	C4-C3-C6-O6
4	A	503	NDP	C5D-O5D-PN-O3
4	A	503	NDP	C5D-O5D-PN-O1N
2	C	501	ICT	O1-C1-C2-O7
2	D	501	ICT	C3-C4-C5-O3
2	D	501	ICT	C3-C4-C5-O4
4	A	503	NDP	PA-O3-PN-O2N
2	A	501	ICT	C4-C3-C6-O5

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	501	ICT	1	0
4	A	503	NDP	1	0
5	A	507	FMT	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	409/414 (98%)	0.73	31 (7%) 13 12	19, 37, 57, 71	0
1	B	412/414 (99%)	0.61	25 (6%) 21 20	21, 35, 47, 60	0
1	C	408/414 (98%)	0.91	48 (11%) 4 4	23, 44, 64, 74	0
1	D	406/414 (98%)	0.92	58 (14%) 2 2	20, 40, 69, 79	0
All	All	1635/1656 (98%)	0.79	162 (9%) 7 6	19, 37, 64, 79	0

All (162) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	409	LEU	5.5
1	C	7	GLY	5.5
1	C	410	ALA	5.5
1	C	84	GLU	5.4
1	C	159	SER	4.9
1	D	410	ALA	4.8
1	C	344	ALA	4.4
1	A	387	GLN	4.3
1	C	406	LYS	4.0
1	C	81	LYS	3.9
1	C	34	TYR	3.9
1	D	44	LEU	3.9
1	C	66	LYS	3.9
1	C	86	PHE	3.9
1	D	409	LEU	3.8
1	C	80	GLU	3.7
1	A	163	GLN	3.7
1	C	412	ALA	3.7
1	B	414	LEU	3.6
1	B	387	GLN	3.6
1	D	309	HIS	3.6

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	D	310	GLY	3.5
1	D	314	ARG	3.5
1	D	411	GLN	3.5
1	D	90	GLN	3.5
1	C	407	ILE	3.4
1	D	384	PRO	3.4
1	D	92	TRP	3.3
1	C	25	LEU	3.3
1	B	156	TYR	3.2
1	D	301	LYS	3.2
1	D	401	LEU	3.2
1	D	83	VAL	3.2
1	D	408	LYS	3.2
1	B	165	VAL	3.1
1	A	409	LEU	3.1
1	A	384	PRO	3.1
1	B	412	ALA	3.1
1	D	86	PHE	3.1
1	C	411	GLN	3.0
1	D	397	PHE	3.0
1	C	149	PRO	3.0
1	D	348	ASN	3.0
1	C	88	LEU	3.0
1	D	151	LYS	3.0
1	D	268	ALA	3.0
1	C	233	LYS	3.0
1	D	371	PHE	2.9
1	B	384	PRO	2.9
1	C	175	GLY	2.9
1	D	88	LEU	2.9
1	A	323	GLN	2.9
1	B	163	GLN	2.9
1	D	149	PRO	2.9
1	C	83	VAL	2.9
1	B	173	GLU	2.9
1	D	175	GLY	2.8
1	A	16	ASP	2.8
1	C	354	PHE	2.8
1	D	7	GLY	2.8
1	C	162	THR	2.8
1	A	408	LYS	2.8
1	D	34	TYR	2.8

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	D	362	VAL	2.7
1	B	164	LYS	2.7
1	D	407	ILE	2.7
1	D	406	LYS	2.7
1	A	410	ALA	2.7
1	C	317	ARG	2.7
1	B	385	ASN	2.7
1	A	159	SER	2.6
1	C	369	ALA	2.6
1	B	159	SER	2.6
1	D	37	LEU	2.6
1	D	40	HIS	2.6
1	B	174	GLU	2.6
1	C	172	PHE	2.6
1	B	151	LYS	2.6
1	C	379	CYS	2.6
1	D	350	LYS	2.5
1	D	31	ILE	2.5
1	A	276	VAL	2.5
1	D	35	VAL	2.5
1	B	389	SER	2.5
1	C	37	LEU	2.5
1	D	346	LEU	2.5
1	D	150	GLY	2.5
1	C	269	CYS	2.4
1	A	77	THR	2.4
1	A	350	LYS	2.4
1	D	317	ARG	2.4
1	D	53	ASN	2.4
1	D	147	PRO	2.4
1	A	182	MET	2.4
1	C	47	GLU	2.4
1	D	276	VAL	2.4
1	C	163	GLN	2.4
1	B	130	ILE	2.4
1	A	73	CYS	2.3
1	C	404	ASN	2.3
1	A	242	GLN	2.3
1	A	156	TYR	2.3
1	D	250	LEU	2.3
1	C	161	GLY	2.3
1	C	276	VAL	2.3

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	A	310	GLY	2.3
1	C	146	VAL	2.3
1	D	52	THR	2.3
1	D	354	PHE	2.3
1	B	255	VAL	2.3
1	C	145	VAL	2.3
1	C	171	ASN	2.3
1	C	50	ASP	2.2
1	B	276	VAL	2.2
1	C	309	HIS	2.2
1	A	391	TYR	2.2
1	A	239	PHE	2.2
1	D	67	HIS	2.2
1	D	313	THR	2.2
1	A	126	LYS	2.2
1	C	33	PRO	2.2
1	C	173	GLU	2.2
1	D	174	GLU	2.2
1	A	307	ALA	2.2
1	D	383	LEU	2.2
1	D	58	LYS	2.2
1	C	214	THR	2.2
1	A	160	ASP	2.2
1	C	9	SER	2.2
1	D	267	TRP	2.2
1	B	160	ASP	2.2
1	D	148	GLY	2.2
1	B	301	LYS	2.2
1	B	388	ARG	2.2
1	B	167	TYR	2.1
1	B	272	TYR	2.1
1	D	391	TYR	2.1
1	D	251	ILE	2.1
1	C	151	LYS	2.1
1	D	66	LYS	2.1
1	A	411	GLN	2.1
1	B	155	THR	2.1
1	A	23	TRP	2.1
1	C	371	PHE	2.1
1	D	369	ALA	2.1
1	D	412	ALA	2.1
1	A	389	SER	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	253	ASP	2.1
1	D	272	TYR	2.1
1	A	289	GLY	2.1
1	D	51	ALA	2.1
1	A	130	ILE	2.0
1	A	383	LEU	2.0
1	C	168	LEU	2.0
1	B	268	ALA	2.0
1	D	81	LYS	2.0
1	C	31	ILE	2.0
1	C	278	SER	2.0
1	D	316	TYR	2.0
1	A	74	ALA	2.0
1	A	75	THR	2.0
1	A	255	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q < 0.9
5	FMT	D	503	3/3	0.73	0.51	17,17,17,21	0
5	FMT	A	506	3/3	0.86	0.29	17,17,17,21	0
5	FMT	A	507	3/3	0.86	0.44	19,19,23,23	0
3	MG	A	502	1/1	0.86	0.14	6,6,6,6	0
3	MG	D	502	1/1	0.87	0.11	29,29,29,29	0
5	FMT	B	503	3/3	0.87	0.33	17,17,17,21	0
3	MG	B	502	1/1	0.87	0.12	8,8,8,8	0
5	FMT	B	504	3/3	0.88	0.29	17,17,17,21	0

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	FMT	A	505	3/3	0.91	0.33	17,17,17,21	0
2	ICT	D	501	13/13	0.91	0.13	17,37,57,58	0
3	MG	C	502	1/1	0.92	0.08	19,19,19,19	0
2	ICT	C	501	13/13	0.94	0.13	21,31,47,47	0
5	FMT	A	504	3/3	0.94	0.38	17,17,17,21	0
2	ICT	B	501	13/13	0.96	0.14	4,12,28,31	0
4	NDP	A	503	40/48	0.96	0.12	11,11,13,13	0
2	ICT	A	501	13/13	0.96	0.14	2,10,18,27	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.