



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 11, 2022 – 10:05 am BST

PDB ID : 7PC8  
Title : The PDZ domain of SNTG1 complexed with the phosphomimetic mutant PDZ-binding motif of RSK1  
Authors : Cousido-Siah, A.; Trave, G.; Gogl, G.  
Deposited on : 2021-08-03  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.27  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0267  
CCP4 : 7.1.010 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.27


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	414	
1	B	414	
2	C	10	
2	D	10	

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 6914 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gamma-1-syntrophin,Annexin A2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	413	Total 3281	C 2056	N 578	O 634	S 13	0	1	0
1	B	409	Total 3253	C 2040	N 572	O 629	S 12	0	1	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	49	GLY	-	expression tag	UNP Q9NSN8
A	50	SER	-	expression tag	UNP Q9NSN8
A	51	HIS	-	expression tag	UNP Q9NSN8
A	52	MET	-	expression tag	UNP Q9NSN8
A	53	GLY	-	expression tag	UNP Q9NSN8
A	144	GLY	-	linker	UNP Q9NSN8
A	189	GLU	ALA	conflict	UNP P07355
B	49	GLY	-	expression tag	UNP Q9NSN8
B	50	SER	-	expression tag	UNP Q9NSN8
B	51	HIS	-	expression tag	UNP Q9NSN8
B	52	MET	-	expression tag	UNP Q9NSN8
B	53	GLY	-	expression tag	UNP Q9NSN8
B	144	GLY	-	linker	UNP Q9NSN8
B	189	GLU	ALA	conflict	UNP P07355

- Molecule 2 is a protein called Ribosomal protein S6 kinase alpha-1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	8	Total 67	C 42	N 12	O 13	0	0	0
2	D	6	Total 47	C 30	N 6	O 11	0	0	0


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	186	GLU	SER	conflict	UNP Q15418
D	186	GLU	SER	conflict	UNP Q15418

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	5	Total Ca 5 5	0	0
3	B	4	Total Ca 4 4	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	B	1	Total C O 6 3 3	0	0
4	B	1	Total C O 6 3 3	0	0
4	B	1	Total C O 6 3 3	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	137	Total 137	O 137	0	0
5	B	83	Total 83	O 83	0	0
5	C	1	Total 1	O 1	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Gamma-1-syntrophin,Annexin A2

Chain A: 




- Molecule 1: Gamma-1-syntrophin,Annexin A2

Chain B: 




- Molecule 2: Ribosomal protein S6 kinase alpha-1

Chain C: 



- Molecule 2: Ribosomal protein S6 kinase alpha-1

Chain D: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	226.58Å 59.82Å 125.21Å 90.00° 117.84° 90.00°	Depositor
Resolution (Å)	48.33 – 2.50 48.33 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.7 (48.33-2.50) 99.7 (48.33-2.50)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.30 (at 2.51Å)	Xtriage
Refinement program	PHENIX 1.14_3260	Depositor
R, $R_{free}$	0.202 , 0.232 0.203 , 0.232	Depositor DCC
$R_{free}$ test set	2584 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	48.7	Xtriage
Anisotropy	0.215	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	6914	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.47% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.27	0/3326	0.42	0/4465
1	B	0.26	0/3297	0.42	0/4427
2	C	0.20	0/67	0.42	0/88
2	D	0.21	0/47	0.52	0/63
All	All	0.26	0/6737	0.42	0/9043

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3281	0	3341	11	0
1	B	3253	0	3317	20	0
2	C	67	0	74	2	0
2	D	47	0	48	0	0
3	A	5	0	0	0	0
3	B	4	0	0	0	0
4	A	18	0	24	0	0
4	B	18	0	24	0	0
5	A	137	0	0	1	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	83	0	0	3	0
5	C	1	0	0	1	0
All	All	6914	0	6828	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:182:ARG:NH2	5:C:200:HOH:O	2.24	0.66
1:B:201:ARG:HG3	1:B:202:THR:HG23	1.87	0.56
1:A:72:ILE:HD11	1:A:122:VAL:HG12	1.86	0.56
1:B:377:ASN:ND2	5:B:606:HOH:O	2.38	0.56
1:A:151:LYS:NZ	5:A:601:HOH:O	2.33	0.55
1:B:324:ALA:HB1	1:B:332:ASP:HB3	1.90	0.53
1:A:60:ILE:HG22	1:A:95:LEU:HD11	1.90	0.53
1:B:92:ARG:HH21	1:B:96:SER:HA	1.76	0.51
1:B:104:ALA:HB2	1:B:139:LEU:HD23	1.93	0.51
1:B:367:ILE:HD13	1:B:379:PHE:HB3	1.93	0.51
1:B:202:THR:HB	1:B:204:LYS:HE3	1.93	0.50
1:A:71:SER:HB2	1:A:86:LYS:HB3	1.93	0.50
1:A:122:VAL:HB	2:C:189:LEU:HD21	1.96	0.48
1:B:327:LYS:NZ	5:B:613:HOH:O	2.47	0.47
1:A:324:ALA:HB1	1:A:332:ASP:HB3	1.97	0.47
1:A:82:VAL:HG21	1:A:121:VAL:HG21	1.96	0.47
1:B:210:LEU:HD12	1:B:226:LEU:HD11	1.98	0.46
1:A:217:HIS:O	1:A:221:VAL:HG23	2.17	0.44
1:A:407:ARG:HE	1:A:409:LYS:HB3	1.83	0.44
1:B:130:GLU:OE1	5:B:601:HOH:O	2.21	0.44
1:B:108:ILE:HD13	1:B:125:LEU:HG	2.00	0.43
1:B:81:PRO:O	1:B:83:VAL:HG23	2.19	0.43
1:B:191:ARG:NH2	1:B:225:LEU:HD22	2.33	0.43
1:B:74:GLY:HA3	1:B:82:VAL:HA	2.01	0.42
1:A:76:ALA:H	1:A:117:ARG:HA	1.84	0.42
1:B:347:HIS:O	1:B:351:VAL:HG23	2.20	0.42
1:B:260:ASN:ND2	1:B:302:ARG:O	2.50	0.41
1:B:371:VAL:O	1:B:372:LYS:HD3	2.20	0.41
1:B:104:ALA:O	1:B:136:VAL:HA	2.20	0.41
1:B:440:TYR:O	1:B:444:GLN:HG2	2.21	0.40
1:A:188:ASN:O	1:A:192:GLN:HG2	2.21	0.40

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:70:LEU:HD23	1:B:70:LEU:HA	1.87	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	412/414 (100%)	406 (98%)	5 (1%)	1 (0%)	47	68
1	B	408/414 (99%)	396 (97%)	10 (2%)	2 (0%)	29	48
2	C	6/10 (60%)	6 (100%)	0	0	100	100
2	D	4/10 (40%)	4 (100%)	0	0	100	100
All	All	830/848 (98%)	812 (98%)	15 (2%)	3 (0%)	34	54

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	174	VAL
1	A	174	VAL
1	B	325	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	358/357 (100%)	347 (97%)	11 (3%)	40	67
1	B	355/357 (99%)	341 (96%)	14 (4%)	32	57
2	C	8/10 (80%)	7 (88%)	1 (12%)	4	8
2	D	6/10 (60%)	6 (100%)	0	100	100
All	All	727/734 (99%)	701 (96%)	26 (4%)	35	61

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	62	ARG
1	A	64	THR
1	A	83	VAL
1	A	90	GLU
1	A	92	ARG
1	A	117	ARG
1	A	119	GLU
1	A	264	GLN
1	A	285	ASP
1	A	407	ARG
1	A	445	ASP
1	B	64	THR
1	B	96	SER
1	B	114	ARG
1	B	119	GLU
1	B	138	PHE
1	B	139	LEU
1	B	141	ARG
1	B	145	SER
1	B	159	GLU
1	B	160	ARG
1	B	172	LYS
1	B	264	GLN
1	B	285	ASP
1	B	445	ASP
2	C	189	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 9 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	GOL	A	506	-	5,5,5	0.90	0	5,5,5	0.99	0
4	GOL	A	507	-	5,5,5	0.90	0	5,5,5	0.96	0
4	GOL	B	505	-	5,5,5	0.86	0	5,5,5	1.00	0
4	GOL	B	506	-	5,5,5	0.85	0	5,5,5	0.99	0
4	GOL	A	508	-	5,5,5	0.93	0	5,5,5	1.02	0
4	GOL	B	507	-	5,5,5	0.89	0	5,5,5	0.99	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	506	-	-	0/4/4/4	-
4	GOL	A	507	-	-	2/4/4/4	-
4	GOL	B	505	-	-	1/4/4/4	-
4	GOL	B	506	-	-	0/4/4/4	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	508	-	-	2/4/4/4	-
4	GOL	B	507	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	507	GOL	C1-C2-C3-O3
4	A	508	GOL	C1-C2-C3-O3
4	A	508	GOL	O2-C2-C3-O3
4	B	507	GOL	O1-C1-C2-C3
4	A	507	GOL	O2-C2-C3-O3
4	B	507	GOL	O1-C1-C2-O2
4	B	505	GOL	C1-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.