



wwPDB EM Validation Summary Report ⓘ

Jul 7, 2024 – 07:23 pm BST

PDB ID : 7PDD
EMDB ID : EMD-13334
Title : Focus refinement of soluble domain of Adenylyl cyclase 9 in complex with DARPin C4 and MANT-GTP
Authors : Qi, C.; Korkhov, V.M.
Deposited on : 2021-08-05
Resolution : 4.20 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev92
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY



The reported resolution of this entry is 4.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1354	
2	B	147	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4516 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Adenylate cyclase 9.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	449	3550	2240	601	677	32	0	0


- Molecule 2 is a protein called DARPin C4.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	128	966	607	171	187	1	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Adenylate cyclase 9



VAL	THR	ASN	ASP	PI038	G1129	E1211	THR	GLU
LEU	LYS	PHE	GLY	Y1039	S1130	E1212	ASP	VAL
SER	ARG	GLN	ILE	H1040	H1131	S1213	GLU	GLY
THR	LEU	LEU	GLU	V1041	F1132	Y1214	ALA	MET
VAL	LEU	SER	ALA	A1042	Q1133	R1215	SER	GLU
PHE	GLU	TRP	SER	E1043	F1140	V1216	LEU	ALA
LEU	TRP	MET	LEU	Q1044	F1140	L1217	VAL	ASN
ILE	ILE	TRP	ILE	L1045	A1143	M1220	PRO	GLU
ALA	GLY	ALA	GLY	K1046	K1144	G1221	SER	LEU
SER	SER	SER	GLN	V1047	E1145	Y1222	VAL	THR
LEU	LEU	TRP	GLU	S1048	M1146	F1223	GLN	LYS
ILE	LEU	PRO	VAL	Q1049	M1147	F1224	ASN	LEU
THR	PRO	ALA	ILE	T1050	R1148	D1225	PRO	ASN
PHE	ARG	THR	VAL	Y1051	V1149	Y1226	ASP	VAL
ARG	PHE	HIS	VAL	S1052	V1150	R1227	GLN	GLU
TYR	ILE	VAL	PHE	I1061	N1154	G1228	PRO	ARG
GLY	ILE	GLY	PHE	F1062	N1155	T1229	PRO	ALA
ALA	ALA	GLY	LEU	A1063	N1156	V1230	GLY	ALA
ALA	ILE	PRO	LEU	M1067	W1159	R1231	SER	GLU
ALA	LEU	LEU	LEU	F1071	F1160	V1232	GLU	ASN
ALA	VAL	LEU	VAL	E1077	N1161	G1236	ASN	ALA
ALA	LEU	LEU	TRP	G1078	F1162	Q1237	ALA	ASN
ALA	PRO	LEU	PHE	G1079	K1163	M1238	GLN	ALA
ALA	ALA	VAL	LEU	K1080	L1164	K1239	THR	ARG
ALA	TYR	VAL	GLU	E1081	R1165	T1240	ARG	ASP
ALA	SER	CYS	PHE	C1082	V1166	Y1241	ALA	ALA
ALA	ASP	PRO	VAL	Y1083	G1167	Y1243	HIS	PRO
PHE	VAL	LEU	VAL	R1084	T1174	P1244	SER	SER
GLY	THR	SER	TYR	V1085	A1175	S1249	ALA	ALA
ALA	GLU	THR	TYR	L1086	I1178	G1250	ARG	ARG
LEU	LEU	VAL	LEU	M1087	G1179	LEU	PRO	PRO
LEU	LEU	ILE	HIS	E1088	T1180	VAL	TRP	TRP
LEU	LEU	PHE	TYR	L1089	T1181	GLN	LYS	LYS
LEU	LEU	ASN	GLY	I1090	K1182	HIS	PRO	GLU
LEU	ASN	ILE	ASP	L1096	L1183	GLN	VAL	VAL
LEU	HIS	HIS	ASP	K1099	L1184	LEU	ARG	GLU
LEU	SER	ALA	VAL	P1100	Y1185	SER	ALA	ALA
LEU	THR	GLN	ALA	D1101	D1186	ILE	GLU	GLU
VAL	MET	ASN	ASP	Y1102	I1187	PRO	ARG	GLU
SER	PHE	PHE	LEU	S1103	W1188	ASP	ARG	ARG
VAL	THR	SER	HIS	S1104	G1189	ILE	CYS	CYS
ARG	SER	THR	ARG	K1107	D1190	ARG	ARG	ARG
MET	ALA	ARG	T1020	I1108	V1192	VAL	PHE	PHE
VAL	VAL	ARG	S1024	T1114	N1193	GLN	GLY	GLY
LEU	LEU	LYS	D1027	Y1115	S1196	VAL	ALA	ALA
THR	THR	CYS	Y1116	M1116	D1199	ASP	ILE	ILE
GLU	VAL	ALA	W1031	M1122	T1200	GLY	GLU	GLU
ASP	VAL	SER	L1032	A1123	E1204	ILE	LYS	SER
VAL	GLN	LEU	L1033	R1206	G1205	GLY	SER	ASP
TYR	PRO	PRO	R1034	I1207	R1206	ARG	CYS	ASP
THR	HIS	HIS	N1035	D1128	I1207	PRO	GLU	GLU

• Molecule 2: DARPin C4



MET	H9	H48	D81
ARG	H10	T49	T82
GLY	G11	P90	P83
SER	G12	L60	L86
HIS	S12	E61	A87
HIS	L14	I62	H88
HIS	G15	V63	R89
H9	K16	E64	R90
H10	K17	V65	G91
G11	E20	L66	L93
G12	R23	L67	H92
S12	A24	K68	L94
L14	G25	A71	E94
G15	Q26	A75	I96
K16	D27	D77	V97
K17	V30	W78	V98
E20	R31	L79	L99
R23	I32	G80	L100
A24	N36		A104
G25	G37		D105
Q26	A38		V106
D27	H48		K111
V30	T49		L119
R31	P90		D122
I32	L51		N123
N36	H52		E126
G37	L53		D127
A38	H59		I128
H48	L60		A129
T49	E61		L132
P90	I62		Q133
L51	V63		A136
H52	E64		LYS
L53	V65		LEU
H59	L66		LEU
L60	L67		ASN
E61	K68		ASN
I62	A71		ASP
V63	A75		TYR
E64	A76		LYS
V65	D77		ASP
L66	W78		ASP
L67	L79		ASP
K68	G80		LYS

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	210729	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	40	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.063	Depositor
Minimum map value	-0.034	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.001	Depositor
Recommended contour level	0.01	Depositor
Map size (\AA)	243.0, 243.0, 243.0	wwPDB
Map dimensions	300, 300, 300	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	0.81, 0.81, 0.81	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/3611	0.48	0/4852
2	B	0.25	0/984	0.40	0/1337
All	All	0.28	0/4595	0.46	0/6189

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3550	0	3486	142	0
2	B	966	0	938	43	0
All	All	4516	0	4424	183	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 183 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1154:ASN:HD21	1:A:1162:PHE:H	1.21	0.88
1:A:392:GLU:N	1:A:493:GLY:O	2.23	0.72

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1206:ARG:HB3	1:A:1242:LEU:HD22	1.72	0.71
1:A:1042:ALA:O	1:A:1046:LYS:HB2	1.90	0.71
1:A:389:GLN:NE2	1:A:494:THR:OG1	2.23	0.71

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	441/1354 (33%)	410 (93%)	31 (7%)	0	100	100
2	B	126/147 (86%)	117 (93%)	9 (7%)	0	100	100
All	All	567/1501 (38%)	527 (93%)	40 (7%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	384/1176 (33%)	383 (100%)	1 (0%)	92	95
2	B	96/114 (84%)	96 (100%)	0	100	100
All	All	480/1290 (37%)	479 (100%)	1 (0%)	93	96

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	348	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1154	ASN
2	B	26	GLN
2	B	48	HIS
1	A	1131	HIS
1	A	420	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-13334. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)

6.2.1 Primary map




The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [\(i\)](#)


6.3.1 Primary map



X Index: 134



Y Index: 160




Z Index: 61


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [\(i\)](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.01. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 51 nm³; this corresponds to an approximate mass of 46 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)




*Reported resolution corresponds to spatial frequency of 0.238\AA^{-1}

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.238 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	4.20	-	-
Author-provided FSC curve	4.07	7.17	4.13
Unmasked-calculated*	-	-	-

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-13334 and PDB model 7PDD. Per-residue inclusion information can be found in section 3 on page 4.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.01 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [\(i\)](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.01).







9.4 Atom inclusion [i](#)



At the recommended contour level, 94% of all backbone atoms, 80% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.01) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.7960	 0.3080
A	 0.8090	 0.3210
B	 0.7480	 0.2600

