



## wwPDB EM Validation Summary Report ⓘ

Sep 18, 2021 – 08:02 AM BST

PDB ID : 7PE8  
EMDB ID : EMD-13348  
Title : cryo-EM structure of DEPTOR bound to human mTOR complex 2, focussed on one protomer  
Authors : Waelchli, M.; Maier, T.  
Deposited on : 2021-08-09  
Resolution : 3.20 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : **FAILED**  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
MolProbity : 4.02b-467  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.1

# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:  
*ELECTRON MICROSCOPY*

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	2571	76% (green), 9% (yellow), 15% (grey)
2	C	326	75% (green), 22% (yellow), . (grey)
3	E	1708	59% (green), 7% (yellow), 35% (grey)
4	G	522	16% (green), . (yellow), 81% (grey)
5	I	409	22% (green), . (yellow), 74% (grey)

## 2 Entry composition i

There are 8 unique types of molecules in this entry. The entry contains 29291 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein kinase mTOR.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	2194	16419	10414	2922	2984	99	0	0

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	246M	GLY	-	insertion	UNP P42345
A	246N	SER	-	insertion	UNP P42345
A	246O	THR	-	insertion	UNP P42345
A	246P	SER	-	insertion	UNP P42345
A	246Q	GLY	-	insertion	UNP P42345
A	246R	SER	-	insertion	UNP P42345
A	246S	GLY	-	insertion	UNP P42345
A	246T	ASP	-	insertion	UNP P42345
A	246U	TYR	-	insertion	UNP P42345
A	246V	LYS	-	insertion	UNP P42345
A	246W	ASP	-	insertion	UNP P42345
A	246X	ASP	-	insertion	UNP P42345
A	246Y	ASP	-	insertion	UNP P42345
A	246Z	ASP	-	insertion	UNP P42345
A	247A	LYS	-	insertion	UNP P42345
A	247B	GLY	-	insertion	UNP P42345
A	247C	SER	-	insertion	UNP P42345
A	247D	THR	-	insertion	UNP P42345
A	247E	SER	-	insertion	UNP P42345
A	247F	GLY	-	insertion	UNP P42345
A	247G	SER	-	insertion	UNP P42345
A	247H	GLY	-	insertion	UNP P42345

- Molecule 2 is a protein called Target of rapamycin complex subunit LST8.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	C	319	Total	C	N	O	S	0	0
			2465	1533	437	477	18		

- Molecule 3 is a protein called Rapamycin-insensitive companion of mTOR.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	E	1115	Total	C	N	O	S	0	0
			8917	5680	1582	1608	47		

- Molecule 4 is a protein called Target of rapamycin complex 2 subunit MAPKAP1.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	G	98	Total	C	N	O	S	0	0
			655	399	125	127	4		


- Molecule 5 is a protein called DEP domain-containing mTOR-interacting protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	I	106	Total	C	N	O	S	0	0
			795	504	138	146	7		

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
I	204	SER	ASN	variant	UNP Q8TB45
I	389	ASN	SER	variant	UNP Q8TB45

- Molecule 6 is INOSITOL HEXAKISPHOSPHATE (three-letter code: IHP) (formula: C<sub>6</sub>H<sub>18</sub>O<sub>24</sub>P<sub>6</sub>).




Mol	Chain	Residues	Atoms				AltConf
			Total	C	O	P	
6	A	1	36	6	24	6	0

- Molecule 7 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		AltConf
			Total	Zn	
7	E	1	1	1	0

- Molecule 8 is ACETYL GROUP (three-letter code: ACE) (formula: C<sub>2</sub>H<sub>4</sub>O).





Mol	Chain	Residues	Atoms			AltConf
			Total	C	O	
8	G	1	3	2	1	0






• Molecule 2: Target of rapamycin complex subunit LST8



• Molecule 3: Rapamycin-insensitive companion of mTOR









## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	750254	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	50	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, IHP, ACE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.23	0/16719	0.43	0/22710
2	C	0.23	0/2523	0.48	0/3438
3	E	0.23	0/9078	0.44	0/12281
4	G	0.22	0/660	0.45	0/900
5	I	0.24	0/811	0.49	0/1104
All	All	0.23	0/29791	0.44	0/40433

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	16419	0	15461	131	0
2	C	2465	0	2351	43	0
3	E	8917	0	9064	70	0
4	G	655	0	519	10	0
5	I	795	0	812	9	0
6	A	36	0	6	0	0
7	E	1	0	0	0	0
8	G	3	0	3	0	0
All	All	29291	0	28216	249	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 249 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2192:LEU:HD22	1:A:2235:GLY:HA3	1.75	0.68
2:C:131:ILE:HA	2:C:147:ASP:HA	1.74	0.68
2:C:146:GLY:HA3	2:C:173:ILE:HD11	1.76	0.67
1:A:737:ILE:HG23	1:A:753:SER:HB2	1.75	0.67
2:C:178:ILE:HG12	2:C:185:MET:HG2	1.76	0.67

There are no symmetry-related clashes.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	2152/2571 (84%)	2100 (98%)	52 (2%)	0	100	100
2	C	317/326 (97%)	298 (94%)	19 (6%)	0	100	100
3	E	1101/1708 (64%)	1064 (97%)	37 (3%)	0	100	100
4	G	94/522 (18%)	90 (96%)	4 (4%)	0	100	100
5	I	104/409 (25%)	95 (91%)	9 (9%)	0	100	100
All	All	3768/5536 (68%)	3647 (97%)	121 (3%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1571/2236 (70%)	1546 (98%)	25 (2%)	62	84
2	C	269/276 (98%)	265 (98%)	4 (2%)	65	85
3	E	985/1539 (64%)	978 (99%)	7 (1%)	84	94
4	G	50/471 (11%)	49 (98%)	1 (2%)	55	80
5	I	89/364 (24%)	86 (97%)	3 (3%)	37	70
All	All	2964/4886 (61%)	2924 (99%)	40 (1%)	70	87

5 of 40 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	325	LEU
3	E	851	LEU
3	E	43	GLN
3	E	524	THR
5	I	315	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1726	GLN
2	C	28	GLN
3	E	607	GLN
3	E	763	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	IHP	A	2601	-	36,36,36	0.79	2 (5%)	54,60,60	0.30	0
8	ACE	G	601	4	1,2,2	0.76	0	1,1,1	0.18	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	IHP	A	2601	-	-	2/30/54/54	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	2601	IHP	P3-O13	3.03	1.65	1.59
6	A	2601	IHP	P1-O11	2.05	1.63	1.59

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	A	2601	IHP	C2-O12-P2-O22
6	A	2601	IHP	C2-O12-P2-O42

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight  $> 250$  and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.