



Full wwPDB X-ray Structure Validation Report i

Jun 17, 2024 – 09:31 pm BST

PDB ID : 8PEQ
Title : Complex of diubiquitin-derived artificial binding protein (Affilin) variant Af2 with its target oncofetal fibronectin (fragment 7B8)
Authors : Parthier, C.; Katschmann, A.; Fiedler, E.; Haupts, U.; Reimann, A.
Deposited on : 2023-06-14
Resolution : 2.32 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.32 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5974 (2.34-2.30)
Clashscore	141614	6604 (2.34-2.30)
Ramachandran outliers	138981	6523 (2.34-2.30)
Sidechain outliers	138945	6523 (2.34-2.30)
RSRZ outliers	127900	5855 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain	
2	M	153	1%	85% 13% •

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 10486 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fibronectin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	274	Total	C 2093	N 1316	O 335	S 441	1	0	0
1	B	275	Total	C 2101	N 1321	O 336	S 442	2	0	0
1	C	275	Total	C 2101	N 1321	O 336	S 442	2	0	0


There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1173	MET	-	initiating methionine	UNP P02751
A	1232	SER	CYS	engineered mutation	UNP P02751
B	1173	MET	-	initiating methionine	UNP P02751
B	1232	SER	CYS	engineered mutation	UNP P02751
C	1173	MET	-	initiating methionine	UNP P02751
C	1232	SER	CYS	engineered mutation	UNP P02751

- Molecule 2 is a protein called Afflin variant Af2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	J	153	Total	C 1226	N 771	O 216	S 237	2	0	0
2	L	151	Total	C 1215	N 765	O 214	S 234	2	0	0
2	M	150	Total	C 1204	N 759	O 210	S 233	2	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	L	1	Total O S 5 4 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	124	Total O 124 124	0	0
4	B	129	Total O 129 129	0	0
4	C	125	Total O 125 125	0	0
4	J	35	Total O 35 35	0	0
4	L	75	Total O 75 75	0	0
4	M	48	Total O 48 48	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Fibronectin




- Molecule 1: Fibronectin




- Molecule 1: Fibronectin



- Molecule 2: Afflin variant Af2





- Molecule 2: Affilin variant Af2

Chain L: 10% ..



- Molecule 2: Affilin variant Af2

Chain M: 13% •



4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	168.65 Å 105.86 Å 78.99 Å 90.00° 93.53° 90.00°	Depositor
Resolution (Å)	44.80 – 2.32 44.80 – 2.32	Depositor EDS
% Data completeness (in resolution range)	99.1 (44.80-2.32) 99.1 (44.80-2.32)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	2.58 (at 2.32 Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R , R_{free}	0.198 , 0.251 0.196 , 0.248	Depositor DCC
R_{free} test set	2983 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	36.1	Xtriage
Anisotropy	0.401	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 41.4	EDS
L-test for twinning ²	$< L > = 0.48$, $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	10486	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section:
SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/2141	0.63	0/2948
1	B	0.41	0/2149	0.65	1/2958 (0.0%)
1	C	0.41	0/2149	0.65	0/2958
2	J	0.40	0/1244	0.66	0/1681
2	L	0.47	0/1233	0.74	0/1667
2	M	0.37	0/1222	0.65	0/1653
All	All	0.42	0/10138	0.66	1/13865 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1390	LEU	CB-CG-CD2	-6.09	100.65	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2093	0	2033	13	0
1	B	2101	0	2042	17	0
1	C	2101	0	2042	20	0
2	J	1226	0	1251	16	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	L	1215	0	1241	13	0
2	M	1204	0	1228	14	0
3	A	5	0	0	0	0
3	L	5	0	0	0	0
4	A	124	0	0	2	0
4	B	129	0	0	5	0
4	C	125	0	0	6	0
4	J	35	0	0	2	0
4	L	75	0	0	4	0
4	M	48	0	0	4	0
All	All	10486	0	9837	89	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (89) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:102:GLU:OE1	4:L:301:HOH:O	1.82	0.97
2:M:86:GLN:OE1	4:M:201:HOH:O	1.99	0.79
1:C:1368:ASN:OD1	1:C:1445:GLN:NE2	2.19	0.76
2:J:92:THR:O	2:J:111:LYS:NZ	2.24	0.70
2:M:118:GLN:NE2	4:M:204:HOH:O	2.26	0.68
2:M:15:LEU:HD22	2:M:29:LYS:HD3	1.75	0.66
1:B:1368:ASN:O	4:B:1501:HOH:O	2.14	0.65
1:C:1424:GLU:OE1	1:C:1444:ARG:NH2	2.29	0.65
1:C:1197:ARG:NH2	4:C:1505:HOH:O	2.31	0.63
1:C:1385:ASP:OD2	2:M:127:GLN:N	2.28	0.62
1:A:1424:GLU:OE1	1:A:1444:ARG:NH2	2.32	0.62
1:B:1399:GLU:O	1:B:1402:VAL:HG23	2.00	0.62
1:C:1365:ARG:NH2	4:C:1507:HOH:O	2.33	0.61
2:M:7:THR:HG22	2:M:71:LEU:HB3	1.83	0.61
1:C:1363:ASP:OD2	4:C:1501:HOH:O	2.16	0.61
1:B:1282:SER:HB2	1:B:1326:THR:HG22	1.83	0.60
1:B:1385:ASP:HB3	2:J:120:ARG:HH12	1.67	0.59
1:B:1279:THR:HG22	1:B:1281:SER:H	1.68	0.59
1:C:1446:LYS:O	4:C:1502:HOH:O	2.16	0.59
2:J:3:ILE:HG12	2:J:15:LEU:HB2	1.84	0.59
2:M:3:ILE:HG22	2:M:67:LEU:HB3	1.86	0.57
1:C:1432:VAL:HG22	1:C:1437:GLU:HG2	1.85	0.57
1:B:1436:HIS:ND1	4:B:1506:HOH:O	2.32	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:J:42:ARG:HD2	2:J:74:ARG:HG2	1.85	0.57
1:A:1181:HIS:NE2	1:A:1183:GLU:OE1	2.28	0.57
1:C:1283:ILE:HD11	1:C:1336:ILE:HD13	1.85	0.57
2:L:66:LYS:NZ	4:L:305:HOH:O	2.37	0.57
2:M:85:THR:HG22	2:M:150:LEU:HD23	1.85	0.56
2:L:150:LEU:O	2:L:151:ARG:NH1	2.37	0.55
2:L:120:ARG:HE	2:L:122:ILE:HD11	1.73	0.54
2:J:48:LYS:HE3	2:J:135:SER:HB2	1.91	0.53
1:B:1176:PRO:HB3	1:B:1257:VAL:O	2.10	0.52
1:C:1311:ILE:HD13	2:M:145:HIS:HB3	1.90	0.51
1:C:1200:THR:HB	1:C:1203:ILE:HD11	1.92	0.51
1:B:1174:LEU:HD12	1:B:1200:THR:HG21	1.91	0.51
2:J:122:ILE:HG13	2:J:151:ARG:HD2	1.93	0.51
2:J:100:THR:HA	2:J:133:THR:HA	1.92	0.51
2:L:131:GLY:N	4:L:301:HOH:O	2.13	0.49
1:A:1394:SER:HB3	1:A:1402:VAL:HA	1.93	0.49
1:C:1299:ARG:HD3	1:C:1315:PHE:CE2	2.48	0.49
2:J:3:ILE:HD11	2:J:15:LEU:HD12	1.95	0.49
2:M:54:ARG:NE	4:M:203:HOH:O	2.25	0.48
1:A:1389:PHE:HB2	1:A:1407:ILE:HB	1.95	0.48
1:A:1409:PRO:HG2	2:L:120:ARG:NH1	2.28	0.48
1:C:1203:ILE:HG22	1:C:1205:GLY:H	1.77	0.48
1:B:1329:GLU:OE2	4:B:1503:HOH:O	2.20	0.47
2:J:99:ASP:OD2	2:J:107:LYS:NZ	2.39	0.47
2:L:8:ASP:HA	4:L:366:HOH:O	2.12	0.47
2:L:22:THR:HG22	2:L:25:ASN:HB2	1.95	0.47
1:B:1432:VAL:HG22	1:B:1437:GLU:HG2	1.96	0.46
1:A:1219:ASN:ND2	4:A:1617:HOH:O	2.49	0.46
1:A:1428:SER:HA	1:A:1441:LEU:O	2.16	0.46
1:C:1174:LEU:HD23	1:C:1174:LEU:HA	1.55	0.46
1:A:1176:PRO:HB3	1:A:1257:VAL:O	2.17	0.45
1:C:1365:ARG:HB2	1:C:1377:THR:HG22	1.97	0.45
1:A:1174:LEU:HD12	1:A:1200:THR:HG21	1.98	0.45
1:B:1331:GLY:O	1:B:1355:GLN:NE2	2.47	0.45
1:C:1176:PRO:HB3	1:C:1257:VAL:O	2.17	0.45
1:C:1405:LEU:HD13	1:C:1414:VAL:HG21	1.98	0.45
2:J:24:GLU:CD	2:J:52:ASP:HB3	2.38	0.45
1:A:1334:TYR:O	1:A:1353:THR:HA	2.16	0.44
1:B:1200:THR:HB	1:B:1203:ILE:HD11	1.99	0.44
2:M:100:THR:HA	2:M:133:THR:HA	1.98	0.44
2:J:107:LYS:NZ	4:J:202:HOH:O	2.23	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:22:THR:HG23	2:L:25:ASN:H	1.82	0.44
1:C:1368:ASN:ND2	4:C:1502:HOH:O	2.50	0.44
2:M:60:ASN:HA	4:M:227:HOH:O	2.18	0.44
1:C:1335:ASP:OD1	1:C:1353:THR:HG22	2.18	0.44
1:B:1183:GLU:OE1	4:B:1504:HOH:O	2.21	0.43
1:B:1242:GLU:HA	1:B:1264:ILE:HG12	1.99	0.43
2:M:47:GLY:O	2:M:140:ASP:HB2	2.18	0.43
2:L:48:LYS:HE2	2:L:135:SER:OG	2.19	0.42
2:J:132:ARG:NH1	4:J:204:HOH:O	2.50	0.42
2:M:122:ILE:HB	2:M:149:ARG:HB2	2.00	0.42
2:J:123:TRP:HZ2	2:J:143:PHE:CZ	2.37	0.42
1:A:1390:LEU:HD12	1:A:1390:LEU:HA	1.92	0.42
1:B:1187:ASP:OD1	4:B:1505:HOH:O	2.22	0.41
1:A:1270:LEU:HD13	1:A:1287:TRP:CG	2.55	0.41
2:J:39:ASP:OD1	2:J:40:GLN:HG2	2.21	0.41
1:C:1277:ASP:HB2	4:C:1527:HOH:O	2.20	0.41
2:L:139:ILE:HG23	2:L:146:LEU:HD13	2.03	0.41
1:B:1279:THR:HG22	1:B:1280:ASP:N	2.35	0.41
2:J:101:ILE:HB	2:J:130:ASP:HA	2.02	0.41
2:M:3:ILE:HG21	2:M:3:ILE:HD13	1.84	0.41
2:L:3:ILE:HD11	2:L:17:VAL:HG21	2.03	0.41
2:L:115:PRO:HA	2:L:116:PRO:HD3	1.98	0.41
1:A:1358:VAL:HG13	4:A:1636:HOH:O	2.21	0.40
1:B:1203:ILE:HG22	1:B:1205:GLY:H	1.85	0.40
2:J:115:PRO:HA	2:J:116:PRO:HD3	1.99	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	272/275 (99%)	265 (97%)	7 (3%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	B	273/275 (99%)	264 (97%)	8 (3%)	1 (0%)	34 41
1	C	273/275 (99%)	267 (98%)	6 (2%)	0	100 100
2	J	151/153 (99%)	150 (99%)	1 (1%)	0	100 100
2	L	149/153 (97%)	148 (99%)	1 (1%)	0	100 100
2	M	148/153 (97%)	146 (99%)	2 (1%)	0	100 100
All	All	1266/1284 (99%)	1240 (98%)	25 (2%)	1 (0%)	51 63

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	1278	ILE

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	248/249 (100%)	246 (99%)	2 (1%)	81 90
1	B	249/249 (100%)	247 (99%)	2 (1%)	81 90
1	C	249/249 (100%)	247 (99%)	2 (1%)	81 90
2	J	135/135 (100%)	129 (96%)	6 (4%)	28 39
2	L	135/135 (100%)	133 (98%)	2 (2%)	65 79
2	M	134/135 (99%)	134 (100%)	0	100 100
All	All	1150/1152 (100%)	1136 (99%)	14 (1%)	71 83

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1203	ILE
1	A	1280	ASP
1	B	1282	SER
1	B	1353	THR
1	C	1299	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	1377	THR
2	J	14	THR
2	J	16	GLU
2	J	57	SER
2	J	93	LEU
2	J	135	SER
2	J	144	LEU
2	L	146	LEU
2	L	151	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	1354	GLN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	A	1501	-	4,4,4	0.22	0	6,6,6	0.23	0
3	SO4	L	201	-	4,4,4	0.08	0	6,6,6	0.18	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	274/275 (99%)	0.09	18 (6%) 18 24	20, 32, 65, 98	1 (0%)
1	B	275/275 (100%)	0.13	16 (5%) 23 29	20, 33, 60, 99	0
1	C	275/275 (100%)	0.30	25 (9%) 9 12	20, 37, 79, 100	0
2	J	153/153 (100%)	0.17	4 (2%) 56 63	29, 47, 71, 96	0
2	L	151/153 (98%)	-0.19	0 100 100	22, 33, 44, 78	0
2	M	150/153 (98%)	0.04	1 (0%) 87 91	24, 44, 62, 90	0
All	All	1278/1284 (99%)	0.12	64 (5%) 28 36	20, 36, 69, 100	1 (0%)

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	1367	THR	5.5
1	B	1173	MET	4.9
1	A	1371	PRO	4.8
1	C	1306	GLY	4.4
1	C	1423	THR	4.1
1	B	1397	LYS	4.0
1	C	1369	ILE	4.0
1	B	1400	GLU	4.0
1	C	1368	ASN	3.8
1	C	1253	ASP	3.8
1	C	1425	TYR	3.6
1	C	1173	MET	3.6
1	A	1253	ASP	3.5
2	J	110	ASP	3.5
1	A	1217	GLN	3.5
1	A	1368	ASN	3.4
1	B	1401	ASP	3.3
1	A	1370	GLY	3.3
1	A	1447	THR	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	1420	LEU	3.2
1	A	1420	LEU	3.1
1	A	1397	LYS	3.0
1	C	1446	LYS	3.0
1	C	1371	PRO	3.0
1	C	1372	ASP	2.9
1	C	1251	LYS	2.9
1	A	1203	ILE	2.8
1	C	1447	THR	2.8
1	A	1202	ASP	2.8
1	C	1398	ASN	2.8
1	A	1199	THR	2.8
1	C	1201	PRO	2.7
1	B	1422	GLY	2.6
1	B	1447	THR	2.6
1	B	1421	PRO	2.6
1	C	1421	PRO	2.6
1	B	1398	ASN	2.6
1	A	1204	THR	2.5
1	A	1251	LYS	2.5
1	C	1399	GLU	2.5
1	B	1370	GLY	2.4
1	A	1446	LYS	2.4
1	A	1400	GLU	2.4
1	C	1252	ASP	2.3
1	C	1204	THR	2.3
1	C	1400	GLU	2.3
1	A	1218	GLY	2.3
2	J	86	GLN	2.3
2	M	111	LYS	2.3
1	B	1278	ILE	2.2
1	A	1398	ASN	2.2
1	B	1201	PRO	2.2
1	B	1396	VAL	2.2
1	C	1397	LYS	2.2
1	C	1426	VAL	2.2
1	C	1205	GLY	2.1
1	B	1202	ASP	2.1
1	B	1275	PHE	2.1
2	J	108	ILE	2.1
1	B	1420	LEU	2.0
2	J	113	GLY	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1425	TYR	2.0
1	B	1286	ARG	2.0
1	C	1271	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	A	1501	5/5	0.92	0.30	55,55,63,69	0
3	SO4	L	201	5/5	0.98	0.11	39,42,44,52	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.