

Full wwPDB X-ray Structure Validation Report (i)

Dec 5, 2023 - 06:08 am GMT

PDB ID	:	8PFC
Title	:	Crystal structure of binary complex between Aster yellows witches'-broom phy-
		toplasma effector SAP05 and the zinc finger domain of SPL5 from Arabidopsis
		thaliana
Authors	:	Huang, W.; Liu, Q.; Maqbool, A.; Stevenson, C.E.M.; Lawson, D.M.; Kamoun,
		S.; Hogenhout, S.A.
Deposited on	:	2023-06-15
Resolution	:	2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\;DIFFRACTION$

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Motria	Whole archive	Similar resolution
wietric	$(\# { m Entries})$	$(\# { m Entries}, { m resolution} { m range}({ m \AA}))$
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594(2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	105	85%	8%	8%
1	С	105	88%	•	9%
1	Е	105	84%	9%	8%
1	G	105	84%	9%	8%



Mol	Chain	Length	Quality of chain		
WIOI	Chan	Length			
1	Ι	105	87%	6%	8%
1	Κ	105	.% 87 %	6%	8%
1	М	105	2% 9 0%	•	8%
1	О	105	88%	5%	8%
2	В	70	83%	13%	•
2	D	70	86%	10%	•
2	F	70	86%	10%	•
2	Н	70	% 87 %	10%	, 0 •
2	J	70	83%	11%	6%
2	L	70	90%	1	.0%
2	Ν	70	86%	9%	6%
2	Р	70	86%	9%	6%



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 11073 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Sequence-variable mosaic (SVM) signal sequence domain-containing protein.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	Λ	07	Total	С	Ν	0	S	0	0	0
	A	91	801	519	130	149	3	0	0	0
1	C	06	Total	С	Ν	0	S	0	0	0
	C	90	795	517	129	146	3	0	0	0
1	F	07	Total	С	Ν	0	S	0	0	0
	E	97	798	518	130	147	3	0	0	0
1	С	G 97	Total	С	Ν	0	S	0	0	0
	G		801	519	130	149	3	0	0	0
1	т	07	Total	С	Ν	0	S	0	0	0
1	1	. 97	793	514	129	147	3	0	0	0
1	K	07	Total	С	Ν	0	S	0	0	0
1	Γ	91	797	517	130	147	3	0	0	0
1	М	07	Total	С	Ν	0	S	0	0	0
1	111	91	801	519	130	149	3	0	0	0
1	0	07	Total	С	Ν	0	S	0	0	0
	U	91	800	520	130	147	3		0	U

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	31	GLY	-	expression tag	UNP Q2NK94
А	32	PRO	-	expression tag	UNP Q2NK94
С	31	GLY	-	expression tag	UNP Q2NK94
С	32	PRO	-	expression tag	UNP Q2NK94
Е	31	GLY	-	expression tag	UNP Q2NK94
E	32	PRO	-	expression tag	UNP Q2NK94
G	31	GLY	-	expression tag	UNP Q2NK94
G	32	PRO	-	expression tag	UNP Q2NK94
Ι	31	GLY	-	expression tag	UNP Q2NK94
Ι	32	PRO	-	expression tag	UNP Q2NK94
K	31	GLY	-	expression tag	UNP Q2NK94
K	32	PRO	_	expression tag	UNP Q2NK94



continuou fronte proceduo pugoni							
Chain	Residue	Modelled	Actual	Comment	Reference		
М	31	GLY	-	expression tag	UNP Q2NK94		
М	32	PRO	-	expression tag	UNP Q2NK94		
0	31	GLY	-	expression tag	UNP Q2NK94		
0	32	PRO	-	expression tag	UNP Q2NK94		

• Molecule 2 is a protein called Squamosa promoter-binding-like protein 5.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
9	В	67	Total	С	Ν	0	S	0	0	0
	D	07	549	333	115	95	6	0	0	0
9	Л	67	Total	С	Ν	Ο	S	0	0	0
	D	07	543	330	112	95	6	0	0	0
0	Б	67	Total	С	Ν	Ο	S	0	0	0
	Г	07	545	331	115	93	6	0	0	0
0	ц	68	Total	С	Ν	Ο	S	0	0	0
	11	00	555	336	116	97	6			
0	т	66	Total	С	Ν	0	S	0	0	0
	J	00	531	321	110	94	6	0	0	0
9	т	70	Total	С	Ν	Ο	\mathbf{S}	0	0	0
		10	566	343	118	99	6	0	0	0
9	N	66	Total	С	Ν	Ο	\mathbf{S}	0	0	0
		00	532	324	108	94	6	0	0	0
2	р	66	Total	С	N	0	S	0	0	0
	Р	00	529	321	108	94	6	U		U

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled Actual		Comment	Reference
В	58	GLY	-	expression tag	UNP $Q9S758$
D	58	GLY	-	expression tag	UNP $Q9S758$
F	58	GLY	-	expression tag	UNP $Q9S758$
Н	58	GLY	-	expression tag	UNP $Q9S758$
J	58	GLY	-	expression tag	UNP $Q9S758$
L	58	GLY	-	expression tag	UNP $Q9S758$
N	58	GLY	-	expression tag	UNP $Q9S758$
Р	58	GLY	-	expression tag	UNP Q9S758

• Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	Total Zn 1 1	0	0
3	В	2	Total Zn 2 2	0	0
3	D	2	Total Zn 2 2	0	0
3	Е	1	Total Zn 1 1	0	0
3	F	2	Total Zn 2 2	0	0
3	Н	3	Total Zn 3 3	0	0
3	J	2	Total Zn 2 2	0	0
3	L	3	Total Zn 3 3	0	0
3	Ν	2	Total Zn 2 2	0	0
3	Р	2	Total Zn 2 2	0	0

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	18	Total O 18 18	0	0
4	В	27	TotalO2727	0	0
4	С	12	Total O 12 12	0	0
4	D	18	Total O 18 18	0	0
4	Е	14	Total O 14 14	0	0
4	F	27	TotalO2727	0	0
4	G	20	TotalO2020	0	0
4	Н	35	$\begin{array}{cc} \text{Total} & \text{O} \\ 35 & 35 \end{array}$	0	0
4	Ι	14	Total O 14 14	0	0
4	J	13	Total O 13 13	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	K	16	Total O 16 16	0	0
4	L	33	Total O 33 33	0	0
4	М	17	Total O 17 17	0	0
4	Ν	21	Total O 21 21	0	0
4	О	21	Total O 21 21	0	0
4	Р	11	Total O 11 11	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Sequence-variable mosaic (SVM) signal sequence domain-containing protein



• Molecule 1: Sequence-variable mosaic (SVM) signal sequence domain-containing protein



• Molecule 1: Sequence-variable mosaic (SVM) signal sequence domain-containing protein



• Molecule 1: Sequence-variable mosaic (SVM) signal sequence domain-containing protein



• Molecule 1: Sequence-variable mosaic (SVM) signal sequence domain-containing protein







Chain J:	83%	11% 6%
QLY PRO SFRO RG7 CG8 RG7 CG8 RG7 CG8 RG7 CG8 RG7 CG8 RG7 CG8 RG7 CG8 CG8 CG8 CG8 CG8 CG8 CG8 CG8 CG8 CG8	L113 L113 L114 LEU	
• Molecule 2: Squam	osa promoter-binding-like protein 5	
Chain L:	90%	10%
G58 V65 N71 173 173 173 173 173 113 114 114		
• Molecule 2: Squam	osa promoter-binding-like protein 5	
Chain N:	86%	9% 6%
CLY PRO SER SER SER SER SER NRG 162 173 173 173 173 173 173 173 173 173 173		
• Molecule 2: Squam	osa promoter-binding-like protein 5	
Chain P:	86%	9% 6%
RLY PRO SER SER R61 L62 L62 L62 L62 H88 H88 H88 H88 H88 H88 H88 H88 H88 H8		



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	78.69Å 165.02Å 80.86Å	Deperitor
a, b, c, α , β , γ	90.00° 109.65° 90.00°	Depositor
$\mathbf{P}_{\text{acclution}}(\hat{\mathbf{A}})$	74.21 - 2.20	Depositor
Resolution (A)	74.10 - 2.20	EDS
% Data completeness	100.0 (74.21-2.20)	Depositor
(in resolution range)	$100.0\ (74.10-2.20)$	EDS
R_{merge}	0.13	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.90 (at 2.20 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
D D	0.223 , 0.251	Depositor
$\mathbf{n}, \mathbf{n}_{free}$	0.227 , 0.253	DCC
R_{free} test set	4853 reflections $(4.94%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	44.5	Xtriage
Anisotropy	0.553	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.33 , 44.7	EDS
L-test for twinning ²	$< L >=0.51, < L^2>=0.34$	Xtriage
Estimated twinning fraction	0.000 for l,-k,h	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11073	wwPDB-VP
Average B, all atoms $(Å^2)$	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 42.95 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.8938e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond lengths		Bond angles		
	Ullalli	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.65	0/820	0.77	0/1105	
1	С	0.63	0/814	0.78	0/1097	
1	Е	0.65	0/817	0.78	0/1101	
1	G	0.66	0/820	0.86	3/1105~(0.3%)	
1	Ι	0.68	0/812	0.80	0/1096	
1	K	0.65	0/816	0.79	0/1100	
1	М	0.64	0/820	0.78	0/1105	
1	0	0.66	0/819	0.79	0/1104	
2	В	0.68	0/557	0.97	0/745	
2	D	0.67	0/551	0.96	2/738~(0.3%)	
2	F	0.68	0/553	0.91	0/740	
2	Н	0.65	0/563	0.92	0/753	
2	J	0.71	0/539	0.93	0/723	
2	L	0.66	0/575	0.91	1/770~(0.1%)	
2	N	0.68	0/540	0.95	0/724	
2	Р	0.71	0/537	0.97	0/720	
All	All	0.66	0/10953	0.86	6/14726~(0.0%)	

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
1	G	104	ARG	NE-CZ-NH1	-8.54	116.03	120.30
1	G	104	ARG	NE-CZ-NH2	7.76	124.18	120.30
2	D	102	ARG	NE-CZ-NH2	-6.40	117.10	120.30
2	D	125	ARG	NE-CZ-NH2	5.68	123.14	120.30
1	G	73	ARG	NE-CZ-NH2	5.08	122.84	120.30
2	L	81	ARG	NE-CZ-NH1	5.01	122.81	120.30

There are no chirality outliers.



There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	${ m H}({ m model})$	H(added)	Clashes	Symm-Clashes
1	А	801	0	771	4	0
1	С	795	0	774	2	0
1	Е	798	0	769	6	0
1	G	801	0	771	5	0
1	Ι	793	0	756	5	0
1	Κ	797	0	767	4	0
1	М	801	0	771	1	0
1	0	800	0	776	4	0
2	В	549	0	536	7	0
2	D	543	0	525	3	0
2	F	545	0	532	4	0
2	Н	555	0	541	5	0
2	J	531	0	503	7	0
2	L	566	0	551	3	0
2	Ν	532	0	512	3	0
2	Р	529	0	503	3	0
3	А	1	0	0	0	0
3	В	2	0	0	0	0
3	D	2	0	0	0	0
3	Е	1	0	0	0	0
3	F	2	0	0	0	0
3	Н	3	0	0	0	0
3	J	2	0	0	0	0
3	L	3	0	0	0	0
3	Ν	2	0	0	0	0
3	Р	2	0	0	0	0
4	А	18	0	0	0	0
4	В	27	0	0	0	0
4	С	12	0	0	0	0
4	D	18	0	0	0	0
4	Е	14	0	0	0	0
4	F	27	0	0	1	0
4	G	20	0	0	0	0
4	Н	35	0	0	0	0
					Continu	ied on next page

PROTEIN DATA BANK

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	Ι	14	0	0	0	0
4	J	13	0	0	0	0
4	Κ	16	0	0	0	0
4	L	33	0	0	0	0
4	М	17	0	0	0	0
4	Ν	21	0	0	0	0
4	0	21	0	0	0	0
4	Р	11	0	0	0	0
All	All	11073	0	10358	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
2:J:98:GLY:O	1:K:121:ASN:HA	1.61	1.01
2:B:107:CYS:SG	2:D:120:LYS:NZ	2.54	0.81
2:B:119:ALA:HB2	1:G:118:GLN:HE22	1.60	0.67
1:E:52:ILE:HD13	1:E:95:LEU:HB3	1.83	0.61
1:A:52:ILE:HD13	1:A:95:LEU:HB3	1.84	0.60
2:J:83:ARG:HD2	1:O:112:ILE:HD11	1.86	0.57
2:D:65:VAL:HG21	2:D:88:HIS:CG	2.42	0.55
2:N:65:VAL:HG21	2:N:88:HIS:CG	2.41	0.55
2:J:65:VAL:HG21	2:J:88:HIS:CG	2.42	0.55
2:P:65:VAL:HG21	2:P:88:HIS:CG	2.43	0.54
2:P:62:LEU:HD12	2:P:72:LEU:HB2	1.90	0.54
2:F:65:VAL:HG21	2:F:88:HIS:CG	2.44	0.53
1:I:85:TRP:CE2	1:I:106:ASP:HB2	2.43	0.53
1:A:85:TRP:CE2	1:A:106:ASP:HB2	2.44	0.53
1:G:85:TRP:CE2	1:G:106:ASP:HB2	2.44	0.53
2:B:119:ALA:HB2	1:G:118:GLN:NE2	2.23	0.53
1:K:85:TRP:CE2	1:K:106:ASP:HB2	2.44	0.53
1:C:85:TRP:CE2	1:C:106:ASP:HB2	2.44	0.53
2:L:65:VAL:HG21	2:L:88:HIS:CG	2.43	0.52
1:M:85:TRP:CE2	1:M:106:ASP:HB2	2.45	0.52
1:O:85:TRP:CE2	1:0:106:ASP:HB2	2.43	0.52
2:B:65:VAL:HG21	2:B:88:HIS:CG	2.45	0.52
2:H:65:VAL:HG21	2:H:88:HIS:CG	2.43	0.52
2:H:126:ARG:CB	1:I:119:ILE:HG22	2.40	0.52
1:E:85:TRP:CE2	1:E:106:ASP:HB2	2.44	0.51



A + am 1	Adam 0	Interatomic	Clash
Atom-1	Atom-2	distance (\AA)	overlap (Å)
1:I:114:GLU:OE1	1:I:114:GLU:N	2.39	0.51
1:E:55:LEU:HD13	1:E:65:PHE:CE2	2.46	0.50
1:A:55:LEU:HD13	1:A:65:PHE:CE2	2.47	0.49
1:A:51:ASN:HA	1:A:131:GLN:HE21	1.76	0.49
1:G:51:ASN:HA	1:G:131:GLN:HE21	1.77	0.49
1:I:52:ILE:HD13	1:I:95:LEU:HD13	1.95	0.47
1:G:55:LEU:HD13	1:G:65:PHE:CE2	2.49	0.47
1:E:51:ASN:HA	1:E:131:GLN:HE21	1.79	0.47
1:K:55:LEU:HD13	1:K:65:PHE:CE2	2.49	0.46
1:E:107:GLU:HG3	2:J:69:THR:HB	1.96	0.46
1:C:92:LEU:H	1:C:92:LEU:CD1	2.28	0.46
2:J:113:LEU:N	2:J:114:PRO:CD	2.79	0.46
2:D:113:LEU:N	2:D:114:PRO:CD	2.79	0.45
2:J:67:ARG:HD2	1:K:40:GLY:O	2.16	0.45
2:P:113:LEU:N	2:P:114:PRO:CD	2.80	0.45
2:F:113:LEU:N	2:F:114:PRO:CD	2.80	0.45
2:N:113:LEU:N	2:N:114:PRO:CD	2.80	0.45
2:H:113:LEU:N	2:H:114:PRO:CD	2.80	0.44
2:B:113:LEU:N	2:B:114:PRO:CD	2.80	0.44
2:L:113:LEU:N	2:L:114:PRO:CD	2.81	0.43
2:F:66:ASP:HA	4:F:318:HOH:O	2.19	0.43
1:E:52:ILE:HD12	1:E:52:ILE:HA	1.95	0.41
1:O:52:ILE:HD12	1:O:55:LEU:HD12	2.01	0.41
2:F:71:ASN:OD1	2:F:73:THR:HB	2.21	0.41
2:B:71:ASN:OD1	2:B:73:THR:HB	2.21	0.41
2:L:71:ASN:OD1	2:L:73:THR:HB	2.20	0.41
2:B:100:ARG:HA	2:B:100:ARG:HD2	1.94	0.41
2:H:71:ASN:OD1	2:H:73:THR:HB	2.20	0.41
2:H:126:ARG:HB2	1:I:119:ILE:HG22	2.02	0.41
2:J:83:ARG:HD2	1:O:112:ILE:CD1	2.51	0.41
2:N:71:ASN:OD1	2:N:73:THR:HB	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



8PFC

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	95/105~(90%)	94~(99%)	1 (1%)	0	100	100
1	С	94/105~(90%)	93~(99%)	1 (1%)	0	100	100
1	Е	95/105~(90%)	94 (99%)	1 (1%)	0	100	100
1	G	95/105~(90%)	94 (99%)	1 (1%)	0	100	100
1	Ι	95/105~(90%)	94 (99%)	1 (1%)	0	100	100
1	Κ	95/105~(90%)	94 (99%)	1 (1%)	0	100	100
1	М	95/105~(90%)	94 (99%)	1 (1%)	0	100	100
1	Ο	95/105~(90%)	94 (99%)	1 (1%)	0	100	100
2	В	65/70~(93%)	64 (98%)	1 (2%)	0	100	100
2	D	65/70~(93%)	65~(100%)	0	0	100	100
2	F	65/70~(93%)	65~(100%)	0	0	100	100
2	Н	66/70~(94%)	65~(98%)	1 (2%)	0	100	100
2	J	64/70~(91%)	63~(98%)	1 (2%)	0	100	100
2	L	68/70~(97%)	67~(98%)	1 (2%)	0	100	100
2	Ν	64/70~(91%)	64 (100%)	0	0	100	100
2	Р	$\overline{64}/70~(91\%)$	64 (100%)	0	0	100	100
All	All	1280/1400~(91%)	1268 (99%)	12 (1%)	0	100	100

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	\mathbf{ntiles}
1	А	85/93~(91%)	85 (100%)	0	100	100
1	С	85/93~(91%)	84 (99%)	1 (1%)	71	83
1	Е	84/93~(90%)	84 (100%)	0	100	100



Mol	Chain	Analysed	Rotameric	Outliers	Percentile	
1	G	85/93~(91%)	85 (100%)	0	100	100
1	Ι	83/93~(89%)	83 (100%)	0	100	100
1	К	84/93~(90%)	84 (100%)	0	100	100
1	М	85/93~(91%)	85 (100%)	0	100	100
1	Ο	85/93~(91%)	85 (100%)	0	100	100
2	В	58/61~(95%)	58 (100%)	0	100	100
2	D	57/61~(93%)	57 (100%)	0	100	100
2	F	57/61~(93%)	57 (100%)	0	100	100
2	Н	59/61~(97%)	59 (100%)	0	100	100
2	J	55/61~(90%)	55 (100%)	0	100	100
2	L	60/61~(98%)	60 (100%)	0	100	100
2	Ν	56/61~(92%)	56 (100%)	0	100	100
2	Р	55/61~(90%)	55 (100%)	0	100	100
All	All	1133/1232 (92%)	1132 (100%)	1 (0%)	93	97

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	С	57	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	89	ASN
1	А	131	GLN
1	Е	131	GLN
1	G	118	GLN
1	G	131	GLN
1	0	89	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 20 ligands modelled in this entry, 20 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$\mathbf{OWAB}(\mathbf{\AA}^2)$	Q<0.9
1	А	97/105~(92%)	0.28	0 100 100	46, 68, 100, 107	0
1	С	96/105~(91%)	0.74	10 (10%) 6 5	52, 71, 102, 119	0
1	Е	97/105~(92%)	0.23	0 100 100	48, 68, 98, 105	0
1	G	97/105~(92%)	0.08	0 100 100	44, 58, 82, 94	0
1	Ι	97/105~(92%)	1.14	18 (18%) 1 1	44, 74, 104, 127	0
1	Κ	97/105~(92%)	0.18	1 (1%) 82 81	45, 68, 100, 108	0
1	М	97/105~(92%)	0.39	2 (2%) 63 61	38, 66, 94, 113	0
1	Ο	97/105~(92%)	0.73	10 (10%) 6 5	41, 66, 93, 126	0
2	В	67/70~(95%)	0.33	0 100 100	41, 55, 83, 123	0
2	D	67/70~(95%)	0.31	3 (4%) 33 32	48, 65, 102, 125	0
2	F	67/70~(95%)	0.37	0 100 100	40, 54, 76, 85	0
2	Н	68/70~(97%)	0.27	1 (1%) 73 72	46, 58, 81, 108	0
2	J	66/70~(94%)	0.33	1 (1%) 73 72	43, 66, 104, 112	0
2	L	70/70~(100%)	0.17	0 100 100	46, 57, 77, 103	0
2	N	66/70~(94%)	0.16	0 100 100	43, 66, 91, 113	0
2	Р	$6\overline{6}/70~(94\%)$	0.26	0 100 100	43, 65, 92, 99	0
All	All	1312/1400 (93%)	0.39	46 (3%) 44 42	38, 65, 97, 127	0

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	Ι	36	GLU	6.2
1	Ι	44	ILE	6.2
1	Ι	38	PHE	5.6
1	0	36	GLU	5.5
1	Ι	37	GLU	3.7



Mol	Chain	Res	Type	RSRZ
1	0	38	PHE	3.6
1	0	44	ILE	3.5
1	0	37	GLU	3.5
1	С	92	LEU	3.5
2	J	61	ARG	3.4
1	Ι	119	ILE	3.4
1	Ι	112	ILE	3.3
1	Ι	97	VAL	3.1
1	С	127	TYR	3.0
1	Ι	70	THR	3.0
1	С	113	LEU	2.9
1	С	124	LYS	2.7
1	С	132	TYR	2.7
1	Ι	42	MET	2.7
2	D	81	ARG	2.6
1	С	52	ILE	2.6
2	D	127	LEU	2.6
1	Κ	65	PHE	2.4
1	Ι	43	ARG	2.4
1	0	39	VAL	2.4
1	Ι	67	PHE	2.3
1	С	49	LEU	2.3
1	С	61	PHE	2.2
2	Н	118	GLU	2.2
2	D	120	LYS	2.2
1	0	49	LEU	2.2
1	Ι	99	PHE	2.2
1	0	43	ARG	2.2
1	С	38	PHE	2.2
1	М	52	ILE	2.2
1	Ι	128	ILE	2.1
1	Ι	39	VAL	2.1
1	Ι	92	LEU	2.1
1	М	127	TYR	2.1
1	Ι	89	ASN	2.1
1	Ο	119	ILE	2.1
1	Ο	113	LEU	2.1
1	0	99	PHE	2.1
1	Ι	91	PRO	2.1
1	С	120	GLY	2.1
1	Ι	81	PHE	2.1

Continued from previous page...



6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
3	ZN	Н	201	1/1	0.92	0.05	121,121,121,121	0
3	ZN	А	201	1/1	0.98	0.17	66, 66, 66, 66	0
3	ZN	J	201	1/1	0.98	0.20	$47,\!47,\!47,\!47$	0
3	ZN	J	202	1/1	0.98	0.15	$66,\!66,\!66,\!66$	0
3	ZN	L	201	1/1	0.98	0.10	76,76,76,76	0
3	ZN	Р	201	1/1	0.98	0.17	$51,\!51,\!51,\!51$	0
3	ZN	F	202	1/1	0.99	0.18	54,54,54,54	0
3	ZN	В	202	1/1	0.99	0.18	$63,\!63,\!63,\!63$	0
3	ZN	Н	202	1/1	0.99	0.20	48,48,48,48	0
3	ZN	D	201	1/1	0.99	0.17	56, 56, 56, 56	0
3	ZN	D	202	1/1	0.99	0.15	60,60,60,60	0
3	ZN	Е	201	1/1	0.99	0.10	83,83,83,83	0
3	ZN	L	202	1/1	0.99	0.20	$50,\!50,\!50,\!50$	0
3	ZN	L	203	1/1	0.99	0.17	54,54,54,54	0
3	ZN	N	201	1/1	0.99	0.17	$53,\!53,\!53,\!53$	0
3	ZN	F	201	1/1	0.99	0.19	48,48,48,48	0
3	ZN	Р	202	1/1	0.99	0.15	$61,\!61,\!61,\!61$	0
3	ZN	N	202	1/1	1.00	0.14	62,62,62,62	0
3	ZN	В	201	1/1	1.00	0.17	47,47,47,47	0
3	ZN	Н	203	1/1	1.00	0.18	53,53,53,53	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

















































































6.5 Other polymers (i)

There are no such residues in this entry.

