



Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 04:07 pm BST

PDB ID : 1PGW
Title : BEAN POD MOTTLE VIRUS (BPMV), TOP COMPONENT
Authors : Lin, T.; Cavarelli, J.; Johnson, J.E.
Deposited on : 2003-05-28
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	1	185	 % 71% 25% 5% 1%
2	2	351	 % 76% 19% 5%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4306 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BEAN POD MOTTLE VIRUS SMALL (S) SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	1	185	1452	924	244	275	9	0	0	0

- Molecule 2 is a protein called BEAN POD MOTTLE VIRUS LARGE (L) SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	2	351	2704	1727	453	497	27	0	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	1	49	Total	O	0	0
			49	49		
3	2	101	Total	O	0	0
			101	101		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: BEAN POD MOTTLE VIRUS SMALL (S) SUBUNIT



- Molecule 2: BEAN POD MOTTLE VIRUS LARGE (L) SUBUNIT



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, α , β , γ	311.20Å 284.20Å 350.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.90 29.95 – 2.90	Depositor EDS
% Data completeness (in resolution range)	46.4 (10.00-2.90) 46.4 (29.95-2.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.86 (at 2.90Å)	Xtrriage
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.212 , (Not available) 0.199 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	26.0	Xtrriage
Anisotropy	0.236	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 60.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.26$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	4306	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	1	0.72	0/1491	0.82	2/2031 (0.1%)
2	2	0.73	2/2765 (0.1%)	0.85	2/3752 (0.1%)
All	All	0.73	2/4256 (0.0%)	0.84	4/5783 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	2	20	MET	SD-CE	7.39	2.19	1.77
2	2	132	CYS	CB-SG	-6.17	1.71	1.82

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	2	243	VAL	CB-CA-C	-6.26	99.51	111.40
2	2	23	LEU	CA-CB-CG	-6.06	101.36	115.30
1	1	42	ASP	CB-CA-C	-5.36	99.68	110.40
1	1	143	THR	N-CA-CB	-5.31	100.20	110.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	1	1452	0	1405	52	0
2	2	2704	0	2715	56	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	1	49	0	0	2	0
3	2	101	0	0	5	0
All	All	4306	0	4120	102	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (102) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:2:20:MET:CE	2:2:20:MET:SD	2.19	1.31
1:1:21:ASP:HB3	1:1:24:LYS:HD3	1.59	0.84
1:1:16:THR:HG21	1:1:154:PRO:O	1.82	0.79
2:2:140:PHE:O	2:2:144:THR:HB	1.81	0.79
1:1:42:ASP:HB3	1:1:44:THR:H	1.52	0.74
2:2:200:LEU:HG	2:2:211:LYS:HG2	1.73	0.69
1:1:21:ASP:O	1:1:24:LYS:HE2	1.93	0.68
1:1:21:ASP:H	1:1:24:LYS:NZ	1.91	0.68
1:1:15:ARG:NH1	1:1:16:THR:HG22	2.10	0.67
2:2:294:LYS:HD2	3:2:437:HOH:O	1.95	0.66
1:1:21:ASP:HB3	1:1:24:LYS:CD	2.26	0.65
1:1:69:CYS:O	1:1:167:SER:HB3	2.00	0.61
2:2:92:MET:HG3	2:2:146:PHE:HE1	1.66	0.59
1:1:143:THR:HG23	1:1:144:SER:N	2.18	0.59
1:1:97:ASN:HB2	1:1:143:THR:CG2	2.33	0.58
1:1:1:SER:HB3	1:1:168:GLN:OE1	2.03	0.57
1:1:97:ASN:HD21	1:1:103:HIS:HD2	1.50	0.57
2:2:247:VAL:HG22	2:2:298:LYS:HD3	1.87	0.56
2:2:239:TRP:O	2:2:310:THR:HG21	2.06	0.56
1:1:134:MET:HE3	2:2:144:THR:O	2.05	0.56
2:2:354:MET:HG2	2:2:355:CYS:N	2.20	0.55
2:2:113:GLU:HB3	2:2:123:MET:CE	2.37	0.55
2:2:20:MET:O	2:2:23:LEU:N	2.38	0.55
2:2:240:ARG:HD3	3:2:432:HOH:O	2.06	0.54
1:1:143:THR:HG22	3:1:221:HOH:O	2.06	0.54
1:1:15:ARG:HB3	1:1:48:ILE:HB	1.90	0.54
2:2:137:SER:HB3	2:2:140:PHE:H	1.73	0.53
2:2:187:PRO:HG3	2:2:239:TRP:CZ2	2.42	0.53
1:1:21:ASP:H	1:1:24:LYS:HZ1	1.56	0.53
2:2:92:MET:HG2	2:2:127:PHE:CE2	2.44	0.52
1:1:97:ASN:HB2	1:1:143:THR:HG23	1.91	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:1:15:ARG:HH11	1:1:16:THR:HG22	1.74	0.52
2:2:256:PRO:HB3	2:2:290:GLU:O	2.09	0.52
2:2:47:LEU:HD13	2:2:92:MET:CE	2.40	0.52
2:2:92:MET:HE3	2:2:148:CYS:HB2	1.90	0.51
2:2:99:ARG:HB3	2:2:99:ARG:HH11	1.76	0.51
2:2:113:GLU:HB3	2:2:123:MET:HE1	1.92	0.51
2:2:212:ARG:NH2	2:2:273:ASP:OD2	2.44	0.51
2:2:354:MET:HG2	2:2:355:CYS:H	1.75	0.51
2:2:243:VAL:HG22	2:2:307:ALA:HB2	1.93	0.50
1:1:130:ASN:H	1:1:132:PHE:H	1.58	0.50
1:1:28:CYS:HB2	1:1:44:THR:HG21	1.93	0.50
2:2:47:LEU:HD13	2:2:92:MET:HE2	1.94	0.49
1:1:25:GLN:NE2	1:1:27:PHE:O	2.44	0.49
1:1:6:THR:O	1:1:6:THR:HG22	2.12	0.49
2:2:114:LEU:HD21	2:2:253:MET:HE3	1.93	0.49
1:1:127:GLY:HA3	1:1:133:GLU:HB2	1.94	0.49
2:2:144:THR:CG2	3:2:396:HOH:O	2.61	0.49
1:1:42:ASP:HB2	1:1:47:TRP:HE1	1.79	0.48
2:2:72:ILE:O	2:2:126:SER:HA	2.14	0.48
2:2:164:VAL:HG12	2:2:165:GLN:N	2.29	0.48
1:1:73:LYS:HG3	1:1:120:ILE:HG12	1.94	0.48
1:1:42:ASP:HB3	1:1:44:THR:HB	1.96	0.48
2:2:83:ASN:N	2:2:83:ASN:HD22	2.11	0.48
2:2:208:ARG:NH2	2:2:275:THR:O	2.40	0.48
1:1:21:ASP:H	1:1:24:LYS:HZ3	1.60	0.47
2:2:204:GLN:HA	2:2:334:SER:HB3	1.97	0.47
1:1:143:THR:HG23	1:1:144:SER:H	1.80	0.47
2:2:230:ASN:HD22	2:2:230:ASN:C	2.18	0.47
1:1:108:ARG:HB2	1:1:108:ARG:HH11	1.79	0.46
1:1:55:PRO:HD3	2:2:362:GLY:HA3	1.96	0.46
1:1:12:ALA:HB1	1:1:39:ILE:HD11	1.96	0.46
1:1:97:ASN:N	1:1:97:ASN:HD22	2.14	0.46
1:1:97:ASN:HB2	1:1:143:THR:HG21	1.98	0.45
1:1:181:PHE:O	2:2:226:ALA:HA	2.17	0.45
2:2:243:VAL:HG22	2:2:307:ALA:CB	2.46	0.45
1:1:58:ASN:HD22	1:1:61:ARG:HH21	1.64	0.45
2:2:124:THR:HG22	2:2:125:MET:N	2.33	0.44
1:1:15:ARG:HD3	1:1:16:THR:N	2.33	0.44
2:2:23:LEU:HA	2:2:23:LEU:HD23	1.47	0.44
1:1:98:SER:HB3	1:1:103:HIS:CD2	2.52	0.44
1:1:24:LYS:HG3	1:1:24:LYS:H	1.52	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:2:144:THR:HG23	3:2:396:HOH:O	2.17	0.43
1:1:97:ASN:ND2	1:1:103:HIS:HD2	2.16	0.43
2:2:52:THR:O	2:2:52:THR:HG23	2.18	0.43
1:1:127:GLY:CA	1:1:133:GLU:HB2	2.48	0.43
2:2:103:ASP:OD2	2:2:105:TYR:HB3	2.18	0.43
2:2:83:ASN:H	2:2:83:ASN:ND2	2.17	0.43
1:1:97:ASN:HD21	1:1:103:HIS:CD2	2.35	0.42
2:2:74:MET:O	2:2:74:MET:HG3	2.20	0.42
1:1:2:ILE:HD13	2:2:239:TRP:CD1	2.55	0.42
1:1:28:CYS:HB2	1:1:42:ASP:OD1	2.18	0.42
2:2:83:ASN:HD22	2:2:83:ASN:H	1.67	0.42
1:1:180:ALA:HB1	2:2:226:ALA:HB1	2.01	0.42
2:2:313:ARG:HD2	3:2:458:HOH:O	2.19	0.42
2:2:193:LEU:HD21	2:2:217:ILE:HD12	2.02	0.41
2:2:318:LEU:HD12	2:2:318:LEU:HA	1.73	0.41
2:2:73:LYS:HG2	2:2:74:MET:N	2.34	0.41
1:1:20:PHE:HD1	1:1:44:THR:HG23	1.86	0.41
1:1:18:LEU:HD23	1:1:18:LEU:HA	1.91	0.41
2:2:258:ILE:HA	2:2:336:SER:HB2	2.01	0.41
2:2:190:PRO:HB3	2:2:352:GLU:HG2	2.02	0.41
1:1:143:THR:CG2	3:1:221:HOH:O	2.68	0.41
2:2:221:ALA:HB1	2:2:369:LEU:HD22	2.03	0.41
2:2:222:GLY:C	2:2:368:LEU:HD23	2.41	0.41
1:1:28:CYS:CB	1:1:44:THR:HG21	2.51	0.41
2:2:231:MET:HE2	2:2:235:VAL:CG1	2.51	0.41
1:1:42:ASP:HB2	1:1:47:TRP:NE1	2.36	0.41
1:1:97:ASN:ND2	1:1:97:ASN:H	2.19	0.41
1:1:183:VAL:HG23	2:2:226:ALA:HA	2.02	0.40
1:1:97:ASN:HD22	1:1:98:SER:N	2.20	0.40
2:2:113:GLU:HB3	2:2:123:MET:HE2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	1	183/185 (99%)	170 (93%)	12 (7%)	1 (0%)	29	61
2	2	349/351 (99%)	334 (96%)	13 (4%)	2 (1%)	25	58
All	All	532/536 (99%)	504 (95%)	25 (5%)	3 (1%)	25	58

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	1	139	TRP
2	2	78	VAL
2	2	21	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	1	165/165 (100%)	146 (88%)	19 (12%)	5	17
2	2	302/302 (100%)	270 (89%)	32 (11%)	6	20
All	All	467/467 (100%)	416 (89%)	51 (11%)	6	19

All (51) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	1	5	GLN
1	1	7	VAL
1	1	9	ASN
1	1	11	MET
1	1	15	ARG
1	1	16	THR
1	1	24	LYS
1	1	34	LEU
1	1	44	THR
1	1	52	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	1	62	VAL
1	1	84	ARG
1	1	91	VAL
1	1	96	THR
1	1	97	ASN
1	1	99	ASN
1	1	108	ARG
1	1	129	ASN
1	1	143	THR
2	2	32	LEU
2	2	42	LEU
2	2	52	THR
2	2	69	LYS
2	2	83	ASN
2	2	88	LEU
2	2	92	MET
2	2	99	ARG
2	2	123	MET
2	2	137	SER
2	2	141	LEU
2	2	144	THR
2	2	162	GLN
2	2	175	GLN
2	2	185	LEU
2	2	186	ASN
2	2	200	LEU
2	2	208	ARG
2	2	228	LEU
2	2	230	ASN
2	2	231	MET
2	2	233	ASN
2	2	243	VAL
2	2	278	PHE
2	2	297	LEU
2	2	310	THR
2	2	312	VAL
2	2	318	LEU
2	2	326	LEU
2	2	335	VAL
2	2	359	LEU
2	2	369	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such

sidechains are listed below:

Mol	Chain	Res	Type
1	1	52	GLN
1	1	58	ASN
1	1	97	ASN
1	1	99	ASN
1	1	103	HIS
1	1	129	ASN
1	1	146	HIS
2	2	49	ASN
2	2	63	GLN
2	2	83	ASN
2	2	128	ASN
2	2	130	ASN
2	2	147	HIS
2	2	230	ASN
2	2	353	ASN
2	2	360	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	1	185/185 (100%)	-1.02	1 (0%) 91 91	5, 14, 40, 98	0
2	2	351/351 (100%)	-1.10	0 100 100	5, 11, 32, 87	0
All	All	536/536 (100%)	-1.07	1 (0%) 95 95	5, 12, 39, 98	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	1	185	GLN	4.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.