



Full wwPDB NMR Structure Validation Report ⓘ

Jan 10, 2022 – 12:04 am GMT

PDB ID : 7PKU
Title : Structure of SARS-CoV-2 nucleoprotein in dynamic complex with its viral partner nsp3a
Authors : Bessa, L.M.; Guseva, S.; Camacho-Zarco, A.R.; Salvi, N.; Blackledge, M.
Deposited on : 2021-08-26

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.24
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.24


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 49%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	96	
2	B	73	

2 Ensemble composition and analysis

This entry contains 10 models. Model 5 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *target function*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:17-A:106, B:219-B:232, B:245-B:254 (114)	0.70	5

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 3 single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 5, 6
2	4, 9
Single-model clusters	7; 8; 10

3 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 2586 atoms, of which 1285 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called 3C-like proteinase.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	96	1493	491	729	112	156	5	0

- Molecule 2 is a protein called Nucleoprotein.


Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
2	B	73	1093	315	556	112	108	2	0

4 Residue-property plots i


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: 3C-like proteinase



- Molecule 2: Nucleoprotein




4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

- Molecule 1: 3C-like proteinase




- Molecule 2: Nucleoprotein




4.2.2 Score per residue for model 2

- Molecule 1: 3C-like proteinase




- Molecule 2: Nucleoprotein




4.2.3 Score per residue for model 3

- Molecule 1: 3C-like proteinase



- Molecule 2: Nucleoprotein



K261
R262
T263

4.2.4 Score per residue for model 4

- Molecule 1: 3C-like proteinase

Chain A:  63% 27% 6%

Q16 G17 Y18 K19 S20 V21 N22 I23 T24 F25 I31 L35 M36 S40 A41 Y42 T43 V44 E45 L46 G47 T48 E49 F53 A60 V61 I62 R63 T64 L65 Q66 P67 V68 L71 L72 I73 P74 L79 M82 G84 E85 F86 Y107 P108 P109 D110 E111

- Molecule 2: Nucleoprotein

Chain B:  14% 11% 8% 67%

R191 N192 S193 S194 R195 M196 S197 T198 P199 S201 S202 R203 G204 T205 S206 P207 A208 R209 N210 A211 G212 N213 G214 G215 D216 A217 A218 L223 L224 D225 R226 L227 N228 Q229 L230 K233 N234 S235 G236 K237 Q238 Q239 Q240 Q241 Q242 Q243 Q244 T245 K249 S250 A251 Q252 E253 A254 S255 K256

K261 P262 Q263

4.2.5 Score per residue for model 5 (medoid)

- Molecule 1: 3C-like proteinase

Chain A:  55% 32% 6% 6%

Q16 G17 Y18 K19 S20 V21 N22 I23 T24 F25 L27 R30 I31 V34 S40 A41 Y42 T43 V44 E45 L46 G47 T48 E49 F53 V57 A58 D59 A60 V61 I62 K63 T64 V68 L71 P74 L75 G76 D91 E92 A99 S100 C104 Y107 P108 P109 D110

E111

- Molecule 2: Nucleoprotein


Chain B:  21% 10% 67%

R191 N192 S193 S194 R195 M196 S197 T198 P199 S201 S202 R203 G204 T205 S206 P207 A208 R209 N210 A211 G212 N213 G214 G215 D216 A217 A218 L219 L223 L224 D225 R226 L227 N228 Q229 L230 K233 N234 S235 G236 K237 Q238 Q239 Q240 Q241 Q242 Q243 Q244 S255 K256 P257 P258 R259 Q260 K261 R262


T263

4.2.6 Score per residue for model 6

- Molecule 1: 3C-like proteinase




- Molecule 2: Nucleoprotein




4.2.7 Score per residue for model 7

- Molecule 1: 3C-like proteinase




- Molecule 2: Nucleoprotein



4.2.8 Score per residue for model 8


- Molecule 1: 3C-like proteinase



P109
D110
E111

- Molecule 2: Nucleoprotein

Chain B:  15% 15% 67%




K257
P258
R259
Q260
K261
R262
T263

4.2.9 Score per residue for model 9


- Molecule 1: 3C-like proteinase

Chain A:  65% 22% 7% 6%



- Molecule 2: Nucleoprotein

Chain B:  19% 12% 67%




K260
K261
R262
T263

4.2.10 Score per residue for model 10


- Molecule 1: 3C-like proteinase

Chain A:  59% 30% 6%



- Molecule 2: Nucleoprotein

Chain B:  19% 11% 67%



R260
R261
R262
T263

5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 1000 calculated structures, 10 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	structure calculation	1.3

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	2
Total number of shifts	1071
Number of shifts mapped to atoms	1071
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	49%

6 Model quality i

6.1 Standard geometry i

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	712	688	687	25±4
2	B	181	199	199	8±2
All	All	8930	8870	8860	292

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:30:ARG:O	1:A:31:ILE:HG12	0.77	1.78	1	2
1:A:22:ASN:HA	1:A:40:SER:O	0.76	1.81	3	10
1:A:68:VAL:HG22	2:B:227:LEU:HD12	0.71	1.60	7	1
2:B:226:ARG:O	2:B:230:LEU:HB2	0.70	1.86	5	8
2:B:224:LEU:O	2:B:228:ASN:HB2	0.70	1.85	10	10
1:A:62:ILE:O	1:A:66:GLN:HB2	0.65	1.90	9	6
1:A:18:TYR:HA	1:A:46:LEU:HD11	0.65	1.69	10	10
1:A:87:TYR:HB2	1:A:104:CYS:SG	0.64	2.33	7	1
2:B:226:ARG:O	2:B:230:LEU:HB3	0.63	1.93	4	1
2:B:225:ASP:O	2:B:229:GLN:HB2	0.63	1.93	2	6
1:A:35:LEU:HA	1:A:39:CYS:SG	0.61	2.36	2	1
1:A:77:ILE:H	1:A:77:ILE:HD13	0.61	1.55	2	1
1:A:58:ALA:O	1:A:62:ILE:HG12	0.61	1.95	7	2

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:68:VAL:HG13	2:B:227:LEU:HD23	0.60	1.72	10	4
2:B:225:ASP:HA	2:B:229:GLN:OE1	0.60	1.96	10	1
1:A:65:LEU:HD11	2:B:223:LEU:HD13	0.59	1.73	10	1
1:A:34:VAL:HB	2:B:224:LEU:HD22	0.59	1.75	3	4
1:A:20:SER:HA	1:A:42:TYR:O	0.59	1.98	5	8
1:A:42:TYR:OH	1:A:60:ALA:HB2	0.58	1.98	9	5
1:A:67:PRO:HB2	2:B:227:LEU:HD11	0.56	1.77	7	1
1:A:36:ASN:HA	1:A:40:SER:OG	0.56	2.00	1	4
1:A:71:LEU:O	1:A:74:PRO:HD2	0.56	2.01	9	10
1:A:32:ASP:O	1:A:36:ASN:HB2	0.56	2.01	3	5
1:A:62:ILE:HG23	1:A:66:GLN:NE2	0.56	2.16	2	1
2:B:223:LEU:O	2:B:227:LEU:HB3	0.54	2.01	4	1
1:A:60:ALA:O	1:A:64:THR:HG22	0.54	2.02	7	4
1:A:27:LEU:HD13	1:A:32:ASP:HA	0.54	1.78	6	1
1:A:89:LEU:HD22	1:A:104:CYS:SG	0.54	2.42	8	5
1:A:25:PHE:HB3	1:A:104:CYS:SG	0.54	2.43	2	1
1:A:21:VAL:O	1:A:41:ALA:HA	0.54	2.03	1	3
1:A:58:ALA:O	1:A:62:ILE:HB	0.54	2.03	5	6
1:A:25:PHE:HA	1:A:104:CYS:O	0.53	2.02	1	3
1:A:90:PHE:HA	1:A:95:GLU:O	0.53	2.03	8	2
1:A:67:PRO:HB2	2:B:227:LEU:HD21	0.53	1.80	10	3
1:A:65:LEU:O	1:A:68:VAL:HG23	0.52	2.04	9	1
1:A:30:ARG:O	1:A:31:ILE:HG13	0.52	2.05	3	2
1:A:71:LEU:O	1:A:75:LEU:HD23	0.51	2.05	2	2
2:B:248:LYS:O	2:B:252:ALA:HB3	0.51	2.06	8	4
1:A:35:LEU:HD21	1:A:65:LEU:HD13	0.51	1.80	3	2
1:A:49:GLU:O	1:A:53:PHE:HB2	0.51	2.06	10	7
2:B:249:LYS:O	2:B:253:GLU:HB2	0.51	2.06	8	3
1:A:64:THR:O	2:B:227:LEU:HD13	0.51	2.06	7	2
1:A:27:LEU:HD22	1:A:30:ARG:HB2	0.50	1.84	3	1
1:A:27:LEU:HD21	1:A:35:LEU:HB2	0.50	1.81	1	1
1:A:53:PHE:O	1:A:56:VAL:HG12	0.50	2.07	2	3
1:A:31:ILE:O	1:A:34:VAL:HG22	0.50	2.06	6	2
1:A:79:LEU:O	1:A:83:SER:HB2	0.50	2.05	7	2
1:A:68:VAL:HG22	2:B:227:LEU:HD22	0.50	1.84	9	1
1:A:64:THR:O	2:B:227:LEU:HD21	0.49	2.08	4	1
1:A:38:LYS:HB2	2:B:224:LEU:HD21	0.49	1.82	9	1
1:A:46:LEU:HD13	1:A:46:LEU:N	0.49	2.23	6	10
1:A:79:LEU:HA	1:A:82:TRP:CD1	0.49	2.42	4	1
2:B:250:SER:O	2:B:254:ALA:HB3	0.49	2.08	7	2
1:A:64:THR:O	2:B:227:LEU:HD23	0.49	2.08	9	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:44:VAL:HG11	1:A:53:PHE:CE2	0.49	2.42	3	2
1:A:44:VAL:HA	1:A:48:THR:OG1	0.48	2.07	4	7
2:B:225:ASP:HA	2:B:229:GLN:NE2	0.48	2.23	2	1
1:A:22:ASN:O	1:A:101:HIS:HA	0.48	2.08	7	3
1:A:42:TYR:OH	1:A:57:VAL:HA	0.48	2.09	3	4
1:A:88:TYR:O	1:A:104:CYS:HA	0.48	2.08	6	3
1:A:44:VAL:HG11	1:A:53:PHE:CE1	0.48	2.44	1	8
1:A:62:ILE:HG23	1:A:66:GLN:OE1	0.48	2.09	7	1
1:A:24:THR:HA	1:A:36:ASN:OD1	0.47	2.09	4	1
1:A:25:PHE:O	1:A:27:LEU:HG	0.47	2.10	5	2
1:A:44:VAL:HG12	1:A:48:THR:HG21	0.47	1.87	9	6
1:A:89:LEU:HA	1:A:103:TYR:O	0.47	2.10	8	2
1:A:52:GLU:OE2	2:B:251:ALA:HA	0.47	2.10	6	1
1:A:43:THR:H	2:B:246:VAL:HG23	0.47	1.70	2	1
1:A:91:ASP:OD1	1:A:92:GLU:HG2	0.46	2.10	1	1
1:A:25:PHE:H	1:A:36:ASN:ND2	0.46	2.08	4	1
1:A:59:ASP:O	1:A:63:LYS:HB2	0.46	2.11	2	4
1:A:36:ASN:O	1:A:40:SER:HB2	0.46	2.11	3	2
2:B:224:LEU:HA	2:B:228:ASN:ND2	0.46	2.26	10	2
1:A:91:ASP:HB3	1:A:92:GLU:OE2	0.46	2.09	5	1
1:A:52:GLU:OE1	2:B:251:ALA:HA	0.46	2.11	3	1
1:A:68:VAL:O	1:A:72:LEU:HD12	0.45	2.10	3	2
1:A:42:TYR:CZ	1:A:57:VAL:HA	0.45	2.46	3	2
1:A:53:PHE:HB3	1:A:89:LEU:HD11	0.45	1.87	3	2
2:B:249:LYS:HA	2:B:253:GLU:OE2	0.45	2.12	7	1
1:A:68:VAL:CG2	2:B:227:LEU:HD22	0.45	2.42	9	1
1:A:68:VAL:HG22	2:B:227:LEU:HD23	0.44	1.88	2	2
1:A:42:TYR:CZ	1:A:56:VAL:HG22	0.44	2.47	2	3
1:A:57:VAL:O	1:A:61:VAL:HG22	0.43	2.13	3	2
2:B:223:LEU:HD12	2:B:224:LEU:N	0.43	2.29	2	1
1:A:21:VAL:HB	1:A:100:SER:O	0.43	2.13	7	2
1:A:23:ILE:HD11	1:A:42:TYR:CE2	0.43	2.48	5	1
1:A:62:ILE:HG13	1:A:66:GLN:NE2	0.43	2.29	8	2
1:A:30:ARG:NH1	1:A:76:GLY:HA3	0.42	2.29	10	1
1:A:30:ARG:NH1	2:B:219:LEU:HD12	0.42	2.29	5	1
1:A:49:GLU:OE1	1:A:50:VAL:HG22	0.42	2.14	6	1
1:A:25:PHE:CD1	1:A:35:LEU:HB3	0.42	2.50	4	1
1:A:19:LYS:HB2	1:A:44:VAL:O	0.42	2.13	6	2
1:A:69:SER:O	1:A:73:THR:HG22	0.42	2.15	1	1
1:A:43:THR:OG1	2:B:245:THR:HA	0.42	2.15	4	1
2:B:222:LEU:O	2:B:226:ARG:HB2	0.41	2.14	1	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:27:LEU:HD23	1:A:32:ASP:HB2	0.41	1.93	2	1
1:A:95:GLU:HG3	1:A:96:PHE:CD1	0.41	2.50	8	1
1:A:53:PHE:O	1:A:57:VAL:HG22	0.41	2.16	5	3
1:A:66:GLN:N	1:A:67:PRO:CD	0.41	2.83	10	2
1:A:87:TYR:HB3	1:A:105:SER:O	0.41	2.15	7	1
1:A:27:LEU:HD23	1:A:32:ASP:O	0.41	2.16	1	1
1:A:64:THR:O	2:B:227:LEU:HD11	0.41	2.15	4	1
1:A:64:THR:O	2:B:227:LEU:HD12	0.41	2.16	6	1
1:A:39:CYS:SG	2:B:224:LEU:HD11	0.41	2.56	8	1
1:A:27:LEU:HD23	1:A:32:ASP:HA	0.41	1.92	1	1
1:A:62:ILE:HD12	1:A:66:GLN:NE2	0.41	2.31	2	1
1:A:57:VAL:O	1:A:61:VAL:HG12	0.41	2.16	5	1
1:A:36:ASN:O	1:A:40:SER:OG	0.41	2.35	8	1
1:A:18:TYR:HA	1:A:46:LEU:CD1	0.41	2.45	7	1
1:A:25:PHE:O	1:A:27:LEU:HD12	0.40	2.15	2	1
1:A:71:LEU:O	1:A:75:LEU:HB2	0.40	2.16	3	1
1:A:21:VAL:HG11	1:A:99:ALA:O	0.40	2.15	5	1
1:A:34:VAL:CG2	1:A:35:LEU:N	0.40	2.84	1	1
1:A:44:VAL:O	1:A:45:GLU:HB2	0.40	2.17	3	1
1:A:67:PRO:HG2	2:B:231:GLU:OE1	0.40	2.16	9	1
2:B:229:GLN:HA	2:B:232:SER:O	0.40	2.16	9	1
1:A:91:ASP:OD2	1:A:97:LYS:HB3	0.40	2.17	10	1
1:A:38:LYS:HE2	2:B:229:GLN:NE2	0.40	2.31	3	1
1:A:25:PHE:O	1:A:26:GLU:O	0.40	2.39	7	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	90/96 (94%)	76±1 (84±1%)	12±2 (13±2%)	3±1 (3±1%)	7	41
2	B	24/73 (33%)	22±1 (90±3%)	2±1 (8±3%)	0±0 (2±2%)	13	56
All	All	1140/1690 (67%)	975 (86%)	135 (12%)	30 (3%)	8	44

All 14 unique Ramachandran outliers are listed below. They are sorted by the frequency of

occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	26	GLU	5
1	A	31	ILE	4
1	A	94	GLY	3
1	A	17	GLY	3
2	B	245	THR	3
1	A	27	LEU	2
1	A	28	ASP	2
1	A	67	PRO	2
1	A	96	PHE	1
1	A	76	GLY	1
1	A	99	ALA	1
1	A	86	THR	1
2	B	232	SER	1
1	A	95	GLU	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	80/86 (93%)	72±2 (90±2%)	8±2 (10±2%)	11	56
2	B	20/57 (35%)	17±1 (86±5%)	3±1 (14±5%)	6	45
All	All	1000/1430 (70%)	889 (89%)	111 (11%)	9	53

All 32 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	23	ILE	10
1	A	46	LEU	10
1	A	53	PHE	10
2	B	229	GLN	9
2	B	227	LEU	8
1	A	62	ILE	7
1	A	24	THR	5
1	A	31	ILE	5
2	B	223	LEU	4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	102	MET	4
1	A	68	VAL	3
1	A	89	LEU	3
1	A	34	VAL	2
1	A	45	GLU	2
2	B	248	LYS	2
1	A	26	GLU	2
1	A	35	LEU	2
1	A	77	ILE	2
1	A	104	CYS	2
1	A	25	PHE	2
2	B	230	LEU	2
1	A	65	LEU	2
2	B	225	ASP	2
1	A	43	THR	2
1	A	87	TYR	2
1	A	59	ASP	1
1	A	78	ASP	1
2	B	224	LEU	1
1	A	27	LEU	1
1	A	82	TRP	1
1	A	75	LEU	1
2	B	253	GLU	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 49% for the well-defined parts and 46% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chemical_shifts_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	734
Number of shifts mapped to atoms	734
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	74	-0.08 ± 0.20	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	59	0.11 ± 0.24	None needed (< 0.5 ppm)
$^{13}\text{C}'$	0	—	None (insufficient data)
^{15}N	91	-0.59 ± 0.33	None needed (imprecise)

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 44%, i.e. 593 atoms were assigned a chemical shift out of a possible 1359. 15 out of 26 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	311/566 (55%)	154/226 (68%)	70/228 (31%)	87/112 (78%)
Sidechain	281/688 (41%)	154/396 (39%)	127/273 (47%)	0/19 (0%)

Continued on next page...

Continued from previous page...

	Total	^1H	^{13}C	^{15}N
Aromatic	1/105 (1%)	0/55 (0%)	0/47 (0%)	1/3 (33%)
Overall	593/1359 (44%)	308/677 (45%)	197/548 (36%)	88/134 (66%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 31%, i.e. 621 atoms were assigned a chemical shift out of a possible 2019. 15 out of 26 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	326/831 (39%)	161/331 (49%)	74/338 (22%)	91/162 (56%)
Sidechain	294/1075 (27%)	161/632 (25%)	133/391 (34%)	0/52 (0%)
Aromatic	1/113 (1%)	0/59 (0%)	0/51 (0%)	1/3 (33%)
Overall	621/2019 (31%)	322/1022 (32%)	207/780 (27%)	92/217 (42%)


7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:



7.2 Chemical shift list 2

File name: working_cs.cif

Chemical shift list name: *assigned_chemical_shifts_2*

7.2.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	337
Number of shifts mapped to atoms	337
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.2.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	47	-0.07 ± 0.38	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	29	0.11 ± 0.18	None needed (< 0.5 ppm)
$^{13}\text{C}'$	0	—	None (insufficient data)
^{15}N	64	-0.77 ± 0.25	Should be applied

7.2.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 5%, i.e. 73 atoms were assigned a chemical shift out of a possible 1359. 0 out of 26 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	60/566 (11%)	29/226 (13%)	9/228 (4%)	22/112 (20%)
Sidechain	13/688 (2%)	5/396 (1%)	8/273 (3%)	0/19 (0%)
Aromatic	0/105 (0%)	0/55 (0%)	0/47 (0%)	0/3 (0%)
Overall	73/1359 (5%)	34/677 (5%)	17/548 (3%)	22/134 (16%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 15%, i.e. 307 atoms were assigned a chemical shift out of a possible 2019. 0 out of 26 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	201/831 (24%)	90/331 (27%)	47/338 (14%)	64/162 (40%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Sidechain	106/1075 (10%)	46/632 (7%)	60/391 (15%)	0/52 (0%)
Aromatic	0/113 (0%)	0/59 (0%)	0/51 (0%)	0/3 (0%)
Overall	307/2019 (15%)	136/1022 (13%)	107/780 (14%)	64/217 (29%)

7.2.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.2.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain B:

