wwPDB X-ray Structure Validation Summary Report (i)

Oct 3, 2023 – 12:08 AM EDT

PDB ID	:	6PMP
Title	:	Crystal structure of a fragment of rat phospholipase Cepsilon EF3-RA1
Authors	:	Rugema, N.Y.; Lyon, A.M.
Deposited on	:	2019-07-02
Resolution	:	2.73 Å(reported)

1   Overall quality at a glance   ⓘ

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.73 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 19693 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 1-phosphatidylinositol 4,5-bisphosphate phosphodiesterase epsilon-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	609	Total	C	N	O	S	0	0	0
			4897	3121	844	901	31			
1	B	611	Total	C	N	O	S	0	0	0
			4902	3127	844	900	31			
1	C	617	Total	C	N	O	S	0	0	0
			4970	3176	852	910	32			
1	D	605	Total	C	N	O	S	0	0	0
			4853	3097	835	890	31			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1281	SER	-	expression tag	UNP Q99P84
A	1282	ASN	-	expression tag	UNP Q99P84
A	1283	ALA	-	expression tag	UNP Q99P84
B	1281	SER	-	expression tag	UNP Q99P84
B	1282	ASN	-	expression tag	UNP Q99P84
B	1283	ALA	-	expression tag	UNP Q99P84
C	1281	SER	-	expression tag	UNP Q99P84
C	1282	ASN	-	expression tag	UNP Q99P84
C	1283	ALA	-	expression tag	UNP Q99P84
D	1281	SER	-	expression tag	UNP Q99P84
D	1282	ASN	-	expression tag	UNP Q99P84
D	1283	ALA	-	expression tag	UNP Q99P84

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total 1	Ca 1	0	0
2	C	1	Total 1	Ca 1	0	0
2	D	1	Total 1	Ca 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	21	Total 21	O 21	0	0
3	B	17	Total 17	O 17	0	0
3	C	19	Total 19	O 19	0	0
3	D	10	Total 10	O 10	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.

3   Data and refinement statistics   (i)

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	93.57Å 127.75Å 139.34Å 90.00° 101.12° 90.00°	Depositor
Resolution (Å)	20.00 – 2.73	Depositor
% Data completeness (in resolution range)	99.3 (20.00-2.73)	Depositor
R_{merge}	0.27	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.55 (at 2.75Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.234 , 0.273	Depositor
Wilson B-factor (Å^2)	36.2	Xtriage
Anisotropy	1.481	Xtriage
L-test for twinning[2]	$< |L| > = 0.47, < L^2 > = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	19693	wwPDB-VP
Average B, all atoms (Å^2)	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.80% of the height of the origin peak. No significant pseudotranslation is detected.*

---

[1]Intensities estimated from amplitudes.
[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality ⓘ

4.1 Standard geometry ⓘ

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts ⓘ

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles ⓘ

4.3.1 Protein backbone ⓘ

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains ⓘ

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA ⓘ

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

4.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

4.7　Other polymers ⓘ

There are no such residues in this entry.

4.8　Polymer linkage issues ⓘ

There are no chain breaks in this entry.

5 Fit of model and data ⓘ

5.1 Protein, DNA and RNA chains ⓘ

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates ⓘ

EDS failed to run properly - this section is therefore empty.

5.4 Ligands ⓘ

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers ⓘ

EDS failed to run properly - this section is therefore empty.