



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 13, 2023 – 03:43 PM EDT

PDB ID : 4PNV  
Title : E. coli sliding clamp apo-crystal in P21 space group with larger cell dimensions  
Authors : Yin, Z.; Oakley, A.J.  
Deposited on : 2014-02-21  
Resolution : 1.86 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : **FAILED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS failed to run properly.

Mol	Chain	Length	Quality of chain
1	A	366	
1	B	366	

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6743 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase III subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	366	2900	1834	496	550	20	0	13	0
1	B	366	2894	1828	493	553	20	0	17	0

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	5	Total 5	Ca 5	0	0
2	B	3	Total 3	Ca 3	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total 1	Cl 1	0	0
3	B	3	Total 3	Cl 3	0	0

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 7 4 3	0	0
4	B	1	Total C O 7 4 3	0	0
4	B	1	Total C O 7 4 3	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	445	Total O 445 445	0	0
5	B	471	Total O 471 471	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS failed to run properly.


- Molecule 1: DNA polymerase III subunit beta

Chain A:  94% 5%



- Molecule 1: DNA polymerase III subunit beta

Chain B:  93% 7%



## 4 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	80.14Å 70.26Å 84.47Å 90.00° 114.67° 90.00°	Depositor
Resolution (Å)	27.55 – 1.86	Depositor
% Data completeness (in resolution range)	90.2 (27.55-1.86)	Depositor
$R_{merge}$	0.03	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.87 (at 1.87Å)	Xtrriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.183 , 0.234	Depositor
Wilson B-factor (Å <sup>2</sup> )	18.5	Xtrriage
Anisotropy	0.084	Xtrriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	6743	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.07% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, CL, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.36	0/2989	0.60	1/4048 (0.0%)
1	B	0.38	0/2995	0.63	1/4063 (0.0%)
All	All	0.37	0/5984	0.62	2/8111 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	197	ARG	NE-CZ-NH2	-5.03	117.79	120.30
1	B	306	LEU	CA-CB-CG	5.03	126.86	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2900	0	2937	20	0
1	B	2894	0	2914	18	0
2	A	5	0	0	0	0
2	B	3	0	0	0	0
3	A	1	0	0	0	0
3	B	3	0	0	2	0
4	A	7	0	10	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	14	0	20	0	0
5	A	445	0	0	2	0
5	B	471	0	0	3	0
All	All	6743	0	5881	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:260[B]:CYS:SG	1:B:336[B]:VAL:HG13	1.44	1.54
1:B:260[B]:CYS:SG	1:B:336[B]:VAL:CG1	2.35	1.13
1:A:35[B]:LEU:CD2	1:A:44[B]:LEU:HD23	2.05	0.85
1:A:35[B]:LEU:HD22	1:A:44[B]:LEU:CD2	2.08	0.83
1:A:35[B]:LEU:CD2	1:A:44[B]:LEU:CD2	2.57	0.82
1:A:147:ALA:O	1:A:197:ARG:NH2	2.17	0.77
1:A:35[B]:LEU:HD22	1:A:44[B]:LEU:HD23	1.65	0.75
1:A:139:ILE:HB	1:A:204:MET:HE1	1.76	0.66
1:A:35[B]:LEU:HD23	1:A:44[B]:LEU:HD23	1.80	0.62
1:A:139:ILE:HB	1:A:204:MET:CE	2.31	0.60
1:A:73:ARG:NH1	5:A:920:HOH:O	2.38	0.55
1:A:33[B]:LEU:HD11	1:A:44[B]:LEU:HB3	1.91	0.52
1:A:132:GLN:HE22	1:A:208:ASP:H	1.57	0.52
1:B:285[B]:VAL:HG22	1:B:310:TYR:CD2	2.45	0.52
1:A:33[B]:LEU:CD2	1:A:35[B]:LEU:HD21	2.39	0.52
1:A:33[B]:LEU:HD21	1:A:35[B]:LEU:HD21	1.94	0.49
1:B:33[B]:LEU:HG	1:B:72:ALA:CB	2.43	0.49
3:B:408:CL:CL	5:B:958:HOH:O	2.56	0.49
1:A:339[A]:MET:HE1	1:A:348:GLN:HG2	1.96	0.47
1:B:35[A]:LEU:HD22	1:B:44:LEU:CD2	2.44	0.46
1:B:184:ILE:HD11	1:B:188:LEU:HD11	1.98	0.45
1:A:170:VAL:HG22	1:A:179:VAL:HG23	1.97	0.45
1:A:48:ASP:O	1:A:49:LEU:HB2	2.16	0.45
1:B:33[A]:LEU:CD2	1:B:35[A]:LEU:HD21	2.47	0.45
1:A:139:ILE:HD12	1:A:204:MET:HE2	1.99	0.44
1:B:306:LEU:HD12	1:B:306:LEU:N	2.33	0.44
1:B:51:MET:HE1	1:B:198:LYS:C	2.38	0.43
1:B:287:GLU:O	3:B:407:CL:CL	2.73	0.43
1:A:251:ASN:HB2	5:A:653:HOH:O	2.17	0.43
1:B:348:GLN:NE2	5:B:684:HOH:O	2.46	0.43

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:51:MET:CE	1:B:198:LYS:HG3	2.48	0.42
1:A:35[B]:LEU:CD2	1:A:44[B]:LEU:HD22	2.43	0.42
1:A:73:ARG:HG2	5:B:670:HOH:O	2.19	0.42
1:B:51:MET:HE2	1:B:232:PHE:HZ	1.85	0.41
1:B:33[A]:LEU:HD23	1:B:35[A]:LEU:HD21	2.01	0.41
1:B:296:ASN:HB2	1:B:297:PRO:CD	2.51	0.41
1:B:51:MET:HE3	1:B:198:LYS:HG3	2.03	0.41
1:B:214:LEU:HD11	1:B:225:ALA:HB1	2.02	0.41
1:B:143:GLN:HG3	1:B:146:MET:CE	2.52	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	377/366 (103%)	372 (99%)	5 (1%)	0	100	100
1	B	381/366 (104%)	377 (99%)	4 (1%)	0	100	100
All	All	758/732 (104%)	749 (99%)	9 (1%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	322/313 (103%)	318 (99%)	4 (1%)	71	62
1	B	323/313 (103%)	316 (98%)	7 (2%)	52	36
All	All	645/626 (103%)	634 (98%)	11 (2%)	62	47

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	27	LEU
1	A	176	ARG
1	A	215	ARG
1	A	238	ASP
1	B	10	LEU
1	B	57	VAL
1	B	176	ARG
1	B	226[A]	HIS
1	B	226[B]	HIS
1	B	306	LEU
1	B	365	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	9	HIS
1	A	132	GLN
1	A	143	GLN
1	A	191	HIS
1	A	289	GLN
1	A	299	GLN
1	B	36	GLN
1	B	212	ASN
1	B	217	GLN
1	B	348	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 12 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	PEG	B	402	-	6,6,6	0.56	0	5,5,5	0.49	0
4	PEG	B	403	-	6,6,6	0.32	0	5,5,5	0.74	0
4	PEG	A	404	-	6,6,6	0.51	0	5,5,5	0.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEG	B	402	-	-	1/4/4/4	-
4	PEG	B	403	-	-	1/4/4/4	-
4	PEG	A	404	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	404	PEG	O1-C1-C2-O2
4	B	403	PEG	O1-C1-C2-O2
4	A	404	PEG	C4-C3-O2-C2
4	B	402	PEG	C1-C2-O2-C3

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS failed to run properly - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS failed to run properly - this section is therefore empty.

### 6.3 Carbohydrates

EDS failed to run properly - this section is therefore empty.

### 6.4 Ligands

EDS failed to run properly - this section is therefore empty.

### 6.5 Other polymers

EDS failed to run properly - this section is therefore empty.