



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 4, 2023 – 10:05 AM EDT

PDB ID : 6PNG
Title : Structure of human neuronal nitric oxide synthase R354A/G357D mutant heme domain in complex with 7-(3-(Aminomethyl)-4-propoxyphenyl)-4-methylquinolin-2-amine
Authors : Li, H.; Poulos, T.L.
Deposited on : 2019-07-02
Resolution : 1.77 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.35.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	9185 (1.80-1.76)
Clashscore	141614	10184 (1.80-1.76)
Ramachandran outliers	138981	10051 (1.80-1.76)
Sidechain outliers	138945	10050 (1.80-1.76)
RSRZ outliers	127900	9032 (1.80-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	421	
1	B	421	

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 7517 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitric oxide synthase, brain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	412	3371	2160	573	616	22	0	4	0
1	B	409	3343	2142	570	610	21	0	2	0

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	354	ALA	ARG	engineered mutation	UNP P29475
A	357	ASP	GLY	engineered mutation	UNP P29475
B	354	ALA	ARG	engineered mutation	UNP P29475
B	357	ASP	GLY	engineered mutation	UNP P29475

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is 7-[3-(aminomethyl)-4-propoxyphenyl]-4-methylquinolin-2-amine (three-letter code: OSG) (formula: C₂₀H₂₃N₃O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			24	20	3	1		
3	A	1	Total	C	N	O	0	0
			24	20	3	1		
3	B	1	Total	C	N	O	0	0
			24	20	3	1		
3	B	1	Total	C	N	O	0	0
			24	20	3	1		

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Zn	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	351	Total 351	O 351	0	0
5	B	269	Total 269	O 269	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitric oxide synthase, brain



- Molecule 1: Nitric oxide synthase, brain



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	52.17Å 125.15Å 164.92Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.16 – 1.77 39.16 – 1.78	Depositor EDS
% Data completeness (in resolution range)	99.1 (39.16-1.77) 99.5 (39.16-1.78)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	0.17	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.95 (at 1.77Å)	Xtrriage
Refinement program	PHENIX (1.11.1-2575_1496: ???)	Depositor
R, R_{free}	0.185 , 0.222 0.185 , 0.222	Depositor DCC
R_{free} test set	5144 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	28.0	Xtrriage
Anisotropy	0.976	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 42.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7517	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, HEM, OSG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/3479	0.50	0/4719
1	B	0.33	0/3444	0.49	0/4671
All	All	0.34	0/6923	0.50	0/9390

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3371	0	3288	19	0
1	B	3343	0	3257	11	0
2	A	43	0	30	2	0
2	B	43	0	30	2	0
3	A	48	0	0	1	0
3	B	48	0	0	1	0
4	B	1	0	0	0	0
5	A	351	0	0	3	0
5	B	269	0	0	1	0
All	All	7517	0	6605	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 2.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:528:LEU:HD22	1:A:536:PRO:HB2	1.63	0.80
1:A:519:ARG:NH1	5:A:904:HOH:O	2.29	0.65
2:B:803:HEM:HHC	2:B:803:HEM:HBB2	1.78	0.65
2:A:801:HEM:O1D	5:A:901:HOH:O	2.15	0.64
1:A:483:GLN:HB2	1:A:486:ARG:HG3	1.83	0.60
1:B:483:GLN:HB2	1:B:486:ARG:HG3	1.85	0.58
2:B:803:HEM:HBC2	2:B:803:HEM:HMC2	1.85	0.58
1:A:610:ASN:HB2	1:A:613:GLU:HG3	1.87	0.57
1:B:606:ASN:HD21	1:B:609:TYR:HB2	1.72	0.54
1:B:513:GLN:NE2	5:B:901:HOH:O	2.25	0.52
1:B:610:ASN:HB2	1:B:613:GLU:HG3	1.93	0.50
1:A:414:TRP:CE3	1:A:426:TRP:HA	2.48	0.49
1:A:625:LYS:HE3	1:A:625:LYS:HB2	1.57	0.47
1:A:572:VAL:HG21	3:A:802:OSG:C07	2.44	0.47
2:A:801:HEM:HBB2	2:A:801:HEM:HHC	1.96	0.47
1:A:341:MET:HG3	1:A:342:HIS:CD2	2.50	0.47
1:A:480:TRP:HB2	1:A:528:LEU:HB3	1.96	0.46
1:B:572:VAL:HG21	3:B:804:OSG:C07	2.45	0.46
1:A:530:GLN:HE21	1:A:533:GLY:HA2	1.82	0.45
1:A:667:MET:HE2	1:A:667:MET:HB3	1.53	0.45
1:A:668:GLU:O	1:A:672:ARG:HD3	2.18	0.44
1:A:408:TYR:CE1	1:A:412:HIS:CE1	3.06	0.44
1:B:689:SER:HB3	1:B:692:ILE:HD11	1.99	0.43
1:A:668:GLU:HB3	1:A:672:ARG:NH1	2.34	0.43
1:A:623:MET:HA	1:A:630:TRP:CD1	2.53	0.43
1:B:542:PRO:HA	1:B:543:PRO:HD3	1.94	0.42
1:A:555:LYS:HE3	1:A:555:LYS:HA	2.02	0.42
1:A:691:SER:HA	1:A:696:PHE:CG	2.55	0.41
1:B:327:LEU:HB2	1:B:704:ARG:HG2	2.02	0.41
1:B:691:SER:HA	1:B:696:PHE:CG	2.55	0.41
1:B:560:LYS:HE3	1:B:560:LYS:HB3	1.64	0.41
1:A:699:GLU:HB3	1:B:340:ILE:HD13	2.02	0.40
1:A:578:GLU:OE1	5:A:902:HOH:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	412/421 (98%)	408 (99%)	4 (1%)	0	100	100
1	B	407/421 (97%)	397 (98%)	9 (2%)	1 (0%)	47	32
All	All	819/842 (97%)	805 (98%)	13 (2%)	1 (0%)	51	35

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	495	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	372/376 (99%)	361 (97%)	11 (3%)	41	24
1	B	367/376 (98%)	359 (98%)	8 (2%)	52	36
All	All	739/752 (98%)	720 (97%)	19 (3%)	47	29

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	342	HIS
1	A	394	ASP
1	A	491	LYS
1	A	519	ARG
1	A	528	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	552	ARG
1	A	555	LYS
1	A	600	VAL
1	A	625	LYS
1	A	673[A]	CYS
1	A	673[B]	CYS
1	B	325	SER
1	B	326	THR
1	B	491	LYS
1	B	552	ARG
1	B	560	LYS
1	B	600	VAL
1	B	617	LYS
1	B	620	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	OSG	A	803	-	26,26,26	0.91	1 (3%)	35,36,36	1.29	4 (11%)
3	OSG	B	801	-	26,26,26	0.95	1 (3%)	35,36,36	1.46	5 (14%)
2	HEM	B	803	1	41,50,50	1.52	7 (17%)	45,82,82	1.57	8 (17%)
3	OSG	B	804	-	26,26,26	0.90	0	35,36,36	1.17	4 (11%)
2	HEM	A	801	1	41,50,50	1.48	6 (14%)	45,82,82	1.66	9 (20%)
3	OSG	A	802	-	26,26,26	0.95	0	35,36,36	1.13	3 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	OSG	A	803	-	-	0/10/19/19	0/3/3/3
3	OSG	B	801	-	-	0/10/19/19	0/3/3/3
2	HEM	B	803	1	-	2/12/54/54	-
3	OSG	B	804	-	-	1/10/19/19	0/3/3/3
2	HEM	A	801	1	-	0/12/54/54	-
3	OSG	A	802	-	-	3/10/19/19	0/3/3/3

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	803	HEM	C3C-C2C	-4.34	1.34	1.40
2	A	801	HEM	C3C-C2C	-3.93	1.34	1.40
2	B	803	HEM	C3C-CAC	3.30	1.54	1.47
2	A	801	HEM	C3C-CAC	3.14	1.54	1.47
2	B	803	HEM	CAB-C3B	2.94	1.55	1.47
3	B	801	OSG	C02-N01	2.74	1.36	1.33
2	A	801	HEM	CAB-C3B	2.74	1.54	1.47
2	B	803	HEM	CMD-C2D	2.31	1.55	1.50
3	A	803	OSG	C02-N01	2.30	1.36	1.33
2	A	801	HEM	CAA-C2A	2.20	1.55	1.52
2	A	801	HEM	FE-NB	2.13	2.07	1.96
2	B	803	HEM	CHA-C4D	2.06	1.40	1.35
2	B	803	HEM	CAA-C2A	2.04	1.55	1.52
2	A	801	HEM	CMC-C2C	2.02	1.56	1.51
2	B	803	HEM	FE-ND	2.00	2.06	1.96

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	801	OSG	O29-C24-C25	4.41	121.55	115.78
2	B	803	HEM	CMA-C3A-C4A	-4.35	121.78	128.46
2	A	801	HEM	CBA-CAA-C2A	-4.12	105.59	112.62
3	B	801	OSG	C04-C05-C10	4.04	120.20	118.01
2	A	801	HEM	C4B-CHC-C1C	4.00	127.84	122.56
3	A	803	OSG	O29-C24-C25	3.78	120.72	115.78
2	B	803	HEM	CBA-CAA-C2A	-3.64	106.40	112.62
3	A	803	OSG	C04-C05-C10	3.54	119.93	118.01
2	B	803	HEM	C4B-CHC-C1C	3.41	127.06	122.56
2	A	801	HEM	C3B-C2B-C1B	3.15	108.82	106.49
2	B	803	HEM	CMA-C3A-C2A	3.14	130.87	124.94
2	A	801	HEM	CMA-C3A-C4A	-2.96	123.92	128.46
3	A	802	OSG	C04-C05-C10	2.89	119.58	118.01
3	B	804	OSG	C04-C05-C10	2.88	119.57	118.01
3	A	802	OSG	C05-C10-N01	-2.80	119.84	122.81
3	B	804	OSG	C05-C10-N01	-2.75	119.90	122.81
3	B	804	OSG	O29-C24-C25	2.66	119.26	115.78
2	B	803	HEM	C3B-C2B-C1B	2.62	108.43	106.49
2	B	803	HEM	C3D-C4D-ND	-2.41	107.49	110.17
3	B	801	OSG	O29-C24-C23	-2.33	118.92	123.97
2	B	803	HEM	C4D-ND-C1D	2.31	107.46	105.07
2	B	803	HEM	CAD-CBD-CGD	-2.29	108.69	113.60
3	A	802	OSG	C03-C04-C05	2.27	120.02	117.78
2	A	801	HEM	C4D-ND-C1D	2.24	107.38	105.07
2	A	801	HEM	CMC-C2C-C3C	2.23	128.86	124.68
3	B	801	OSG	C05-C10-N01	-2.19	120.49	122.81
3	A	803	OSG	C26-C25-C24	2.19	120.56	118.26
2	A	801	HEM	C2C-C3C-C4C	2.15	108.40	106.90
3	B	804	OSG	C08-C09-C10	-2.09	119.61	121.44
3	B	801	OSG	C26-C25-C24	2.08	120.45	118.26
2	A	801	HEM	C4C-CHD-C1D	2.06	125.27	122.56
3	A	803	OSG	O29-C24-C23	-2.02	119.61	123.97
2	A	801	HEM	C1B-NB-C4B	2.01	107.14	105.07

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	802	OSG	C24-C25-C27-N28
2	B	803	HEM	C1A-C2A-CAA-CBA
2	B	803	HEM	C3A-C2A-CAA-CBA
3	A	802	OSG	O29-C30-C31-C32
3	A	802	OSG	C26-C25-C27-N28

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	B	804	OSG	C31-C30-O29-C24



There are no ring outliers.



4 monomers are involved in 6 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	803	HEM	2	0
3	B	804	OSG	1	0
2	A	801	HEM	2	0
3	A	802	OSG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	412/421 (97%)	0.39	33 (8%) 12 11	26, 37, 69, 124	0
1	B	409/421 (97%)	0.64	52 (12%) 3 3	27, 42, 80, 112	0
All	All	821/842 (97%)	0.51	85 (10%) 6 6	26, 39, 75, 124	0

All (85) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	609	TYR	9.0
1	B	354	ALA	8.4
1	A	608	ARG	8.2
1	A	607	SER	8.2
1	B	608	ARG	7.4
1	A	604	CYS	6.2
1	A	606	ASN	6.1
1	B	609	TYR	5.8
1	B	495	GLY	5.7
1	B	493	PRO	5.5
1	B	721	TRP	5.5
1	A	353	VAL	4.9
1	B	720	VAL	4.8
1	A	605	ASP	4.3
1	B	304	ARG	4.2
1	A	305	PHE	4.1
1	B	607	SER	3.9
1	A	493	PRO	3.6
1	A	696	PHE	3.5
1	A	685	VAL	3.4
1	B	683	TRP	3.4
1	B	360	PHE	3.4
1	B	722	LYS	3.4
1	B	375	LYS	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	491	LYS	3.3
1	A	327	LEU	3.3
1	B	326	THR	3.2
1	B	355	THR	3.2
1	B	374	ILE	3.2
1	B	604	CYS	3.1
1	B	624	ARG	3.1
1	B	357	ASP	3.1
1	B	606	ASN	3.0
1	A	360	PHE	3.0
1	B	393	ILE	3.0
1	B	571	ALA	2.9
1	B	494	ASP	2.9
1	B	496	SER	2.9
1	B	386	LEU	2.8
1	B	474	LYS	2.8
1	B	598	ILE	2.8
1	A	572	VAL	2.8
1	B	327	LEU	2.8
1	A	603	TYR	2.7
1	B	605	ASP	2.7
1	B	356	LYS	2.7
1	B	420	CYS	2.7
1	B	685	VAL	2.7
1	B	358	GLN	2.7
1	A	624	ARG	2.7
1	B	376	ARG	2.6
1	A	635	LEU	2.6
1	B	719	HIS	2.6
1	A	304	ARG	2.6
1	B	572	VAL	2.6
1	A	303	PRO	2.5
1	B	690	GLY	2.5
1	A	357	ASP	2.5
1	A	690	GLY	2.5
1	A	682	VAL	2.5
1	B	718	THR	2.4
1	B	589	PHE	2.4
1	A	355	THR	2.4
1	A	686	PRO	2.4
1	B	684	ILE	2.2
1	A	354	ALA	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	692	ILE	2.2
1	A	596	THR	2.2
1	B	625	LYS	2.2
1	A	687	PRO	2.2
1	B	688	MET	2.2
1	B	394	ASP	2.2
1	A	334	TYR	2.1
1	B	717	ASN	2.1
1	B	504	VAL	2.1
1	B	512	GLN	2.1
1	B	689	SER	2.1
1	B	711	TYR	2.1
1	B	359	LEU	2.1
1	B	686	PRO	2.1
1	A	394	ASP	2.1
1	A	352	ASP	2.0
1	A	587	CYS	2.0
1	A	495	GLY	2.0
1	B	577	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	OSG	B	801	24/24	0.86	0.15	26,47,67,68	0
3	OSG	A	803	24/24	0.90	0.16	29,48,64,69	0
3	OSG	A	802	24/24	0.94	0.14	22,37,50,55	0
3	OSG	B	804	24/24	0.95	0.14	26,35,52,61	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	HEM	A	801	43/43	0.97	0.14	22,28,42,49	0
2	HEM	B	803	43/43	0.98	0.14	23,30,42,46	0
4	ZN	B	802	1/1	1.00	0.06	31,31,31,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around OSG A 803:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around OSG A 802:**



$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM A 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.