

wwPDB EM Validation Summary Report (i)

Apr 17, 2024 - 08:17 pm BST

PDB ID	:	7PNV
EMDB ID	:	EMD-13553
Title	:	Assembly intermediate of mouse mitochondrial ribosome small subunit without
		mS37 in complex with RbfA and Mettl15 $$
Authors	:	Itoh, Y.; Khawaja, A.; Laptev, I.; Sergiev, P.; Rorbach, J.; Amunts, A.
Deposited on	:	2021-09-08
Resolution	:	3.06 Å(reported)
Based on initial model	:	6RW4

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/EMValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis	:	0.0.1.dev92
Mogul	:	1.8.4, CSD as541be (2020)
MolProbity	:	4.02b-467
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 3.06 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f EM} {f structures} \ (\#{f Entries})$
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826
RNA backbone	4643	859

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for $\geq=3, 2, 1$ and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq=5\%$ The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	956	82%	16% ••
2	В	291	73% •	23%
3	С	167	74% 5%	21%
4	D	432	75% 5%	21%
5	Е	125	91%	• 5%
6	F	242	83%	• 14%
7	G	390	79% 5	% 17%



Mol	Chain	Length	Quality of chain	
8	Η	160	79%	8% 12%
9	Ι	191	6 9% •••	28%
10	J	139	75%	• 22%
11	К	128	73%	6% 21%
12	L	258	5% 66% ·	32%
13	М	135	92%	• 7%
14	Ν	120	91%	• 6%
15	О	254	- 78%	22%
16	Р	143	• 66% •	32%
17	Q	86	97%	•
18	R	359	• 80%	• 18%
19	S	177	75%	• 23%
20	Т	171	• 95%	
21	U	200	87%	13%
22	V	415	85%	• 12%
23	W	186	•	46%
24	X	391	⊷ 86%	5% 9%
25	Y	384	7%	
26	Z	106	7%	• 5%
27	0	218	97%	
28	1	320	₹ 	. 13%
29	3	200	36% 6404	- 15/0
30	4	685	17%	1 / 0/
21		350	8%	• 14%
91 90	a h	000 406	55% 57%	43%
54	U	400	80%	20%

Continued from previous page...



2 Entry composition (i)

There are 39 unique types of molecules in this entry. The entry contains 131939 atoms, of which 60861 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a RNA chain called 12S mitochondrial rRNA.

Mol	Chain	Residues				AltConf	Trace			
1	А	951	Total 30430	C 9078	H 10223	N 3629	O 6549	Р 951	0	0

• Molecule 2 is a protein called 28S ribosomal protein S2, mitochondrial.

Mol	Chain	Residues			AltConf	Trace				
2	В	223	Total	C	H 1700	N	0	S	0	0
			3590	1142	1799	326	315	8		

• Molecule 3 is a protein called 28S ribosomal protein S24, mitochondrial.

Mol	Chain	Residues	Atoms						AltConf	Trace
3	С	132	Total 2163	C 690	Н 1091	N 197	0 180	${S \atop 5}$	0	0

• Molecule 4 is a protein called 28S ribosomal protein S5, mitochondrial.

Mol	Chain	Residues			AltConf	Trace				
4	Л	3/3	Total	С	Η	Ν	0	\mathbf{S}	0	0
4	D	040	5527	1716	2794	527	480	10	0	0

• Molecule 5 is a protein called 28S ribosomal protein S6, mitochondrial.

Mol	Chain	Residues			AltConf	Trace				
5	Е	119	Total 1916	C 599	Н 968	N 171	0 175	${ m S} { m 3}$	0	0

• Molecule 6 is a protein called 28S ribosomal protein S7, mitochondrial.

Mol	Chain	Residues			AltConf	Trace				
6	F	208	Total 3472	C 1096	Н 1750	N 316	O 299	S 11	0	0



• Molecule 7 is a protein called 28S ribosomal protein S9, mitochondrial.

Mol	Chain	Residues			AltConf	Trace				
7	G	325	Total 5304	C 1689	Н 2630	N 480	0 491	S 14	0	0

• Molecule 8 is a protein called 28S ribosomal protein S10, mitochondrial.

Mol	Chain	Residues			Atom	.s			AltConf	Trace
8	Н	140	Total	C	H	N	0	S	0	0
			2346	742	1193	200	207	4		

• Molecule 9 is a protein called 28S ribosomal protein S11, mitochondrial.

Mol	Chain	Residues			Atom	S			AltConf	Trace
9	Ι	137	Total 2047	C 629	Н 1038	N 191	0 184	${f S}{5}$	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
I	181	5F0	ASN	conflict	UNP Q9DCA2

• Molecule 10 is a protein called 28S ribosomal protein S12, mitochondrial.

Mol	Chain	Residues			Aton	ns			AltConf	Trace
10	J	108	Total 1749	C 528	Н 903	N 172	0 141	${ m S}{ m 5}$	0	0

• Molecule 11 is a protein called 28S ribosomal protein S14, mitochondrial.

Mol	Chain	Residues			Aton	ıs			AltConf	Trace
11	K	101	Total 1743	C 534	H 888	N 175	0 140	S 6	0	0

• Molecule 12 is a protein called 28S ribosomal protein S15, mitochondrial.

Mol	Chain	Residues			Atom	IS			AltConf	Trace
12	L	176	Total 3041	C 930	H 1576	N 274	O 255	S 6	0	0

• Molecule 13 is a protein called 28S ribosomal protein S16, mitochondrial.



Mol	Chain	Residues			Atom	S			AltConf	Trace
13	М	126	Total 2004	C 623	Н 1009	N 194	O 172	${ m S}{ m 6}$	0	0

• Molecule 14 is a protein called 28S ribosomal protein S17, mitochondrial.

Mol	Chain	Residues			AltConf	Trace				
14	Ν	113	Total 1839	C 575	Н 951	N 160	O 150	${ m S} { m 3}$	0	0

• Molecule 15 is a protein called 28S ribosomal protein S18b, mitochondrial.

Mol	Chain	Residues			Atoms	S			AltConf	Trace
15	О	197	Total 3146	C 1014	Н 1548	N 289	O 286	S 9	0	0

• Molecule 16 is a protein called 28S ribosomal protein S18c, mitochondrial.

Mol	Chain	Residues	Atoms						AltConf	Trace
16	Р	97	Total 1610	C 505	Н 818	N 140	O 139	S 8	0	0

• Molecule 17 is a protein called 28S ribosomal protein S21, mitochondrial.

Mol	Chain	Residues			Aton	ns			AltConf	Trace
17	Q	86	Total 1482	C 453	Н 750	N 146	0 126	S 7	0	0

• Molecule 18 is a protein called 28S ribosomal protein S22, mitochondrial.

Mol	Chain	Residues			Atom	S			AltConf	Trace
18	R	294	Total 4816	C 1526	Н 2416	N 418	0 449	${ m S} 7$	0	0

• Molecule 19 is a protein called 28S ribosomal protein S23, mitochondrial.

Mol	Chain	Residues			Atom	IS			AltConf	Trace
19	S	136	Total 2257	C 722	Н 1133	N 199	O 201	${f S}{2}$	0	0

• Molecule 20 is a protein called 28S ribosomal protein S25, mitochondrial.



Mol	Chain	Residues			Aton	ıs			AltConf	Trace
20	Т	170	Total 2801	C 892	Н 1413	N 238	O 246	S 12	0	0

• Molecule 21 is a protein called 28S ribosomal protein S26, mitochondrial.

Mol	Chain	Residues			Atom	IS			AltConf	Trace
21	U	174	Total 2908	C 894	Н 1459	N 283	O 270	$\frac{S}{2}$	0	0

• Molecule 22 is a protein called 28S ribosomal protein S27, mitochondrial.

Mol	Chain	Residues			Atom	S			AltConf	Trace
22	V	365	Total 5970	C 1911	Н 2972	N 506	O 570	S 11	0	0

• Molecule 23 is a protein called 28S ribosomal protein S28, mitochondrial.

Mol	Chain	Residues			Aton	ıs			AltConf	Trace
23	W	100	Total 1606	C 503	Н 813	N 141	0 146	${ m S} { m 3}$	0	0

• Molecule 24 is a protein called 28S ribosomal protein S29, mitochondrial.

Mol	Chain	Residues			Atom	S			AltConf	Trace
24	Х	357	Total 5762	C 1834	Н 2881	N 515	O 522	S 10	0	0

• Molecule 25 is a protein called 28S ribosomal protein S31, mitochondrial.

Mol	Chain	Residues			Atom	IS			AltConf	Trace
25	Y	149	Total 2439	C 809	Н 1193	N 201	0 233	${ m S} { m 3}$	0	0

• Molecule 26 is a protein called 28S ribosomal protein S33, mitochondrial.

Mol	Chain	Residues			Aton	ns			AltConf	Trace
26	Z	101	Total 1682	C 526	H 848	N 157	0 148	${ m S} { m 3}$	0	0

• Molecule 27 is a protein called 28S ribosomal protein S34, mitochondrial.



Mol	Chain	Residues			Atoms	5			AltConf	Trace
27	0	216	Total 3649	C 1139	Н 1838	N 355	0 313	$\frac{S}{4}$	0	0

• Molecule 28 is a protein called 28S ribosomal protein S35, mitochondrial.

Mol	Chain	Residues			Atom	.s			AltConf	Trace
28	1	277	Total	С	Η	Ν	0	\mathbf{S}	0	0
	-	211	4440	1397	2228	380	423	12		

• Molecule 29 is a protein called Aurora kinase A-interacting protein.

Mol	Chain	Residues			Atom	S			AltConf	Trace
29	3	72	Total 1371	C 414	Н 728	N 135	O 93	S 1	0	0

• Molecule 30 is a protein called Pentatricopeptide repeat domain-containing protein 3, mitochondrial.

Mol	Chain	Residues			Atom	s			AltConf	Trace
30	4	589	Total 9558	C 3064	Н 4800	N 800	0 872	S 22	0	0

• Molecule 31 is a protein called Putative ribosome-binding factor A, mitochondrial.

Mol	Chain	Residues			Atom	S			AltConf	Trace
31	a	191	Total	С	Н	Ν	0	\mathbf{S}	0	0
01	a	101	3129	977	1583	280	284	5	Ŭ,	Ŭ

• Molecule 32 is a protein called 12S rRNA N4-methylcytidine methyltransferase.

Mol	Chain	Residues			Atom	S			AltConf	Trace
32	b	323	Total 5114	C 1592	Н 2595	N 454	0 462	S 11	0	0

• Molecule 33 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	AltConf
33	А	46	Total Mg 46 46	0
33	В	1	Total Mg 1 1	0



Continued from previous page...

Mol	Chain	Residues	Atoms	AltConf
33	Х	1	Total Mg 1 1	0
33	3	1	Total Mg 1 1	0

• Molecule 34 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	AltConf
34	А	13	Total K 13 13	0

• Molecule 35 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	AltConf
35	О	1	Total Zn 1 1	0

• Molecule 36 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe₂S₂).



Mol	Chain	Residues	Atoms	AltConf
26	D	1	Total Fe S	0
- 50	Г	1	4 2 2	0
26	Т	1	Total Fe S	0
- 50			4 2 2	0

• Molecule 37 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula:



 $C_{10}H_{16}N_5O_{13}P_3).$



Mol	Chain	Residues	Atoms					AltConf	
27	v	1	Total	С	Η	Ν	Ο	Р	0
- 57	Λ	1	43	10	12	5	13	3	0

• Molecule 38 is S-ADENOSYL-L-HOMOCYSTEINE (three-letter code: SAH) (formula: $\rm C_{14}H_{20}N_6O_5S).$



Mol	Chain	Residues	Atoms				AltConf		
90	h	1	Total	С	Η	Ν	Ο	\mathbf{S}	0
30	D	1	46	14	20	6	5	1	0



• Molecule 39 is water.

Mol	Chain	Residues	Atoms	AltConf
39	А	549	Total O 549 549	0
39	В	23	Total O 23 23	0
39	С	19	Total O 19 19	0
39	D	34	Total O 34 34	0
39	Е	2	Total O 2 2	0
39	F	7	Total O 7 7	0
39	G	10	Total O 10 10	0
39	Н	19	Total O 19 19	0
39	Ι	5	Total O 5 5	0
39	J	14	Total O 14 14	0
39	К	25	Total O 25 25	0
39	L	6	Total O 6 6	0
39	М	16	Total O 16 16	0
39	Ν	8	Total O 8 8	0
39	О	25	Total O 25 25	0
39	Р	3	Total O 3 3	0
39	Q	8	Total O 8 8	0
39	R	13	Total O 13 13	0
39	S	4	Total O 4 4	0
39	Т	19	Total O 19 19	0
39	U	3	Total O 3 3	0



Continued from previous page...

Mol	Chain	Residues	Atoms	AltConf
39	W	2	Total O 2 2	0
39	Х	5	Total O 5 5	0
39	Y	7	Total O 7 7	0
39	Ζ	18	Total O 18 18	0
39	0	7	Total O 7 7	0
39	1	5	Total O 5 5	0
39	3	7	Total O 7 7	0
39	a	5	$\begin{array}{cc} \text{Total} & \text{O} \\ 5 & 5 \end{array}$	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 12S mitochondrial rRNA









Chain I:	69%	28%	
MET GLN CLN VAL LEU ARG ASN ASN SER CLY TRP LEU	LEU TRP TRP ALA ALA ALA ALA ALA ALA ARG CLY ALA ARG ALA ASN ASN ASN ASN ASN ASN ASN ASN ASN AS	ALA ALA ALA ALA ALA ALA ALA ALA ALA ALA	V148 T173 D174
N175 T176 5F0181 L191			
• Molecule 10: 2	28S ribosomal protein S12, mitochondrial	l	
Chain J:	75%	• 22%	
MET SER TRP PRO CLY CLY CLY CLY CLY CLY CLU TYR CLU TYR THR	SER SER SEU SEU SEU SEU ALA ALA ALA ALA ALA ALA ALA ALA ALA AL	2	
• Molecule 11: 2	28S ribosomal protein S14, mitochondrial	l	
Chain K:	73%	6% 21%	
MET ALA ALA ALA SER VAL LEU CLEU CLEU LEU LEU ARG THR	PHE ALA ALA ALA ALA ALA PRO PRO PRO CLY CLA CLA CLA CLA CLA CLA CLA CLA CLA CLA	112	
• Molecule 12: 2	28S ribosomal protein S15, mitochondrial	l	
Chain L:	66%	32%	
MET LEU ARG ALA ALA ARG ALA ALA LEU SER SER SER VAL	ARG GLN ALA ALA ALA ALA ALA ALA ALA ALA ALA A	PRO SER LEU LEU ALA ALA ALA ALA ALA ALA VAL	GLN LYS PRO VAL GLN
PR0 K62 Q63 D64 P193 P193 P193 P226 A227	K228 K229 A230 A231 A235 A235 A235 A235 A235 A235 A235 A235	CTAR VAL ASN	
• Molecule 13: 2	28S ribosomal protein S16, mitochondrial	l	
Chain M:	92%	• 7%	
MET MET VAL GLN LEU THR THR THR THR THR THR CKO K10 K10	1104 E126 S127 E131 E131 A134 A134 S135		
• Molecule 14: 2	28S ribosomal protein S17, mitochondrial	l	
Chain N:	91%	• 6%	1
MET SER 13 13 14 14 14 14 14 14 14 14 14 14 14 14 14	P113 R114 E115 E115 LTS LTS VAL		

• Molecule 15: 28S ribosomal protein S18b, mitochondrial



Chain O:	78%	22%
MET ALA ALA PRO PRO LEU HIS HIS LEU LEU	LEU VAL VAL LEU LEU LEU LEU ARG ALA CLN CAL CAL CAL CAL CAL CAL CAL CAL CAL CAL	843 V46 V46 PR0 PR0 ALA ALA ALA CLU CLU CLU CLU CLU
GLY SER GLN SER ALA		
• Molecule 16:	28S ribosomal protein S18c, mitochondrial	l
Chain P:	66% ·	32%
MET ALA ALA ALA LEU VAL ALA CYS SER SER GLY	ARG ARG ARG ASS ALA ALA ALA ALA ALA ALA ASS ASS ASS	
• Molecule 17:	28S ribosomal protein S21, mitochondrial	
Chain Q:	97%	
A2 139 R42 R50 R50 C87		
• Molecule 18:	28S ribosomal protein S22, mitochondrial	
Chain R:	80%	• 18%
MET ALA ALA ALA VAL ARG PRO PRO LEU SER LEU	ARG GLY GLY GLY SER GLY SER ARG ARG ARG ARG ARG ARG ARG ARG ARG AR	PHU PHU OLU VAL OLY GLY GLN SER SER SER SER SER SER SER SER SER SER
SER SER F63 V64 K65 K65 K65 S161	R169 V207 V207 D237 D237 D235 C 1366 C 1351 V355 S356 ALA ALA ALA ALA ALA ALA	
• Molecule 19:	28S ribosomal protein S23, mitochondrial	
Chain S:	75%	• 23%
MET A2 V9 1112 E114 E114	H136 H136 H136 H14 ARLA ARLA ARLA ARLA ARLA ARLA ARLA ARC CLU CLU CLU CLU CLU CLU ARP ARP CLU CLU CLU CLU CLU CLU CLU CLU CLU CLU	VAL LYS GLN GLU PRO PRO PRO PRO PRO
• Molecule 20:	28S ribosomal protein S25, mitochondrial	
Chain T:	95%	• •
MET P2 129 192 192 1160		

• Molecule 21: 28S ribosomal protein S26, mitochondrial



Chain U:	87%		13%
MET LEU LEU ARG ALA ARG ARG ALA ARG PRO PRO PRO	ARA ARA PRO PRO PRO LEU LEU LEU LEU VAL ARG ARG ARG ARG ARG ARG		
• Molecule 22: 285	5 ribosomal protein S27	mitochondrial	
Chain V:	85%		• 12%
MET ALA ALA ALA PRO PRO PRO CVS CVS CVS CVS CVS CVS CVS ALA	ARM ARM LEU LEU ARF ARG CYS CYS CYS CYS CYS CYS CYS CYS CYS CYS	V175 P176 S177 S177 L181 V187 Y187 Y187 F196	Y226 D272 V279 S282 P294 VML CUN THR SBR GLU GLN
PRO CLIM CLIM CLIU CLIV CLIV CLIV CLIV CLIV CLIV CLIV CLIV	q397 R399 E399 E400 K400 K400 K400 C403 C403 C403 C403 C405 C406 A407 C406 C406 C406 C406 C406 C406 C406 C406	ALA ALA GLU LYS ALA ALA ALA	
• Molecule 23: 28	S ribosomal protein S28	, mitochondrial	
Chain W:	52%	• 4	5%
MET ALA ALA ALA CYS CYS CYS ALA ALA ALA ALA GLY	ARG PHE LEU LEU ALA ALA PHE SFER SFER ALA ASN ALA ASN ALA	GLU GLY GLY GLY GLY GLU GLU ASP ASP SER SER SER SER	ARA ARA ALA ALA ALA CLY CLY ALA ALA ALA ALA CLU CLU CLU CLU CLU SER ARC
ASP LEU LEU ARG ARG ALA ALA CLU CLU CLU SER SER SER	PRO VT6 VT6 L153 L153 D158 ASP SER SER SER SER	GLU GLN PRO SER LYS	
• Molecule 24: 285	5 ribosomal protein S29	, mitochondrial	
Chain X:	86%		5% 9%
MET LEU LIEU THR CLY CLY LEU PHE SER SER ARG CVAL CLA	LLIS LLIS ASP ASP ASP ARC CYS ARC ARC ARC ARC ALA ALA ALA ALA ALA SER SER SER SER SER SER SER SER ASP	GLN VAL PRO P35 A35 B35 P38 P38 P38 P38 P38 P38 P38 P38 P38 P38	H112 H116 E122 E122 L144 L144 L146 B126
K205 E210 K226 K226 K227 L237 L237	L266 L267 K268 L274 L274 L376 L309 L330 L330 L330 L330 L333 L331 L333 L333		
• Molecule 25: 28	S ribosomal protein S31	, mitochondrial	
Chain Y:	38% •	61%	
MET LEU HIS ARG TLE PHE LEU ARG PRO PRO	FIRE GLY GLY FRO FRO CYS SER ASP ASP ASP ASP VAL ASP VAL VAL LEU LEU	PRO ALA ALA ALA GLN SER GLY ALA ARG GLU ASN	GLLE ARG ARG CYS CYS ARG ARG SER ILE LYS CYS SER LYS VAL
ASP GLN SER VAL VAL ALA ALA ALA ALA CLU CLU SER CLU ALA	ALA GLU SER CLN CLN CLN CLN CLN CLN CLN CLN CLN CLN	ILE LYS ASP MET LYS VAL SEU SEU SEU ALA ALA ALA	THI THR PRO LLYS PRO ARG ARG CLY ARG ALA ALA ALA ALA ALA ALA
THR VAL ASP ARG LEU CLEU CLEU CLY ARD PRO PRO PRO	LTS ARG ARG ARG ARG ALU CEU CEU CEU CEU ALA ALA ALA ALA ALA ALA ALA ALA ALA	ALA ASP SER SER LEU PHE ASP CLYS GLN THR THR LYS SER	LEU LEU ARG GLN GLN GLU GLU CLU CLU CLU CLU CLU CLU CLU CLU CLU C



LYS ASP ARG GLU LYS ARG TLYS SER PHE THR HIS HIS TLE	ILLE SER MET MET LYS ILYS ALA ALA LYS SER PRO SER SER SER	ARG ALA SER THR ARG ARG CLN GLN GLN GLN GLN	PHE ASP GLU GLU ASP ASP SER SER CLU CLU GLU	PRU ASP PHE LVS K237 Y238 F240 F240
K241 ♦ C242 ♦ K243 ♦ K244 ♦ 1.245 ♦ F248 ♦ A249 ♦ K251 ♦	P252 P253 A254 D255 P255 P255 P255 P258 P258	E261 A262 8263 P264 B268 C304 I321	◆ ◆ 13380 13384	
• Molecule 26: 28	S ribosomal pro	tein S33, mitoc	hondrial	
Chain Z:		94%		• 5%
MET S2 P3 A23 A23 K24 K24 K24 K22 K23 C920	A100 A102 LYS LYS LYS LYS			
• Molecule 27: 28	S ribosomal pro	tein S34, mitoc	hondrial	
Chain 0:		97%		••
MET ALA ALA R3 K5 K5 K7 B37 L41 L41 R48 R48	R99 P157 E158 D159 R160 R160 P215	<mark>↓218</mark>		
• Molecule 28: 28	S ribosomal pro	tein S35, mitoc	hondrial	
Chain 1:		85%	•	13%
MET ALA ALA ALA ALA ALA ALA CEU ALA CEU SER CEU SER CTS CTS	PRO GLY ARG VAL VAL LEU LEU LEU LEU THR PHE SER SER VAL	ALA SER PRO ALA ALA ALA ARG ARG PLY PRO PRO	THR ALA SER SER ARC CUU CUU ARC P44 M45 M45 M45 P33	1103 11103 1111 1160 1160 1244
D246 271 277 277 4274 4320 4320				
• Molecule 29: Au	ırora kinase A-iı	nteracting prote	ein	
Chain 3:	36%		64%	
MET PHE LEU ALA ARG LEU THR SER ARG ALA ALA ALA THR	VAL VAL PRO TRP ALA GLY PHE SER SER SER CYS	GLY SER GLY VAL TLE GLY SER ALA ALA ARG	PRO LEU TYR SER LEU GLN PRO PRO SER ARG ARG	ALA SER LEU PRO GLY CYS ARG THR GLN SER GLU
LEU GLU GLU GLU PHE LEU VAL PRO ARG LYS MET ALA ALA SER SER	PRO LEU GLU GLU SER TRP TRP LEU GLN LEU	PRO ARG ARG ASN VAL CUU PRO VAL THR THR	ALA PRO SER CLN PHE CLN CVS CVS CVS PRO PRO PRO PRO CLN CLN CLN CLN	GLU GLU GLU GLN GLN GLN CLN ARG GLU ALA
TRP ALA ALA ALA ALA PRO VAL CTS CTS CTS CTS K156 K156	N199 K200			
• Molecule 30: Pe	entatricopeptide	repeat domain-	-containing protein	3, mitochondrial
Chain 4:	_	84%	·	14%











4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	59111	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{\AA}^2)$	31	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	3600	Depositor
Magnification	165000	Depositor
Image detector	GATAN K2 SUMMIT $(4k \ge 4k)$	Depositor
Maximum map value	1.795	Depositor
Minimum map value	-0.950	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.032	Depositor
Recommended contour level	0.1	Depositor
Map size (Å)	398.40192, 398.40192, 398.40192	wwPDB
Map dimensions	576, 576, 576	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.69167, 0.69167, 0.69167	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: B8T, MG, FES, SAH, ZN, AYA, ATP, 5MC, 5F0, MA6, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bo	nd lengths	Bond angles		
	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.26	1/22518~(0.0%)	0.67	0/35045	
2	В	0.25	0/1832	0.41	0/2479	
3	С	0.25	0/1100	0.43	0/1485	
4	D	0.25	0/2789	0.43	0/3739	
5	Ε	0.25	0/966	0.42	0/1308	
6	F	0.24	0/1763	0.37	0/2368	
7	G	0.25	0/2734	0.39	0/3669	
8	Н	0.25	0/1179	0.41	0/1597	
9	Ι	0.25	0/1018	0.45	0/1374	
10	J	0.26	0/862	0.45	0/1155	
11	Κ	0.24	0/871	0.41	0/1167	
12	L	0.23	0/1485	0.36	0/1980	
13	М	0.25	0/1017	0.42	0/1366	
14	Ν	0.25	0/907	0.45	0/1228	
15	Ο	0.25	0/1653	0.39	0/2254	
16	Р	0.26	0/809	0.41	0/1085	
17	Q	0.25	0/735	0.40	0/980	
18	R	0.24	0/2449	0.38	0/3311	
19	S	0.26	0/1148	0.40	0/1541	
20	Т	0.25	0/1420	0.39	0/1903	
21	U	0.24	0/1470	0.37	0/1976	
22	V	0.23	0/3059	0.35	0/4135	
23	W	0.25	0/805	0.43	0/1084	
24	Х	0.24	0/2952	0.38	0/3995	
25	Y	0.25	0/1283	0.36	0/1730	
26	Ζ	0.25	0/851	0.37	0/1133	
27	0	0.24	0/1856	0.41	0/2511	
28	1	0.24	0/2260	0.38	0/3063	
29	3	0.24	0/654	0.38	0/860	
30	4	0.24	0/4868	0.36	0/6597	
31	a	0.24	0/1573	0.37	0/2115	
32	b	0.24	$0/2\overline{561}$	0.39	0/3457	



Mal	Chain	Bo	nd lengths	Bond angles		
IVIOI		RMSZ	# Z > 5	RMSZ	# Z > 5	
All	All	0.25	1/73447~(0.0%)	0.50	0/103690	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
9	Ι	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	А	1	А	OP3-P	-10.78	1.48	1.61

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
9	Ι	181	5F0	Mainchain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	20207	10223	10209	40	0
2	В	1791	1799	1797	8	0
3	С	1072	1091	1087	7	0
4	D	2733	2794	2790	13	0
5	Е	948	968	968	3	0
6	F	1722	1750	1748	8	0
7	G	2674	2630	2626	11	0
8	Н	1153	1193	1190	10	0
9	I	1009	1038	1029	3	0
10	J	846	903	901	2	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
11	K	855	888	887	7	0
12	L	1465	1576	1574	3	0
13	М	995	1009	1006	2	0
14	N	888	951	947	4	0
15	0	1598	1548	1547	0	0
16	Р	792	818	817	2	0
17	Q	732	750	750	4	0
18	R	2400	2416	2415	6	0
19	S	1124	1133	1132	3	0
20	Т	1388	1413	1413	5	0
21	U	1449	1459	1456	0	0
22	V	2998	2972	2967	8	0
23	W	793	813	811	3	0
24	Х	2881	2881	2879	12	0
25	Y	1246	1193	1191	3	0
26	Z	834	848	847	1	0
27	0	1811	1838	1834	3	0
28	1	2212	2228	2226	5	0
29	3	643	728	726	1	0
30	4	4758	4800	4793	11	0
31	a	1546	1583	1579	0	0
32	b	2519	2595	2593	0	0
33	3	1	0	0	0	0
33	А	46	0	0	0	0
33	В	1	0	0	0	0
33	Х	1	0	0	0	0
34	А	13	0	0	0	0
35	0	1	0	0	0	0
36	Р	4	0	0	0	0
36	Т	4	0	0	0	0
37	Х	31	12	12	0	0
38	b	26	20	19	0	0
39	0	7	0	0	0	0
39	1	5	0	0	0	0
39	3	7	0	0	0	0
39	A	549	0	0	8	0
39	B	23	0	0	0	0
39	C	19	0	0	1	0
39	D	34	0	0	1	0
39	E	2	0	0	0	0
39	F	7	0	0	0	0
39	G	10	0	0	0	0

Continued from previous page...



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
39	Н	19	0	0	0	0
39	Ι	5	0	0	0	0
39	J	14	0	0	0	0
39	K	25	0	0	3	0
39	L	6	0	0	0	0
39	М	16	0	0	0	0
39	N	8	0	0	0	0
39	0	25	0	0	0	0
39	Р	3	0	0	0	0
39	Q	8	0	0	0	0
39	R	13	0	0	0	0
39	S	4	0	0	0	0
39	Т	19	0	0	0	0
39	U	3	0	0	0	0
39	W	2	0	0	0	0
39	Х	5	0	0	0	0
39	Y	7	0	0	0	0
39	Ζ	18	0	0	0	0
39	a	5	0	0	0	0
All	All	71078	60861	60766	152	0

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 152 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)	
1:A:465:A:OP1	2:B:97:ARG:NH1	2.23	0.72	
1:A:705:G:N7	39:A:1112:HOH:O	2.27	0.67	
4:D:196:SER:O	39:D:501:HOH:O	2.13	0.65	
1:A:942:G:O6	39:A:1102:HOH:O	2.12	0.64	
13:M:55:ASP:OD2	20:T:146:GLN:NE2	2.31	0.64	

There are no symmetry-related clashes.



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	s Percenti	
2	В	221/291~(76%)	217 (98%)	4(2%)	0	100	100
3	\mathbf{C}	130/167~(78%)	124 (95%)	6~(5%)	0	100	100
4	D	341/432~(79%)	336 (98%)	5(2%)	0	100	100
5	Ε	117/125~(94%)	115 (98%)	2(2%)	0	100	100
6	F	206/242~(85%)	202 (98%)	4 (2%)	0	100	100
7	G	321/390~(82%)	317 (99%)	4 (1%)	0	100	100
8	Н	138/160~(86%)	135 (98%)	2(1%)	1 (1%)	22	52
9	Ι	134/191~(70%)	129 (96%)	5(4%)	0	100	100
10	J	106/139~(76%)	105 (99%)	1 (1%)	0	100	100
11	K	99/128~(77%)	99 (100%)	0	0	100	100
12	L	174/258~(67%)	173 (99%)	1 (1%)	0	100	100
13	М	124/135~(92%)	124 (100%)	0	0	100	100
14	Ν	111/120 (92%)	108 (97%)	3(3%)	0	100	100
15	Ο	195/254~(77%)	193 (99%)	2(1%)	0	100	100
16	Р	95/143~(66%)	95 (100%)	0	0	100	100
17	Q	84/86~(98%)	83 (99%)	1 (1%)	0	100	100
18	R	292/359~(81%)	283 (97%)	9~(3%)	0	100	100
19	S	134/177~(76%)	132 (98%)	2(2%)	0	100	100
20	Т	168/171~(98%)	167 (99%)	1 (1%)	0	100	100
21	U	172/200~(86%)	171 (99%)	1 (1%)	0	100	100
22	V	361/415~(87%)	356 (99%)	5 (1%)	0	100	100
23	W	98/186~(53%)	95 (97%)	3 (3%)	0	100	100
24	Х	355/391 (91%)	349 (98%)	6 (2%)	0	100	100
25	Y	147/384~(38%)	145 (99%)	2 (1%)	0	100	100
26	Z	99/106~(93%)	98 (99%)	1 (1%)	0	100	100



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
27	0	214/218~(98%)	209~(98%)	5 (2%)	0	100	100
28	1	275/320~(86%)	273~(99%)	2(1%)	0	100	100
29	3	70/200~(35%)	70 (100%)	0	0	100	100
30	4	585/685~(85%)	580 (99%)	5 (1%)	0	100	100
31	a	183/350~(52%)	180 (98%)	3 (2%)	0	100	100
32	b	319/406~(79%)	310 (97%)	9 (3%)	0	100	100
All	All	6068/7829~(78%)	5973 (98%)	94 (2%)	1 (0%)	100	100

Continued from previous page...

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
8	Н	85	ILE

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
2	В	195/247~(79%)	195 (100%)	0	100 100		
3	С	113/139~(81%)	113 (100%)	0	100 100		
4	D	282/354~(80%)	282 (100%)	0	100 100		
5	Е	104/110~(94%)	104 (100%)	0	100 100		
6	F	180/204~(88%)	180 (100%)	0	100 100		
7	G	282/335~(84%)	281 (100%)	1 (0%)	91 95		
8	Η	130/150~(87%)	130 (100%)	0	100 100		
9	Ι	102/143~(71%)	102 (100%)	0	100 100		
10	J	93/117~(80%)	93 (100%)	0	100 100		
11	К	90/110~(82%)	90 (100%)	0	100 100		
12	L	161/224 (72%)	161 (100%)	0	100 100		
13	М	103/112~(92%)	103 (100%)	0	100 100		



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
14	Ν	97/104~(93%)	97~(100%)	0	100	100	
15	Ο	176/225~(78%)	176 (100%)	0	100	100	
16	Р	89/125~(71%)	89 (100%)	0	100	100	
17	Q	77/77~(100%)	77~(100%)	0	100	100	
18	R	261/313~(83%)	261 (100%)	0	100	100	
19	S	117/152~(77%)	117 (100%)	0	100	100	
20	Т	153/154~(99%)	153~(100%)	0	100	100	
21	U	149/171~(87%)	149 (100%)	0	100	100	
22	V	326/362~(90%)	325~(100%)	1 (0%)	92	96	
23	W	87/156~(56%)	87~(100%)	0	100	100	
24	Х	314/346~(91%)	312~(99%)	2(1%)	86	93	
25	Υ	133/341~(39%)	133 (100%)	0	100	100	
26	Ζ	88/93~(95%)	88 (100%)	0	100	100	
27	0	190/191~(100%)	189 (100%)	1 (0%)	88	94	
28	1	247/279~(88%)	247~(100%)	0	100	100	
29	3	68/176~(39%)	68~(100%)	0	100	100	
30	4	519/599~(87%)	519 (100%)	0	100	100	
31	a	174/299~(58%)	174 (100%)	0	100	100	
32	b	270/343~(79%)	270 (100%)	0	100	100	
All	All	5370/6751~(80%)	5365 (100%)	5 (0%)	93	97	

Continued from previous page...

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
7	G	375	LYS
22	V	226	TYR
24	Х	74	HIS
24	Х	112	HIS
27	0	48	ARG

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such side chains are listed below:

Mol	Chain	Res	Type
7	G	139	GLN



Continued from previous page...

Mol	Chain	Res	Type
26	Ζ	82	GLN
30	4	372	HIS
28	1	182	HIS
4	D	279	HIS

5.3.3 RNA (i)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	А	946/956~(98%)	123~(13%)	0

5 of 123 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	А	3	А
1	А	7	U
1	А	32	G
1	А	40	А
1	А	41	U

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Iol Type Chain		Dog	Tink	Bond lengths		Bond angles			
	Type Chain	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2	
17	AYA	Q	2	17	6,7,8	0.77	0	$5,\!8,\!10$	0.57	0
1	MA6	А	937	1	18,26,27	0.75	0	$19,\!38,\!41$	0.54	0
1	5MC	А	842	1	18,22,23	0.32	0	$26,\!32,\!35$	0.40	0
1	B8T	А	840	1	19,22,23	0.30	0	$26,\!31,\!34$	0.32	0
9	5F0	Ι	181	9	8,8,9	0.55	0	7,9,11	1.05	1 (14%)
1	MA6	А	938	1	18,26,27	0.75	0	19,38,41	0.58	0



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
17	AYA	Q	2	17	-	0/4/6/8	-
1	MA6	А	937	1	-	0/7/29/30	0/3/3/3
1	5MC	А	842	1	-	2/7/25/26	0/2/2/2
1	B8T	А	840	1	-	0/7/27/28	0/2/2/2
9	5F0	Ι	181	9	-	0/9/9/10	-
1	MA6	А	938	1	-	1/7/29/30	0/3/3/3

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
9	Ι	181	$5\mathrm{F0}$	O-C-CB	-2.44	118.31	125.43

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	А	938	MA6	C4'-C5'-O5'-P
1	А	842	5MC	O4'-C4'-C5'-O5'
1	А	842	5MC	C3'-C4'-C5'-O5'

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 67 ligands modelled in this entry, 63 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



Mal	Turne	Chain	Dec	Tiple	Link Bond lengths				Bond angles		
	туре	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2	
37	ATP	Х	401	33	26,33,33	0.75	0	31,52,52	0.66	0	
36	FES	Т	201	13,20	0,4,4	-	-	-			
38	SAH	b	501	-	24,28,28	0.70	0	25,40,40	0.78	1 (4%)	
36	FES	Р	201	5,16	0,4,4	-	-	-			

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
37	ATP	Х	401	33	-	0/18/38/38	0/3/3/3
36	FES	Т	201	13,20	-	-	0/1/1/1
38	SAH	b	501	-	-	2/11/31/31	0/3/3/3
36	FES	Р	201	5,16	-	-	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
38	b	501	SAH	C5-C6-N6	2.35	123.92	120.35

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
38	b	501	SAH	C4'-C5'-SD-CG
38	b	501	SAH	CA-CB-CG-SD

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.



Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-13553. These allow visual inspection of the internal detail of the map and identification of artifacts.

Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections (i)

6.1.1 Primary map



6.1.2 Raw map



The images above show the map projected in three orthogonal directions.



6.2 Central slices (i)

6.2.1 Primary map



X Index: 288



Y Index: 288



Z Index: 288

6.2.2 Raw map



X Index: 240

Y Index: 240



The images above show central slices of the map in three orthogonal directions.



6.3 Largest variance slices (i)

6.3.1 Primary map



X Index: 301





Z Index: 276

6.3.2 Raw map



X Index: 262

Y Index: 256



The images above show the largest variance slices of the map in three orthogonal directions.



6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map



6.4.2 Raw map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.



6.5 Orthogonal surface views (i)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.1. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.



Mask visualisation (i) 6.6

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency

A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure

$emd_{13553}msk_{1.map}$ (i) 6.6.1





7 Map analysis (i)

This section contains the results of statistical analysis of the map.

7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)



The volume at the recommended contour level is 564 nm^3 ; this corresponds to an approximate mass of 509 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)



*Reported resolution corresponds to spatial frequency of 0.327 \AA^{-1}



8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.327 ${\rm \AA^{-1}}$



8.2 Resolution estimates (i)

$\begin{bmatrix} Bosolution ostimato (Å) \end{bmatrix}$	Estim	Estimation criterion (FSC cut-off			
resolution estimate (A)	0.143	0.5	Half-bit		
Reported by author	3.06	-	-		
Author-provided FSC curve	3.05	3.91	3.10		
Unmasked-calculated*	3.93	8.48	4.13		

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.93 differs from the reported value 3.06 by more than 10 %



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-13553 and PDB model 7PNV. Per-residue inclusion information can be found in section 3 on page 13.

9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.1 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.



9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.1).



9.4 Atom inclusion (i)



At the recommended contour level, 85% of all backbone atoms, 87% of all non-hydrogen atoms, are inside the map.



1.0

0.0 <0.0

9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.1) and Q-score for the entire model and for each chain.

\mathbf{Chain}	Atom inclusion	Q-score
All	0.8700	0.5440
0	0.8980	0.5550
1	0.8530	0.5440
3	0.8640	0.5440
4	0.6620	0.4290
А	0.9580	0.5940
В	0.9470	0.6110
С	0.9740	0.6360
D	0.9070	0.5720
Е	0.9240	0.5700
F	0.9000	0.5420
G	0.8570	0.5340
Н	0.9090	0.5860
Ι	0.9480	0.5800
J	0.9500	0.5900
К	0.9590	0.6300
L	0.8420	0.5400
М	0.9000	0.6030
Ν	0.9200	0.5940
О	0.9360	0.5950
Р	0.9360	0.5840
Q	0.9700	0.6090
R	0.8850	0.5620
S	0.8390	0.5160
Т	0.9200	0.5780
U	0.8310	0.5230
V	0.7750	0.4790
W	0.9260	0.5870
Х	0.8480	0.5230
Y	0.7040	0.4890
Ζ	0.8660	0.5490
a	0.6750	0.4090
b	0.2530	0.2420

