



Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 04:13 pm BST

PDB ID : 1PO1
Title : POLIOVIRUS (TYPE 1, MAHONEY) IN COMPLEX WITH R80633, AN INHIBITOR OF VIRAL REPLICATION
Authors : Hiremath, C.N.; Filman, D.J.; Grant, R.A.; Hogle, J.M.
Deposited on : 1997-01-08
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	0	5	100%
2	1	302	79% 14% • 6%
3	2	272	85% 12% ••
4	3	238	87% 12% •
5	4	68	75% 16% 9%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6690 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called POLIOVIRUS TYPE 1 MAHONEY.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
1	0	5	29	15	5	9	0	0	0

- Molecule 2 is a protein called POLIOVIRUS TYPE 1 MAHONEY.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	1	283	2222	1416	378	423	5	0	0	0

- Molecule 3 is a protein called POLIOVIRUS TYPE 1 MAHONEY.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	2	268	2085	1317	358	396	14	0	0	0

- Molecule 4 is a protein called POLIOVIRUS TYPE 1 MAHONEY.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	3	235	1834	1169	299	349	17	0	0	0


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
3	123	SER	PHE	CONFLICT	UNP P03300

- Molecule 5 is a protein called POLIOVIRUS TYPE 1 MAHONEY.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	4	62	476	293	81	101	1	0	0	0

- Molecule 6 is (METHYLPYRIDAZINE PIPERIDINE BUTYLOXYPHENYL)ETHYLACETATE (three-letter code: J80) (formula: $C_{23}H_{31}N_3O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
6	1	1	29	23	3	3	0	0

- Molecule 7 is MYRISTIC ACID (three-letter code: MYR) (formula: $C_{14}H_{28}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
7	4	1	15	14	1	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: POLIOVIRUS TYPE 1 MAHONEY

Chain 0:  100%


There are no outlier residues recorded for this chain.



- Molecule 2: POLIOVIRUS TYPE 1 MAHONEY

Chain 1:  79% 14% • 6%



- Molecule 3: POLIOVIRUS TYPE 1 MAHONEY

Chain 2:  85% 12% ••



- Molecule 4: POLIOVIRUS TYPE 1 MAHONEY

Chain 3:  87% 12% •



- Molecule 5: POLIOVIRUS TYPE 1 MAHONEY

Chain 4:  75% 16% 9%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	322.94Å 358.04Å 380.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 2.90 30.95 – 2.87	Depositor EDS
% Data completeness (in resolution range)	(Not available) ((Not available)-2.90) 26.7 (30.95-2.87)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.90 (at 2.85Å)	Xtrriage
Refinement program	X-PLOR	Depositor
R, R_{free}	0.241 , (Not available) 0.213 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	30.8	Xtrriage
Anisotropy	0.079	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , -39.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.22	EDS
Total number of atoms	6690	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: J80, MYR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	0	0.88	0/28	1.62	0/36
2	1	0.76	0/2285	1.39	16/3124 (0.5%)
3	2	0.77	0/2142	1.45	26/2928 (0.9%)
4	3	0.76	0/1881	1.32	12/2562 (0.5%)
5	4	0.75	0/483	1.43	2/651 (0.3%)
All	All	0.76	0/6819	1.39	56/9301 (0.6%)

There are no bond length outliers.

All (56) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	1	269	TRP	CD1-CG-CD2	9.20	113.66	106.30
3	2	227	TRP	CD1-CG-CD2	9.00	113.50	106.30
3	2	38	TRP	CD1-CG-CD2	8.72	113.28	106.30
3	2	78	TRP	CD1-CG-CD2	8.71	113.27	106.30
4	3	110	TRP	CD1-CG-CD2	8.47	113.08	106.30
3	2	79	TRP	CD1-CG-CD2	8.25	112.90	106.30
2	1	269	TRP	CE2-CD2-CG	-8.03	100.88	107.30
4	3	110	TRP	CE2-CD2-CG	-8.01	100.89	107.30
2	1	175	TRP	CD1-CG-CD2	7.99	112.69	106.30
3	2	80	TRP	CD1-CG-CD2	7.98	112.68	106.30
3	2	80	TRP	CE2-CD2-CG	-7.92	100.96	107.30
3	2	227	TRP	CE2-CD2-CG	-7.92	100.97	107.30
4	3	170	TRP	CE2-CD2-CG	-7.91	100.97	107.30
3	2	71	TRP	CD1-CG-CD2	7.80	112.54	106.30
3	2	71	TRP	CE2-CD2-CG	-7.79	101.07	107.30
3	2	78	TRP	CE2-CD2-CG	-7.74	101.11	107.30
2	1	170	TRP	CD1-CG-CD2	7.71	112.47	106.30
4	3	170	TRP	CD1-CG-CD2	7.64	112.41	106.30
2	1	175	TRP	CE2-CD2-CG	-7.57	101.25	107.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	2	38	TRP	CE2-CD2-CG	-7.55	101.26	107.30
2	1	108	TRP	CD1-CG-CD2	7.38	112.21	106.30
2	1	108	TRP	CE2-CD2-CG	-7.37	101.41	107.30
2	1	170	TRP	CE2-CD2-CG	-7.15	101.58	107.30
3	2	79	TRP	CE2-CD2-CG	-6.99	101.71	107.30
2	1	275	ARG	NE-CZ-NH1	6.81	123.70	120.30
4	3	156	TRP	CD1-CG-CD2	6.76	111.71	106.30
4	3	226	ARG	NE-CZ-NH1	6.59	123.60	120.30
4	3	156	TRP	CE2-CD2-CG	-6.59	102.03	107.30
2	1	120	ARG	NE-CZ-NH1	6.51	123.56	120.30
2	1	258	ARG	NE-CZ-NH1	6.44	123.52	120.30
4	3	226	ARG	NE-CZ-NH2	-6.43	117.08	120.30
3	2	204	ASN	CB-CA-C	-6.39	97.62	110.40
3	2	172	ARG	NE-CZ-NH1	6.20	123.40	120.30
5	4	34	ARG	NE-CZ-NH2	-6.07	117.27	120.30
2	1	269	TRP	CB-CG-CD1	-5.91	119.32	127.00
3	2	37	ARG	NE-CZ-NH1	5.85	123.22	120.30
4	3	170	TRP	CG-CD2-CE3	5.83	139.15	133.90
4	3	170	TRP	CB-CG-CD1	-5.72	119.56	127.00
3	2	79	TRP	CB-CG-CD1	-5.67	119.62	127.00
5	4	34	ARG	NE-CZ-NH1	5.67	123.14	120.30
3	2	8	GLY	O-C-N	-5.65	113.65	122.70
4	3	223	ARG	NE-CZ-NH2	-5.61	117.49	120.30
3	2	227	TRP	CG-CD2-CE3	5.53	138.87	133.90
3	2	227	TRP	CG-CD1-NE1	-5.48	104.62	110.10
2	1	280	TYR	CB-CG-CD2	-5.38	117.77	121.00
4	3	110	TRP	CG-CD2-CE3	5.35	138.72	133.90
2	1	129	ARG	NE-CZ-NH1	5.34	122.97	120.30
2	1	269	TRP	CG-CD1-NE1	-5.32	104.78	110.10
3	2	8	GLY	N-CA-C	5.25	126.21	113.10
3	2	38	TRP	CG-CD1-NE1	-5.21	104.89	110.10
2	1	175	TRP	CG-CD1-NE1	-5.18	104.92	110.10
3	2	78	TRP	CG-CD1-NE1	-5.15	104.95	110.10
3	2	79	TRP	CG-CD1-NE1	-5.12	104.98	110.10
3	2	62	ARG	NE-CZ-NH2	-5.10	117.75	120.30
3	2	78	TRP	CB-CG-CD1	-5.10	120.37	127.00
3	2	71	TRP	CG-CD2-CE3	5.06	138.45	133.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	0	29	0	24	0	0
2	1	2222	0	2173	20	0
3	2	2085	0	2000	14	0
4	3	1834	0	1816	13	0
5	4	476	0	457	4	0
6	1	29	0	31	3	0
7	4	15	0	27	1	0
All	All	6690	0	6528	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:1:158:MET:SD	2:1:177:THR:HG23	2.29	0.71
2:1:177:THR:HG22	2:1:180:ASN:HB2	1.74	0.70
4:3:51:THR:HG21	4:3:99:MET:H	1.61	0.65
3:2:143:THR:HG23	3:2:173:ARG:HA	1.82	0.60
3:2:37:ARG:HG3	4:3:37:PRO:HB3	1.85	0.58
2:1:177:THR:HG21	2:1:182:SER:OG	2.04	0.57
2:1:183:ILE:HG13	6:1:0:J80:H12	1.85	0.57
3:2:213:VAL:HG22	4:3:37:PRO:HG2	1.86	0.55
3:2:30:ASN:HD21	5:4:59:ASP:HB2	1.70	0.55
2:1:191:PRO:HG2	4:3:13:TYR:HB2	1.90	0.53
2:1:109:LYS:HA	2:1:239:ILE:HG22	1.91	0.52
5:4:10:VAL:HG21	5:4:25:ILE:HD12	1.91	0.52
2:1:22:THR:HG22	2:1:24:ARG:H	1.75	0.52
2:1:107:VAL:HG13	2:1:239:ILE:HD13	1.92	0.51
4:3:87:LEU:HD11	4:3:114:LEU:HD12	1.93	0.49
2:1:48:GLU:HA	3:2:197:ILE:HB	1.94	0.49
4:3:120:PHE:HA	4:3:210:ILE:HG22	1.94	0.49
4:3:167:VAL:O	4:3:169:PRO:HD3	2.12	0.48
5:4:30:ILE:HD13	7:4:1:MYR:H72	1.95	0.48
3:2:5:GLU:HB3	3:2:9:TYR:H	1.78	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:4:57:ILE:HD11	5:4:61:LEU:HB3	1.96	0.48
2:1:273:PRO:HB3	3:2:189:ASN:HB2	1.96	0.48
2:1:96:ALA:HA	2:1:249:ASN:O	2.14	0.47
2:1:237:PHE:HB3	6:1:0:J80:H132	1.95	0.47
4:3:61:LYS:HD3	4:3:66:GLU:HB3	1.98	0.46
4:3:55:PHE:HE2	4:3:212:GLY:HA3	1.82	0.45
3:2:5:GLU:HG3	3:2:7:CYS:H	1.81	0.45
4:3:53:ILE:HD11	4:3:214:VAL:HB	1.99	0.45
3:2:32:VAL:HB	3:2:208:LEU:HD23	1.99	0.44
2:1:181:PRO:HB2	6:1:0:J80:H3	1.98	0.44
3:2:27:GLU:HB2	3:2:204:ASN:OD1	2.18	0.43
3:2:187:LEU:HD22	4:3:65:MET:CE	2.49	0.43
2:1:160:VAL:HB	2:1:239:ILE:HG13	2.01	0.42
4:3:64:THR:O	4:3:67:MET:HG2	2.19	0.42
2:1:302:TYR:CE1	4:3:189:TYR:HB3	2.55	0.42
2:1:286:TYR:HB2	2:1:291:LEU:HD21	2.02	0.42
2:1:24:ARG:HA	2:1:71:SER:OG	2.20	0.42
2:1:216:PRO:HA	2:1:225:GLY:HA3	2.02	0.42
3:2:66:LEU:HD12	3:2:251:LEU:HD23	2.02	0.41
2:1:89:ILE:HG12	2:1:258:ARG:HG2	2.02	0.41
3:2:201:ARG:HH11	3:2:201:ARG:HD3	1.77	0.40
2:1:95:PRO:O	2:1:248:HIS:HB3	2.21	0.40
3:2:134:GLY:HA2	3:2:174:PHE:HA	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	0	3/5 (60%)	3 (100%)	0	0	100 100
2	1	281/302 (93%)	268 (95%)	12 (4%)	1 (0%)	34 66

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	2	266/272 (98%)	248 (93%)	16 (6%)	2 (1%)	19	51
4	3	233/238 (98%)	220 (94%)	13 (6%)	0	100	100
5	4	58/68 (85%)	52 (90%)	4 (7%)	2 (3%)	3	15
All	All	841/885 (95%)	791 (94%)	45 (5%)	5 (1%)	25	58

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	2	240	ALA
3	2	48	ASN
5	4	11	GLY
2	1	270	CYS
5	4	60	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	0	4/4 (100%)	4 (100%)	0	100	100
2	1	245/261 (94%)	235 (96%)	10 (4%)	30	64
3	2	228/232 (98%)	219 (96%)	9 (4%)	32	66
4	3	210/212 (99%)	205 (98%)	5 (2%)	49	79
5	4	54/57 (95%)	52 (96%)	2 (4%)	34	68
All	All	741/766 (97%)	715 (96%)	26 (4%)	36	70

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	1	83	ARG
2	1	99	THR
2	1	132	MET
2	1	147	ASN
2	1	149	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	1	177	THR
2	1	181	PRO
2	1	194	ILE
2	1	220	GLN
2	1	295	SER
3	2	52	GLN
3	2	66	LEU
3	2	168	THR
3	2	201	ARG
3	2	204	ASN
3	2	238	ASN
3	2	241	SER
3	2	264	ARG
3	2	272	GLN
4	3	51	THR
4	3	85	LEU
4	3	163	SER
4	3	208	MET
4	3	218	ASN
5	4	16	SER
5	4	69	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
2	1	100	ASN
4	3	6	ASN
4	3	218	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	MYR	4	1	5	14,14,15	0.36	0	13,13,15	0.58	0
6	J80	1	0	-	31,31,31	1.46	3 (9%)	39,40,40	2.16	10 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	MYR	4	1	5	-	4/11/12/13	-
6	J80	1	0	-	-	5/19/29/29	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	1	0	J80	O25-C23	5.06	1.46	1.33
6	1	0	J80	C20-C23	-4.74	1.38	1.50
6	1	0	J80	O25-C26	-2.02	1.40	1.46

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	1	0	J80	C1-C2-N3	6.24	119.18	116.24
6	1	0	J80	C4-C5-N4	-4.87	116.59	123.86
6	1	0	J80	C11-N6-C7	4.81	122.14	111.52
6	1	0	J80	O25-C26-C27	4.74	125.82	108.42
6	1	0	J80	C5-N4-N3	4.39	123.46	118.97
6	1	0	J80	C7-C8-C9	2.91	119.37	111.99
6	1	0	J80	O25-C23-C20	2.89	117.17	112.14
6	1	0	J80	C8-C7-N6	2.68	116.63	111.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	1	0	J80	C13-C14-C15	-2.50	102.41	113.49
6	1	0	J80	C3-C4-C5	2.36	120.78	117.53

There are no chirality outliers.

All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	1	0	J80	C18-C17-O16-C15
7	4	1	MYR	C6-C7-C8-C9
6	1	0	J80	C22-C17-O16-C15
7	4	1	MYR	C11-C12-C13-C14
6	1	0	J80	O24-C23-O25-C26
7	4	1	MYR	C10-C11-C12-C13
6	1	0	J80	C14-C15-O16-C17
6	1	0	J80	C13-C14-C15-O16
7	4	1	MYR	C5-C6-C7-C8

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	4	1	MYR	1	0
6	1	0	J80	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [\(i\)](#)

6.1 Protein, DNA and RNA chains [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.


6.3 Carbohydrates [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers

Unable to reproduce the depositor's R factor - this section is therefore empty.