



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 4, 2024 – 12:12 AM EST

PDB ID : 1PP1
Title : Crystal structure of the Borna Disease Virus Nucleoprotein
Authors : Rudolph, M.G.; Kraus, I.; Dickmanns, A.; Garten, W.; Ficner, R.
Deposited on : 2003-06-16
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 1.20.1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	7293 (1.90-1.90)
Clashscore	180529	8090 (1.90-1.90)
Ramachandran outliers	177936	8022 (1.90-1.90)
Sidechain outliers	177891	8022 (1.90-1.90)
RSRZ outliers	164620	7292 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	X	375	 6% 81% 7% • 11%

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 2988 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called p40 nucleoprotein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	X	335	2606	1660	444	488	4	10	0	7	0

There are 17 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
X	-4	VAL	-	cloning artifact	UNP Q01552
X	-3	GLU	-	cloning artifact	UNP Q01552
X	-2	PHE	-	cloning artifact	UNP Q01552
X	-1	GLY	-	cloning artifact	UNP Q01552
X	0	SER	-	cloning artifact	UNP Q01552
X	1	MSE	MET	modified residue	UNP Q01552
X	14	MSE	MET	modified residue	UNP Q01552
X	68	MSE	MET	modified residue	UNP Q01552
X	169	MSE	MET	modified residue	UNP Q01552
X	170	MSE	MET	modified residue	UNP Q01552
X	187	MSE	MET	modified residue	UNP Q01552
X	223	MSE	MET	modified residue	UNP Q01552
X	235	MSE	MET	modified residue	UNP Q01552
X	249	MSE	MET	modified residue	UNP Q01552
X	317	MSE	MET	modified residue	UNP Q01552
X	361	MSE	MET	modified residue	UNP Q01552
X	363	MSE	MET	modified residue	UNP Q01552


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	X	382	Total	O	0	0
			382	382		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: p40 nucleoprotein



4 Data and refinement statistics i

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, α , β , γ	100.26Å 100.26Å 103.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.23 – 1.90 29.23 – 1.90	Depositor EDS
% Data completeness (in resolution range)	96.1 (29.23-1.90) 96.0 (29.23-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.07 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.1.19	Depositor
R, R_{free}	0.153 , 0.182 0.165 , 0.193	Depositor DCC
R_{free} test set	2018 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	19.4	Xtriage
Anisotropy	0.141	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 46.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.002 for l,-k,h 0.014 for -l,-k,-h 0.017 for -h,-l,-k 0.003 for -h,l,k 0.037 for -h,k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2988	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	X	0.68	0/2686	0.80	7/3620 (0.2%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	X	54	ARG	NE-CZ-NH2	-5.61	117.49	120.30
1	X	187[A]	MSE	CG-SE-CE	5.54	111.08	98.90
1	X	187[B]	MSE	CG-SE-CE	5.54	111.08	98.90
1	X	62	ASP	CB-CG-OD2	5.36	123.13	118.30
1	X	338	ARG	NE-CZ-NH1	5.25	122.92	120.30
1	X	342	ARG	NE-CZ-NH2	-5.19	117.71	120.30
1	X	287	ARG	NE-CZ-NH2	-5.17	117.72	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	X	2606	0	2617	19	0
2	X	382	0	0	5	1
All	All	2988	0	2617	19	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:55:GLN:NE2	1:X:123:THR:HG23	2.06	0.70
1:X:186:GLN:HG3	1:X:187[A]:MSE:HE3	1.76	0.68
1:X:92:HIS:ND1	2:X:569:HOH:O	2.27	0.68
1:X:130:GLU:OE1	1:X:175:ARG:NH2	2.29	0.64
1:X:155:ILE:O	2:X:581:HOH:O	2.19	0.54
1:X:55:GLN:HE21	1:X:123:THR:HG23	1.76	0.50
1:X:40:GLY:N	2:X:454:HOH:O	2.47	0.47
1:X:160:GLU:CG	2:X:576:HOH:O	2.62	0.47
1:X:123:THR:O	1:X:123:THR:HG22	2.17	0.45
1:X:344:ILE:HD12	1:X:344:ILE:C	2.38	0.44
1:X:309:TRP:CE2	1:X:313:GLU:HG3	2.53	0.44
1:X:62:ASP:OD2	1:X:64:SER:OG	2.26	0.43
1:X:160:GLU:HG2	2:X:576:HOH:O	2.18	0.43
1:X:155:ILE:HD12	1:X:155:ILE:N	2.34	0.42
1:X:186:GLN:HG3	1:X:187[A]:MSE:CE	2.47	0.42
1:X:289:PRO:O	1:X:292:ILE:HG22	2.20	0.42
1:X:47:ILE:HD12	1:X:87:HIS:CD2	2.56	0.41
1:X:154:LYS:O	1:X:158:GLY:N	2.54	0.41
1:X:103:THR:HA	1:X:104:PRO:HD3	1.94	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:X:564:HOH:O	2:X:695:HOH:O[3_655]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	X	338/375 (90%)	331 (98%)	5 (2%)	2 (1%)	22 13

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	X	122	LYS
1	X	158	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	X	287/304 (94%)	283 (99%)	4 (1%)	62 62

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	X	38	VAL
1	X	103	THR
1	X	338	ARG
1	X	348	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	X	55	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	X	326/375 (86%)	-0.23	22 (6%) 25 26	12, 19, 45, 61	6 (1%)

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	X	158	GLY	5.7
1	X	106	THR	4.3
1	X	123	THR	4.2
1	X	351	ALA	4.1
1	X	108	GLY	4.0
1	X	121	GLU	3.7
1	X	348	GLU	3.6
1	X	39	GLY	3.5
1	X	350	GLY	3.5
1	X	124	THR	3.3
1	X	157	ALA	3.2
1	X	323	SER	3.1
1	X	107	ARG	3.0
1	X	110	GLN	3.0
1	X	105	VAL	2.9
1	X	155	ILE	2.8
1	X	349	ASP	2.7
1	X	109	GLU	2.5
1	X	314	ASN	2.4
1	X	159	ALA	2.3
1	X	122	LYS	2.2
1	X	38	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.