



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 23, 2021 – 06:05 pm GMT

PDB ID : 7PQ7
Title : Crystal structure of Campylobacter jejuni DsbA1
Authors : Wilk, P.; Orlikowska, M.; Banas, A.M.; Bocian-Ostrzycka, K.M.; Jagusztyn-Krynicka, E.K.
Deposited on : 2021-09-16
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.24
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.24


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1483 (1.56-1.56)
Clashscore	141614	1529 (1.56-1.56)
Ramachandran outliers	138981	1498 (1.56-1.56)
Sidechain outliers	138945	1495 (1.56-1.56)
RSRZ outliers	127900	1465 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	202	 2% 83% 11% • 5%
1	B	202	 6% 83% 10% 7%

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 6716 atoms, of which 3147 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Thiol:disulfide interchange protein DsbA.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	191	3129	1016	1549	258	298	8	0	5	0
1	B	188	3149	1021	1566	256	298	8	0	12	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP A0A1J6PBD5
A	195	LEU	-	expression tag	UNP A0A1J6PBD5
A	196	GLU	-	expression tag	UNP A0A1J6PBD5
A	197	HIS	-	expression tag	UNP A0A1J6PBD5
A	198	HIS	-	expression tag	UNP A0A1J6PBD5
A	199	HIS	-	expression tag	UNP A0A1J6PBD5
A	200	HIS	-	expression tag	UNP A0A1J6PBD5
A	201	HIS	-	expression tag	UNP A0A1J6PBD5
A	202	HIS	-	expression tag	UNP A0A1J6PBD5
B	1	MET	-	initiating methionine	UNP A0A1J6PBD5
B	195	LEU	-	expression tag	UNP A0A1J6PBD5
B	196	GLU	-	expression tag	UNP A0A1J6PBD5
B	197	HIS	-	expression tag	UNP A0A1J6PBD5
B	198	HIS	-	expression tag	UNP A0A1J6PBD5
B	199	HIS	-	expression tag	UNP A0A1J6PBD5
B	200	HIS	-	expression tag	UNP A0A1J6PBD5
B	201	HIS	-	expression tag	UNP A0A1J6PBD5
B	202	HIS	-	expression tag	UNP A0A1J6PBD5

- Molecule 2 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: C₆H₁₄O₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
2	A	1	24	6	14	4	0	0

- Molecule 3 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: C₈H₁₈O₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
3	A	1	31	8	18	5	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	196	Total 196	O 196	0	0
4	B	187	Total 187	O 187	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Thiol:disulfide interchange protein DsbA



- Molecule 1: Thiol:disulfide interchange protein DsbA



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	120.88Å 51.73Å 75.54Å 90.00° 125.14° 90.00°	Depositor
Resolution (Å)	42.65 – 1.55 42.65 – 1.55	Depositor EDS
% Data completeness (in resolution range)	98.9 (42.65-1.55) 99.0 (42.65-1.55)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.97 (at 1.55Å)	Xtriage
Refinement program	PHENIX 1.16_3549	Depositor
R, R_{free}	0.177 , 0.200 0.177 , 0.201	Depositor DCC
R_{free} test set	2101 reflections (3.81%)	wwPDB-VP
Wilson B-factor (Å ²)	26.0	Xtriage
Anisotropy	0.473	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6716	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PGE, PG4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.11	9/1632 (0.6%)	1.85	5/2200 (0.2%)
1	B	0.87	0/1655	0.89	2/2230 (0.1%)
All	All	1.00	9/3287 (0.3%)	1.45	7/4430 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	110	ARG	CZ-NH1	-15.53	1.12	1.33
1	A	110	ARG	CD-NE	-10.43	1.28	1.46
1	A	110	ARG	NE-CZ	-8.16	1.22	1.33
1	A	55	TYR	CG-CD1	-7.67	1.29	1.39
1	A	188	LYS	CE-NZ	7.32	1.67	1.49
1	A	77[A]	TYR	CD1-CE1	6.39	1.49	1.39
1	A	77[B]	TYR	CD1-CE1	6.39	1.49	1.39
1	A	105	PHE	CE2-CZ	5.78	1.48	1.37
1	A	188	LYS	CD-CE	5.65	1.65	1.51

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	110	ARG	NE-CZ-NH1	-64.56	88.02	120.30
1	A	110	ARG	NE-CZ-NH2	39.44	140.02	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	110	ARG	NH1-CZ-NH2	10.24	130.67	119.40
1	A	60	MET	CG-SD-CE	5.58	109.14	100.20
1	B	126	MET	CG-SD-CE	5.37	108.79	100.20
1	A	55	TYR	CA-CB-CG	5.28	123.43	113.40
1	B	177	ILE	N-CA-C	-5.18	97.02	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	177	ILE	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1580	1549	1549	15	2
1	B	1583	1566	1564	14	2
2	A	10	14	14	0	0
3	A	13	18	18	0	0
4	A	196	0	0	7	0
4	B	187	0	0	6	0
All	All	3569	3147	3145	26	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:150:GLN:NE2	4:A:401:HOH:O	1.70	0.97
1:B:17:GLU:OE1	4:B:301:HOH:O	1.83	0.95
1:B:111[A]:ASN:OD1	4:B:302:HOH:O	1.87	0.92
1:A:150:GLN:CD	4:A:403:HOH:O	2.14	0.85
1:B:150:GLN:NE2	4:B:303:HOH:O	2.11	0.82

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:113:SER:OG	1:B:117:GLU:OE1	2.05	0.73
1:A:150:GLN:NE2	4:A:403:HOH:O	2.19	0.72
1:B:111[B]:ASN:ND2	4:B:304:HOH:O	2.20	0.64
1:A:41:GLU:OE2	4:A:402:HOH:O	2.15	0.63
1:B:143[A]:LYS:NZ	4:B:305:HOH:O	2.26	0.59
1:A:48:PRO:HB2	1:B:48:PRO:HB2	1.89	0.54
1:B:60:MET:HE3	4:B:463:HOH:O	2.09	0.52
1:A:68[B]:MET:HG2	1:A:118:PHE:HZ	1.75	0.52
1:A:49:ASN:HA	1:B:49:ASN:HA	1.95	0.48
1:B:16[A]:SER:OG	1:B:168:GLY:HA3	2.12	0.48
1:A:111:ASN:ND2	4:A:417:HOH:O	2.47	0.47
1:A:16:SER:OG	1:A:168:GLY:HA3	2.15	0.46
1:A:32:TYR:CE2	1:A:36:LYS:HE2	2.50	0.46
1:A:48:PRO:HB2	1:B:48:PRO:CB	2.46	0.46
1:A:90:ASP:O	1:A:95:LYS:HE2	2.19	0.43
1:A:150:GLN:OE1	4:A:403:HOH:O	2.19	0.43
1:A:48:PRO:O	4:A:405:HOH:O	2.21	0.42
1:A:120:ASP:O	1:A:124:LYS:HG3	2.19	0.41
1:B:67:GLU:HG2	1:B:115[B]:LEU:HD11	2.03	0.40
1:B:23:ALA:HA	1:B:55:TYR:O	2.22	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:110:ARG:HH12	1:B:177:ILE:O[3_455]	1.21	0.39
1:A:110:ARG:NH1	1:B:177:ILE:O[3_455]	2.03	0.17

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	194/202 (96%)	193 (100%)	1 (0%)	0	100	100
1	B	196/202 (97%)	194 (99%)	2 (1%)	0	100	100
All	All	390/404 (96%)	387 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	178/183 (97%)	173 (97%)	5 (3%)	43	14
1	B	181/183 (99%)	177 (98%)	4 (2%)	52	23
All	All	359/366 (98%)	350 (98%)	9 (2%)	50	18

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	77[A]	TYR
1	A	77[B]	TYR
1	A	119	TYR
1	A	151	ARG
1	A	194	LYS
1	B	30	HIS
1	B	89	SER
1	B	119	TYR
1	B	151	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	127	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PGE	A	301	-	9,9,9	0.21	0	8,8,8	0.16	0
3	PG4	A	302	-	12,12,12	0.23	0	11,11,11	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PGE	A	301	-	-	3/7/7/7	-
3	PG4	A	302	-	-	3/10/10/10	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	302	PG4	O3-C5-C6-O4
2	A	301	PGE	C4-C3-O2-C2
3	A	302	PG4	O1-C1-C2-O2
2	A	301	PGE	O3-C5-C6-O4
3	A	302	PG4	O2-C3-C4-O3
2	A	301	PGE	O2-C3-C4-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	191/202 (94%)	0.33	4 (2%) 63 69	21, 33, 52, 89	0
1	B	188/202 (93%)	0.66	13 (6%) 16 20	21, 35, 60, 96	0
All	All	379/404 (93%)	0.50	17 (4%) 33 38	21, 34, 56, 96	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	177	ILE	4.7
1	B	113	SER	4.2
1	A	194	LYS	4.1
1	B	118	PHE	3.8
1	A	193	MET	3.8
1	A	77[A]	TYR	3.4
1	B	193	MET	3.4
1	B	112	PHE	3.3
1	B	111[A]	ASN	3.0
1	B	192	ASN	2.8
1	B	106	LEU	2.5
1	A	112	PHE	2.5
1	B	185	ASP	2.3
1	B	77	TYR	2.2
1	B	51[A]	HIS	2.2
1	B	115[A]	LEU	2.1
1	B	178	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PG4	A	302	13/13	0.77	0.17	39,56,75,80	0
2	PGE	A	301	10/10	0.80	0.12	45,64,73,74	0

6.5 Other polymers [i](#)

There are no such residues in this entry.