



Full wwPDB X-ray Structure Validation Report ⓘ

May 18, 2020 – 08:54 pm BST

PDB ID : 3PTW
Title : CRYSTAL STRUCTURE OF malonyl CoA-acyl carrier protein transacylase from *Clostridium perfringens* Atcc 13124
Authors : Malashkevich, V.N.; Toro, R.; Ramagopal, U.; Seidel, R.; Foti, R.; Almo, S.C.; New York Structural Genomics Research Consortium (NYSGRC)
Deposited on : 2010-12-03
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	336	

2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 2506 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Malonyl CoA-acyl carrier protein transacylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	316	2416	1540	391	471	14	0	0	0

There are 23 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	VAL	-	EXPRESSION TAG	UNP Q0TRG9
A	315	ALA	-	EXPRESSION TAG	UNP Q0TRG9
A	316	GLU	-	EXPRESSION TAG	UNP Q0TRG9
A	317	ASN	-	EXPRESSION TAG	UNP Q0TRG9
A	318	LEU	-	EXPRESSION TAG	UNP Q0TRG9
A	319	TYR	-	EXPRESSION TAG	UNP Q0TRG9
A	320	PHE	-	EXPRESSION TAG	UNP Q0TRG9
A	321	GLN	-	EXPRESSION TAG	UNP Q0TRG9
A	322	SER	-	EXPRESSION TAG	UNP Q0TRG9
A	323	HIS	-	EXPRESSION TAG	UNP Q0TRG9
A	324	HIS	-	EXPRESSION TAG	UNP Q0TRG9
A	325	HIS	-	EXPRESSION TAG	UNP Q0TRG9
A	326	HIS	-	EXPRESSION TAG	UNP Q0TRG9
A	327	HIS	-	EXPRESSION TAG	UNP Q0TRG9
A	328	HIS	-	EXPRESSION TAG	UNP Q0TRG9
A	329	TRP	-	EXPRESSION TAG	UNP Q0TRG9
A	330	SER	-	EXPRESSION TAG	UNP Q0TRG9
A	331	HIS	-	EXPRESSION TAG	UNP Q0TRG9
A	332	PRO	-	EXPRESSION TAG	UNP Q0TRG9
A	333	GLN	-	EXPRESSION TAG	UNP Q0TRG9
A	334	PHE	-	EXPRESSION TAG	UNP Q0TRG9
A	335	GLU	-	EXPRESSION TAG	UNP Q0TRG9
A	336	LYS	-	EXPRESSION TAG	UNP Q0TRG9


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	90	Total	O	0	0
			90	90		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Malonyl CoA-acyl carrier protein transacylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	39.70Å 87.08Å 87.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.85 – 2.10 19.85 – 2.10	Depositor EDS
% Data completeness (in resolution range)	97.5 (19.85-2.10) 97.5 (19.85-2.10)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.19 (at 2.09Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.218 , 0.265 0.217 , 0.263	Depositor DCC
R_{free} test set	907 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	34.6	Xtrriage
Anisotropy	0.184	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 29.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.015 for -h,l,k	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2506	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/2448	0.61	1/3290 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	90	LEU	CA-CB-CG	5.48	127.90	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2416	0	2473	32	0
2	A	90	0	0	2	0
All	All	2506	0	2473	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:176:GLU:HG2	2:A:347:HOH:O	1.74	0.87

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:262:MET:O	1:A:267:VAL:HG12	1.85	0.77
1:A:222:LEU:HD21	1:A:241:ILE:HD11	1.69	0.74
1:A:59:THR:HG21	1:A:121:GLN:NE2	2.09	0.68
1:A:159:ASN:HD22	1:A:167:VAL:HB	1.62	0.63
1:A:309:ARG:HA	1:A:312:GLU:HG2	1.84	0.58
1:A:103:ALA:HB2	1:A:222:LEU:HD22	1.85	0.58
1:A:262:MET:HB3	1:A:267:VAL:HG11	1.85	0.57
1:A:59:THR:HG21	1:A:121:GLN:HE21	1.68	0.56
1:A:149:SER:HB2	1:A:155:VAL:HG11	1.88	0.55
1:A:53:GLU:HB2	1:A:55:LEU:HD22	1.87	0.55
1:A:90:LEU:HD22	1:A:255:PHE:HB2	1.89	0.55
1:A:194:VAL:HG12	1:A:196:ALA:H	1.71	0.54
1:A:124:VAL:CG1	1:A:128:ILE:HG12	2.38	0.54
1:A:67:ILE:HG13	1:A:113:VAL:HG21	1.91	0.52
1:A:90:LEU:HD13	1:A:253:VAL:HG11	1.91	0.52
1:A:59:THR:CG2	1:A:121:GLN:NE2	2.72	0.52
1:A:262:MET:C	1:A:267:VAL:HG12	2.31	0.50
1:A:222:LEU:N	1:A:222:LEU:HD23	2.27	0.49
1:A:59:THR:CG2	1:A:121:GLN:HE21	2.26	0.48
1:A:255:PHE:CZ	1:A:259:ILE:HD11	2.50	0.47
1:A:75:THR:HG22	2:A:363:HOH:O	2.14	0.46
1:A:77:LEU:O	1:A:82:VAL:HG13	2.17	0.45
1:A:90:LEU:HD21	1:A:161:ASN:OD1	2.17	0.44
1:A:302:GLU:HA	1:A:302:GLU:OE1	2.18	0.44
1:A:154:ILE:O	1:A:170:GLY:HA3	2.18	0.44
1:A:53:GLU:HB2	1:A:55:LEU:CD2	2.50	0.41
1:A:180:GLU:O	1:A:184:GLU:HG3	2.21	0.41
1:A:99:ILE:HD11	1:A:109:GLY:HA3	2.03	0.41
1:A:112:LEU:HD21	1:A:214:LEU:CD2	2.52	0.40
1:A:21:GLU:O	1:A:25:ASN:ND2	2.33	0.40
1:A:214:LEU:HB3	1:A:242:ILE:HG12	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	314/336 (94%)	308 (98%)	4 (1%)	2 (1%)	25 21

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	3	LYS
1	A	91	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	266/286 (93%)	244 (92%)	22 (8%)	11 7

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	LYS
1	A	40	LEU
1	A	55	LEU
1	A	58	LYS
1	A	77	LEU
1	A	82	VAL
1	A	90	LEU
1	A	128	ILE
1	A	136	ARG
1	A	173	VAL
1	A	175	LEU
1	A	176	GLU
1	A	177	LYS
1	A	217	ILE
1	A	222	LEU
1	A	237	GLU
1	A	244	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	245	LEU
1	A	254	LEU
1	A	260	GLU
1	A	298	LEU
1	A	313	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	121	GLN
1	A	159	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	316/336 (94%)	0.18	8 (2%) 57 62	27, 36, 53, 90	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	313	VAL	5.7
1	A	315	ALA	5.7
1	A	2	ALA	4.7
1	A	316	GLU	4.5
1	A	314	LEU	3.8
1	A	317	ASN	2.8
1	A	195	SER	2.3
1	A	73	ILE	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.