



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 7, 2023 – 06:57 PM EDT

PDB ID : 1PYW
Title : Human class II MHC protein HLA-DR1 bound to a designed peptide related to influenza virus hemagglutinin, FVKQNA(MAA)AL, in complex with staphylococcal enterotoxin C3 variant 3B2 (SEC3-3B2)
Authors : Zavala-Ruiz, Z.; Sundberg, E.J.; Stone, J.D.; DeOliveira, D.B.; Chan, I.C.; Svendsen, J.; Mariuzza, R.A.; Stern, L.J.
Deposited on : 2003-07-09
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	182	 5% 80% 18% ..
2	B	190	 6% 74% 23% .
3	C	10	 70% 20% 10%
4	D	239	 5% 79% 16% . .

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5396 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class II histocompatibility antigen, DR alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	179	1470	952	239	274	5	0	0	0

- Molecule 2 is a protein called HLA class II histocompatibility antigen, DR-1 beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	190	1557	979	279	293	6	37	0	0

- Molecule 3 is a protein called 9-residue influenza virus hemagglutinin related peptide FVKQNA(MAA)AL

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	10	71	47	12	12	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	1	PHE	TYR	engineered mutation	GB 3212739
C	6	ALA	THR	engineered mutation	GB 3212739
C	7	MAA	LEU	engineered mutation	GB 3212739
C	8	ALA	LYS	engineered mutation	GB 3212739

- Molecule 4 is a protein called Enterotoxin type C-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	230	1884	1196	306	372	10	18	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	43	SER	LYS	engineered mutation	UNP P23313
D	45	PHE	LEU	engineered mutation	UNP P23313
D	46	LYS	ALA	engineered mutation	UNP P23313
D	47	TRP	HIS	engineered mutation	UNP P23313


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	135	Total O 135 135	0	0
5	B	113	Total O 113 113	0	0
5	C	12	Total O 12 12	0	0
5	D	154	Total O 154 154	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HLA class II histocompatibility antigen, DR alpha chain




- Molecule 2: HLA class II histocompatibility antigen, DR-1 beta chain




- Molecule 3: 9-residue influenza virus hemagglutinin related peptide FVKQNA(MAA)AL



- Molecule 4: Enterotoxin type C-3





4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	172.51Å 172.51Å 121.65Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 2.10 21.31 – 2.08	Depositor EDS
% Data completeness (in resolution range)	98.3 (20.00-2.10) 97.4 (21.31-2.08)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.35 (at 2.08Å)	Xtrriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.209 , 0.235 0.196 , 0.219	Depositor DCC
R_{free} test set	7813 reflections (9.95%)	wwPDB-VP
Wilson B-factor (Å ²)	27.0	Xtrriage
Anisotropy	0.108	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 48.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.018 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5396	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACE, MAA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.36	0/1515	0.63	0/2065
2	B	0.36	0/1597	0.63	0/2168
3	C	0.49	0/62	0.84	0/81
4	D	0.35	0/1926	0.59	0/2592
All	All	0.36	0/5100	0.62	0/6906

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1470	0	1406	26	0
2	B	1557	0	1488	49	0
3	C	71	0	76	2	0
4	D	1884	0	1811	32	0
5	A	135	0	0	3	0
5	B	113	0	0	0	0
5	C	12	0	0	0	0
5	D	154	0	0	1	0
All	All	5396	0	4781	97	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (97) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:150:ASN:HD22	2:B:154:THR:HG22	1.19	1.05
2:B:105:LYS:HE3	2:B:105:LYS:H	1.12	1.04
4:D:117:LYS:HE2	4:D:119:GLU:HB3	1.53	0.89
2:B:65:LYS:H	2:B:65:LYS:HE3	1.36	0.87
2:B:105:LYS:H	2:B:105:LYS:CE	1.92	0.83
1:A:94:ASN:HD22	1:A:104:VAL:HB	1.47	0.80
1:A:147:LYS:HE3	1:A:149:HIS:HE1	1.47	0.80
4:D:88:ASN:H	4:D:88:ASN:HD22	1.29	0.80
2:B:150:ASN:ND2	2:B:154:THR:HG22	1.96	0.79
4:D:215:MET:O	4:D:218:ASN:HB2	1.82	0.79
4:D:121:ASN:HD21	4:D:153:SER:H	1.31	0.78
2:B:105:LYS:HE3	2:B:105:LYS:N	1.98	0.72
2:B:65:LYS:H	2:B:65:LYS:CE	2.03	0.71
2:B:8:LEU:O	2:B:32:TYR:O	2.12	0.68
2:B:2:ASP:OD1	2:B:4:ARG:HD3	1.94	0.67
4:D:59:LYS:HA	4:D:59:LYS:HE2	1.79	0.65
2:B:152:ASP:OD1	2:B:154:THR:HB	1.97	0.64
1:A:81:PRO:HB3	2:B:5:PRO:HB2	1.82	0.62
5:A:187:HOH:O	4:D:46:LYS:HG2	1.99	0.62
2:B:145:THR:HG22	2:B:158:LEU:H	1.65	0.62
1:A:16:PRO:HD2	2:B:6:ARG:HD3	1.81	0.62
2:B:150:ASN:HD22	2:B:154:THR:CG2	2.06	0.61
4:D:88:ASN:H	4:D:88:ASN:ND2	1.99	0.60
2:B:94:ARG:HH11	2:B:94:ARG:HG3	1.66	0.59
2:B:97:PRO:HB3	2:B:122:PHE:HB3	1.86	0.58
1:A:95:SER:HB2	1:A:96:PRO:HD2	1.85	0.58
1:A:98:GLU:HB2	1:A:101:GLU:HB3	1.87	0.56
2:B:27:LEU:HG	2:B:29:ARG:HD2	1.86	0.56
2:B:65:LYS:HE3	2:B:65:LYS:N	2.15	0.56
2:B:2:ASP:CG	2:B:6:ARG:HH22	2.10	0.55
4:D:83:ASP:OD1	4:D:118:HIS:HD2	1.89	0.55
2:B:114:LEU:HD21	2:B:160:MET:HB3	1.86	0.55
2:B:18:PHE:HB2	2:B:23:ARG:HB3	1.87	0.55
4:D:122:HIS:O	4:D:152:LYS:HE3	2.05	0.55
2:B:94:ARG:HG3	2:B:94:ARG:NH1	2.22	0.54
4:D:37:LYS:HG3	4:D:81:VAL:CG1	2.38	0.54
4:D:2:SER:HB2	4:D:195:THR:H	1.73	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:143:HIS:HD2	2:B:12:LYS:NZ	2.06	0.53
1:A:128:VAL:HG12	1:A:129:THR:N	2.23	0.53
4:D:58:LEU:N	4:D:58:LEU:HD22	2.23	0.53
1:A:38:LYS:O	4:D:212:LYS:HE2	2.09	0.52
1:A:17:ASP:O	1:A:18:GLN:HB2	2.12	0.50
1:A:67:LYS:O	1:A:71:GLU:HG3	2.12	0.50
1:A:182:ALA:HA	2:B:105:LYS:HG2	1.93	0.49
1:A:94:ASN:ND2	1:A:104:VAL:HB	2.21	0.48
2:B:52:GLU:OE1	2:B:55:ARG:HD2	2.13	0.48
4:D:121:ASN:ND2	4:D:153:SER:H	2.05	0.47
2:B:103:PRO:HD3	2:B:188:TRP:CZ3	2.50	0.47
2:B:61:TRP:CZ2	3:C:9:LEU:HD23	2.50	0.47
2:B:116:VAL:HG22	2:B:160:MET:HG3	1.97	0.46
4:D:64:VAL:HG22	4:D:109:THR:HG22	1.97	0.46
2:B:23:ARG:HH11	2:B:23:ARG:HG2	1.80	0.46
1:A:76:ARG:HH22	2:B:57:ASP:CG	2.18	0.46
1:A:76:ARG:NH2	2:B:57:ASP:OD2	2.46	0.45
1:A:12:PHE:C	1:A:12:PHE:CD1	2.90	0.45
2:B:2:ASP:OD1	2:B:6:ARG:NH2	2.49	0.45
1:A:70:LEU:HD13	2:B:9:TRP:HB2	1.98	0.45
4:D:231:GLU:OE1	4:D:233:HIS:HE1	1.98	0.45
4:D:6:PRO:HB3	4:D:197:TRP:CZ2	2.51	0.45
4:D:37:LYS:HG3	4:D:81:VAL:HG11	1.98	0.45
4:D:88:ASN:HD22	4:D:88:ASN:N	2.05	0.44
4:D:226:LYS:HG2	5:D:286:HOH:O	2.16	0.44
2:B:105:LYS:CE	2:B:105:LYS:N	2.69	0.44
1:A:110:ASP:OD1	1:A:146:ARG:HG2	2.18	0.44
4:D:2:SER:CB	4:D:195:THR:H	2.30	0.44
1:A:14:LEU:HD11	2:B:6:ARG:HB3	1.99	0.44
1:A:130:THR:HG23	5:A:292:HOH:O	2.16	0.44
2:B:134:ASN:ND2	2:B:170:VAL:H	2.15	0.44
4:D:129:GLN:NE2	4:D:226:LYS:HB3	2.33	0.43
2:B:74:ALA:O	2:B:78:TYR:HB3	2.19	0.43
4:D:55:ASP:OD1	4:D:60:ASN:HB2	2.19	0.43
3:C:6:ALA:HA	3:C:7:MAA:HM1	1.53	0.43
1:A:140:ARG:HG3	1:A:146:ARG:HG3	2.00	0.43
4:D:121:ASN:HD21	4:D:153:SER:N	2.08	0.42
2:B:23:ARG:HH11	2:B:23:ARG:CG	2.33	0.42
1:A:122:LEU:HD23	1:A:127:PRO:HA	2.01	0.42
2:B:143:VAL:HG13	2:B:160:MET:HB2	2.02	0.42
4:D:231:GLU:HB3	4:D:233:HIS:CE1	2.55	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:105:LEU:HG	1:A:153:PHE:CE1	2.54	0.41
2:B:2:ASP:OD1	2:B:4:ARG:CD	2.64	0.41
4:D:6:PRO:HB3	4:D:197:TRP:CH2	2.55	0.41
4:D:94:TYR:O	4:D:95:PHE:HB3	2.21	0.41
4:D:125:ASN:O	4:D:127:ASN:N	2.53	0.41
2:B:104:SER:HB3	2:B:107:GLN:HE21	1.86	0.41
1:A:49:GLY:HA3	5:A:293:HOH:O	2.20	0.41
2:B:118:SER:HA	2:B:158:LEU:HD23	2.02	0.41
2:B:166:ARG:HB2	2:B:166:ARG:HH11	1.86	0.41
2:B:133:ARG:O	2:B:134:ASN:HB2	2.21	0.41
4:D:1:GLU:HG3	4:D:2:SER:N	2.36	0.41
1:A:82:ILE:HG13	2:B:33:ASN:HB3	2.02	0.40
2:B:87:GLU:HG2	2:B:92:GLN:NE2	2.35	0.40
2:B:104:SER:HB3	2:B:107:GLN:NE2	2.35	0.40
1:A:143:HIS:HD2	2:B:12:LYS:HZ1	1.68	0.40
2:B:166:ARG:CB	2:B:166:ARG:NH1	2.85	0.40
4:D:54:SER:HA	4:D:62:ASP:HA	2.02	0.40
4:D:64:VAL:HG22	4:D:109:THR:CG2	2.51	0.40
4:D:91:VAL:O	4:D:92:ASN:HB2	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	177/182 (97%)	173 (98%)	4 (2%)	0	100	100
2	B	188/190 (99%)	178 (95%)	6 (3%)	4 (2%)	7	3
3	C	7/10 (70%)	7 (100%)	0	0	100	100
4	D	226/239 (95%)	217 (96%)	7 (3%)	2 (1%)	17	12
All	All	598/621 (96%)	575 (96%)	17 (3%)	6 (1%)	15	11

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	108	PRO
2	B	110	GLN
2	B	111	HIS
2	B	32	TYR
4	D	57	LYS
4	D	126	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	163/166 (98%)	161 (99%)	2 (1%)	71	77
2	B	171/171 (100%)	163 (95%)	8 (5%)	26	25
3	C	6/6 (100%)	6 (100%)	0	100	100
4	D	212/220 (96%)	209 (99%)	3 (1%)	67	73
All	All	552/563 (98%)	539 (98%)	13 (2%)	49	53

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	GLU
1	A	101	GLU
2	B	23	ARG
2	B	29	ARG
2	B	53	LEU
2	B	65	LYS
2	B	68	LEU
2	B	105	LYS
2	B	109	LEU
2	B	114	LEU
4	D	56	LYS
4	D	88	ASN
4	D	218	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	78	ASN
1	A	143	HIS
1	A	149	HIS
2	B	64	GLN
2	B	92	GLN
2	B	113	ASN
2	B	134	ASN
2	B	150	ASN
2	B	156	GLN
4	D	60	ASN
4	D	88	ASN
4	D	92	ASN
4	D	118	HIS
4	D	121	ASN
4	D	125	ASN
4	D	127	ASN
4	D	157	GLN
4	D	177	ASN
4	D	218	ASN
4	D	233	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	MAA	C	7	3	4,5,6	2.05	1 (25%)	1,5,7	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MAA	C	7	3	-	1/1/4/6	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	7	MAA	CM-N	-3.90	1.36	1.46

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	7	MAA	CB-CA-N-CM

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	7	MAA	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	179/182 (98%)	0.02	10 (5%) 24 29	15, 26, 61, 84	0
2	B	186/190 (97%)	0.19	12 (6%) 18 23	14, 30, 55, 79	0
3	C	8/10 (80%)	-0.41	0 100 100	18, 21, 24, 37	0
4	D	228/239 (95%)	0.06	11 (4%) 30 36	17, 28, 53, 70	0
All	All	601/621 (96%)	0.08	33 (5%) 25 31	14, 28, 57, 84	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	182	ALA	10.1
4	D	125	ASN	5.6
1	A	181	ASP	5.5
2	B	1	GLY	5.4
4	D	58	LEU	5.3
2	B	108	PRO	4.6
4	D	238	ASN	4.3
4	D	239	GLY	4.2
4	D	106	GLY	4.2
1	A	157	THR	4.1
1	A	158	GLU	3.8
4	D	126	GLY	3.6
4	D	124	ASP	3.6
2	B	106	THR	3.6
2	B	139	LYS	3.5
4	D	1	GLU	3.5
1	A	180	PHE	3.4
1	A	130	THR	3.1
2	B	136	GLN	2.9
2	B	181	THR	2.8
2	B	164	VAL	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	101	GLU	2.6
2	B	166	ARG	2.6
1	A	131	GLY	2.6
4	D	177	ASN	2.6
4	D	127	ASN	2.5
4	D	7	MET	2.5
2	B	19	ASN	2.4
2	B	145	THR	2.4
1	A	100	ARG	2.3
2	B	162	GLU	2.1
2	B	135	GLY	2.1
1	A	129	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MAA	C	7	6/7	0.97	0.09	17,19,19,20	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.