

Full wwPDB X-ray Structure Validation Report (i)

Aug 21, 2023 – 05:14 PM EDT

PDB ID	:	2PYD
Title	:	The crystal structure of Glycogen phosphorylase in complex with glucose at
		100 K
Authors	:	Alexacou, K.M.; Tiraidis, C.; Zographos, S.E.; Chrysina, E.D.; Hayes, J.;
		Oikonomakos, N.G.
Deposited on	:	2007-05-16
Resolution	:	1.93 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 1.93 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
Clashscore	141614	1023 (1.94-1.94)
Ramachandran outliers	138981	1007 (1.94-1.94)
Sidechain outliers	138945	1007 (1.94-1.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain		
1	А	843	89%	6%	·



2PYD

2 Entry composition (i)


There are 4 unique types of molecules in this entry. The entry contains 7653 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Glycogen phosphorylase, muscle form.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	А	808	Total 6609	C 4207	N 1171	O 1200	Р 1	S 30	0	4	0


• Molecule 2 is alpha-D-glucopyranose (three-letter code: GLC) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	Total C O 12 6 6	0	0

• Molecule 3 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C_2H_6OS).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	$\begin{array}{cccc} \text{Total} & \text{C} & \text{O} & \text{S} \\ 4 & 2 & 1 & 1 \end{array}$	0	0
3	А	1	$\begin{array}{cccc} \text{Total} & \text{C} & \text{O} & \text{S} \\ 4 & 2 & 1 & 1 \end{array}$	0	0
3	А	1	$\begin{array}{cccc} \text{Total} & \text{C} & \text{O} & \text{S} \\ 4 & 2 & 1 & 1 \end{array}$	0	0
3	А	1	$\begin{array}{cccc} \text{Total} & \text{C} & \text{O} & \text{S} \\ 4 & 2 & 1 & 1 \end{array}$	0	0
3	А	1	$\begin{array}{cccc} \text{Total} & \text{C} & \text{O} & \text{S} \\ 4 & 2 & 1 & 1 \end{array}$	0	0
3	А	1	$\begin{array}{cccc} \text{Total} & \text{C} & \text{O} & \text{S} \\ 4 & 2 & 1 & 1 \end{array}$	0	0
3	А	1	$\begin{array}{cccc} \text{Total} & \text{C} & \text{O} & \text{S} \\ 4 & 2 & 1 & 1 \end{array}$	0	0
3	А	1	$\begin{array}{cccc} \text{Total} & \text{C} & \text{O} & \text{S} \\ 4 & 2 & 1 & 1 \end{array}$	0	0
3	А	1	$\begin{array}{cccc} \text{Total} & \text{C} & \text{O} & \text{S} \\ 4 & 2 & 1 & 1 \end{array}$	0	0
3	А	1	$\begin{array}{cccc} \text{Total} & \text{C} & \text{O} & \text{S} \\ 4 & 2 & 1 & 1 \end{array}$	0	0
3	А	1	$\begin{array}{cccc} \text{Total} & \text{C} & \text{O} & \text{S} \\ 4 & 2 & 1 & 1 \end{array}$	0	0
3	А	1	$\begin{array}{cccc} \text{Total} & \text{C} & \text{O} & \text{S} \\ 4 & 2 & 1 & 1 \end{array}$	0	0
3	А	1	$\begin{array}{cccc} \text{Total} & \text{C} & \text{O} & \text{S} \\ 4 & 2 & 1 & 1 \end{array}$	0	0
3	А	1	$\begin{array}{cccc} \text{Total} & \text{C} & \text{O} & \text{S} \\ 4 & 2 & 1 & 1 \end{array}$	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	$\begin{array}{cccc} \text{Total} & \text{C} & \text{O} & \text{S} \\ 4 & 2 & 1 & 1 \end{array}$	0	0
			$\begin{array}{c ccccccccccccccccccccccccccccccccccc$		
3	А	1	$4 \ 2 \ 1 \ 1$	0	0
3	А	1	Total C O S	0	0
			$\begin{array}{c ccccccccccccccccccccccccccccccccccc$		
3	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} & \text{S} \\ 4 & 2 & 1 & 1 \end{array}$	0	0

• Molecule 4 is water.

Μ	ol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	1	А	960	Total 960	O 960	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: Glycogen phosphorylase, muscle form





4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	125.73Å 125.73Å 114.93Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 - 1.93	Depositor
% Data completeness (in resolution range)	99.6 (30.00-1.93)	Depositor
R _{merge}	(Not available)	Depositor
R _{sym}	0.07	Depositor
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.190 , 0.235	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	7653	wwPDB-VP
Average B, all atoms $(Å^2)$	20.0	wwPDB-VP



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: LLP, DMS, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bond angles		
	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.37	1/6754~(0.0%)	0.55	3/9134~(0.0%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	А	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	Observed(Å)	Ideal(Å)
1	А	250	ASN	C-O	5.44	1.33	1.23

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
1	А	250	ASN	CA-C-N	15.66	151.65	117.20
1	А	250	ASN	O-C-N	-12.72	102.34	122.70
1	А	250	ASN	CA-C-O	-12.32	94.23	120.10

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	А	250	ASN	Mainchain



5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	6609	0	6546	43	0
2	А	12	0	12	0	0
3	А	72	0	108	3	0
4	А	960	0	0	7	0
All	All	7653	0	6666	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

	A	Interatomic	Clash
Atom-1	Atom-2	distance (\AA)	overlap (Å)
1:A:16:ARG:HH22	1:A:497:PRO:HB3	1.32	0.94
1:A:16:ARG:HH22	1:A:497:PRO:CB	1.97	0.78
1:A:455:VAL:H	1:A:459:HIS:HD2	1.35	0.72
1:A:16:ARG:NH2	1:A:497:PRO:HB3	2.04	0.70
1:A:93[B]:ARG:NH2	4:A:1320:HOH:O	2.26	0.69
1:A:34:HIS:HD2	1:A:38:THR:OG1	1.80	0.64
1:A:16:ARG:NH2	1:A:497:PRO:CB	2.64	0.60
1:A:224:MET:HE1	4:A:1832:HOH:O	2.01	0.60
1:A:283:ASP:O	1:A:571[B]:HIS:CE1	2.56	0.59
1:A:159:ILE:CG1	1:A:299:VAL:HG22	2.34	0.57
1:A:159:ILE:HG13	1:A:299:VAL:CG2	2.35	0.55
1:A:450:HIS:HD2	4:A:1072:HOH:O	1.88	0.55
1:A:34:HIS:HE1	1:A:61:ASP:OD1	1.89	0.55
1:A:583:VAL:HG11	1:A:642:VAL:HG21	1.90	0.53
1:A:680:LLP:O3	1:A:680:LLP:NZ	2.41	0.53
1:A:250:ASN:O	1:A:251:ASP:OD1	2.26	0.53
1:A:184:ARG:HD2	1:A:185:TYR:CE2	2.44	0.53
1:A:159:ILE:CG1	1:A:299:VAL:CG2	2.87	0.52
1:A:224:MET:CE	4:A:1832:HOH:O	2.58	0.51
1:A:122:LEU:HA	1:A:125:ILE:HD12	1.92	0.51
1:A:235:ASN:H	1:A:235:ASN:HD22	1.57	0.51
1:A:11:LYS:HE3	1:A:19:ALA:HB2	1.94	0.50
1:A:224:MET:HE3	1:A:247:LYS:HD2	1.93	0.49

Continued on next page...



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:786:ARG:NH2	4:A:1222:HOH:O	2.45	0.49
1:A:159:ILE:HG12	1:A:299:VAL:HG22	1.95	0.48
1:A:85:LEU:HD13	1:A:335:ILE:HG23	1.96	0.48
1:A:159:ILE:HG13	1:A:299:VAL:HG22	1.96	0.47
1:A:283:ASP:O	1:A:571[B]:HIS:HE1	1.96	0.47
1:A:648:TYR:HA	1:A:652:LEU:HD23	1.99	0.45
1:A:566:GLN:HE22	1:A:576:GLN:HA	1.82	0.45
1:A:450:HIS:HE1	4:A:1062:HOH:O	2.00	0.44
1:A:431:VAL:HG11	1:A:437:LYS:HE2	1.99	0.44
1:A:470:ASP:HB2	3:A:983:DMS:H23	2.01	0.43
1:A:466:THR:OG1	1:A:467:ILE:HD12	2.19	0.43
1:A:150:LEU:HD12	1:A:817:ILE:HG22	2.01	0.43
1:A:693:ASP:O	1:A:696:ASN:HB2	2.19	0.42
1:A:726:TYR:OH	1:A:774:PHE:HB2	2.19	0.42
1:A:467:ILE:HD12	1:A:467:ILE:H	1.84	0.42
1:A:16:ARG:HD3	4:A:1754:HOH:O	2.19	0.42
1:A:270:ASN:ND2	3:A:978:DMS:O	2.47	0.41
1:A:49:ARG:HA	1:A:125:ILE:HG21	2.02	0.41
1:A:786:ARG:CZ	3:A:975:DMS:O	2.69	0.41
1:A:275:ILE:O	1:A:295:GLN:HG2	2.20	0.41

Continued from previous page...

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured Allowed		Outliers	Percentiles	
1	А	805/843~(96%)	783~(97%)	21 (3%)	1 (0%)	51 43	

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	250	ASN



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	А	703/731~(96%)	694 (99%)	9(1%)	69 62	

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	10	ARG
1	А	11	LYS
1	А	90	TYR
1	А	128	ASP
1	А	235	ASN
1	А	250	ASN
1	А	576	GLN
1	А	579	ASN
1	А	613	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	34	HIS
1	А	57	HIS
1	А	97	ASN
1	А	106	ASN
1	А	235	ASN
1	А	450	HIS
1	А	453	ASN
1	А	459	HIS
1	А	477	HIS
1	А	481	ASN
1	А	484	ASN
1	А	566	GLN
1	А	576	GLN
1	А	579	ASN



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Type Chain		Chain Res Link		Bo	Bond lengths			Bond angles		
Moi Type Chai	Ullaili	anam Res Lin		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2		
1	LLP	А	680	1	23,24,25	1.71	3 (13%)	25,32,34	1.26	5 (20%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	А	680	1	-	2/16/17/19	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	А	680	LLP	O3-C3	-5.59	1.24	1.37
1	А	680	LLP	C4-C4'	2.75	1.51	1.46
1	А	680	LLP	C2-N1	2.47	1.38	1.33

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
1	А	680	LLP	C4-C4'-NZ	-2.82	111.36	124.31
1	А	680	LLP	CE-NZ-C4'	-2.54	111.11	118.90
1	А	680	LLP	OP4-P-OP1	-2.33	99.95	106.47
1	А	680	LLP	C5-C6-N1	-2.17	120.20	123.82
1	А	680	LLP	OP3-P-OP2	2.10	115.68	107.64



There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	А	680	LLP	C6-C5-C5'-OP4
1	А	680	LLP	C4-C5-C5'-OP4

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	А	680	LLP	1	0

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

19 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Turne	Chain	Res	Link	Bo	ond leng	ths	B	ond ang	les
	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
2	GLC	А	998	-	$12,\!12,\!12$	0.54	0	17,17,17	0.45	0
3	DMS	А	978	-	$3,\!3,\!3$	2.61	1 (33%)	3, 3, 3	0.48	0
3	DMS	А	981	-	$3,\!3,\!3$	2.66	1 (33%)	3,3,3	0.48	0
3	DMS	А	971	-	$3,\!3,\!3$	2.65	1 (33%)	3,3,3	0.50	0
3	DMS	А	980	-	3,3,3	2.66	1 (33%)	3,3,3	0.54	0
3	DMS	А	975	-	3,3,3	2.54	1 (33%)	3,3,3	0.38	0
3	DMS	А	974	-	3,3,3	2.64	1 (33%)	3,3,3	0.45	0
3	DMS	А	986	-	3,3,3	2.68	1 (33%)	3,3,3	0.54	0
3	DMS	А	987	-	3,3,3	2.62	1 (33%)	3,3,3	0.62	0
3	DMS	А	982	-	3,3,3	2.68	1 (33%)	3,3,3	0.61	0



Mol	Turne	Chain	Res	Link	Bo	ond leng	ths	В	ond ang	les
	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
3	DMS	А	976	-	$3,\!3,\!3$	2.63	1 (33%)	$3,\!3,\!3$	0.49	0
3	DMS	А	972	-	3,3,3	2.59	1 (33%)	3,3,3	0.53	0
3	DMS	А	977	-	3,3,3	2.67	1 (33%)	3,3,3	0.50	0
3	DMS	А	979	-	$3,\!3,\!3$	2.57	1 (33%)	3,3,3	0.42	0
3	DMS	А	984	-	$3,\!3,\!3$	2.69	1 (33%)	$3,\!3,\!3$	0.51	0
3	DMS	А	973	-	3,3,3	2.64	1 (33%)	3,3,3	0.52	0
3	DMS	А	983	-	3,3,3	2.71	1 (33%)	3,3,3	0.45	0
3	DMS	А	985	-	3,3,3	2.66	1 (33%)	3,3,3	0.50	0
3	DMS	А	970	-	3,3,3	2.64	1 (33%)	3,3,3	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	А	998	-	-	0/2/22/22	0/1/1/1

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	А	983	DMS	O-S	4.55	1.81	1.50
3	А	984	DMS	O-S	4.52	1.80	1.50
3	А	986	DMS	O-S	4.50	1.80	1.50
3	А	982	DMS	O-S	4.49	1.80	1.50
3	А	977	DMS	O-S	4.48	1.80	1.50
3	А	985	DMS	O-S	4.47	1.80	1.50
3	А	980	DMS	O-S	4.47	1.80	1.50
3	А	981	DMS	O-S	4.46	1.80	1.50
3	А	971	DMS	O-S	4.46	1.80	1.50
3	А	974	DMS	O-S	4.44	1.80	1.50
3	А	970	DMS	O-S	4.43	1.80	1.50
3	А	973	DMS	O-S	4.43	1.80	1.50
3	А	976	DMS	O-S	4.42	1.80	1.50
3	А	978	DMS	O-S	4.38	1.79	1.50
3	А	987	DMS	O-S	4.38	1.79	1.50
3	А	972	DMS	O-S	4.32	1.79	1.50
3	А	979	DMS	O-S	4.31	1.79	1.50
3	А	975	DMS	O-S	4.27	1.79	1.50



There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	А	978	DMS	1	0
3	А	975	DMS	1	0
3	А	983	DMS	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

