



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 18, 2025 – 12:06 pm GMT

PDB ID : 8Q3D
Title : HsNMT1 in complex with both MyrCoA and GNCFSKPR(NH2) inhibitor peptide
Authors : Dian, C.; Giglione, C.; Meinnel, T.
Deposited on : 2023-08-04
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.41


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1881 (2.16-2.16)
Clashscore	180529	2047 (2.16-2.16)
Ramachandran outliers	177936	2027 (2.16-2.16)
Sidechain outliers	177891	2026 (2.16-2.16)
RSRZ outliers	164620	1882 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	401	 91% 7%
2	B	401	 94%
3	C	9	 67% 33%
3	D	9	 78% 22%

2 Entry composition i

There are 8 unique types of molecules in this entry. The entry contains 7237 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycylpeptide N-tetradecanoyltransferase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	393	3197	2070	540	570	17	0	4	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	96	GLY	-	expression tag	UNP P30419
A	97	SER	-	expression tag	UNP P30419
A	98	GLU	-	expression tag	UNP P30419

- Molecule 2 is a protein called Glycylpeptide N-tetradecanoyltransferase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	392	3176	2060	530	569	17	0	4	0


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	96	GLY	-	expression tag	UNP P30419
B	97	SER	-	expression tag	UNP P30419
B	98	GLU	-	expression tag	UNP P30419
B	195	GLN	GLU	conflict	UNP P30419

- Molecule 3 is a protein called GNCFSKPR(NH2) inhibitor peptide.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	9	63	38	14	10	1	0	0	1
3	D	9	63	38	14	10	1	0	0	1

- Molecule 4 is COENZYME A (three-letter code: COA) (formula: $C_{21}H_{36}N_7O_{16}P_3S$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	P			S
4	A	1	48	21	7	16	3	1	0	0
4	B	1	48	21	7	16	3	1	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Cl 1 1	0	0
6	B	2	Total Cl 2 2	0	0

- Molecule 7 is MYRISTIC ACID (three-letter code: MYR) (formula: C₁₄H₂₈O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	C	1	Total C O 15 14 1	0	0
7	D	1	Total C O 15 14 1	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	271	Total O 271 271	0	2
8	B	301	Total O 301 301	0	0
8	C	6	Total O 6 6	0	0
8	D	7	Total O 7 7	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glycylpeptide N-tetradecanoyltransferase 1




- Molecule 2: Glycylpeptide N-tetradecanoyltransferase 1



- Molecule 3: GNCFSKPR(NH2) inhibitor peptide



- Molecule 3: GNCFSKPR(NH2) inhibitor peptide



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	79.80Å 178.34Å 58.26Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.67 – 2.15 47.67 – 2.15	Depositor EDS
% Data completeness (in resolution range)	99.5 (47.67-2.15) 99.5 (47.67-2.15)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	10.10 (at 2.16Å)	Xtrriage
Refinement program	PHENIX 1.17_3644	Depositor
R, R_{free}	0.142 , 0.187 0.142 , 0.187	Depositor DCC
R_{free} test set	2526 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	18.8	Xtrriage
Anisotropy	0.459	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 42.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7237	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 25.13 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.3575e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NH2, MYR, GOL, CL, COA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.36	0/3288	0.52	0/4474
2	B	0.39	0/3269	0.54	0/4449
3	C	1.98	3/63 (4.8%)	0.68	0/82
3	D	2.00	3/63 (4.8%)	0.63	0/82
All	All	0.46	6/6683 (0.1%)	0.53	0/9087

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	9	ARG	NE-CZ	10.31	1.46	1.33
3	C	9	ARG	NE-CZ	10.28	1.46	1.33
3	C	9	ARG	CZ-NH1	9.91	1.46	1.33
3	D	9	ARG	CZ-NH1	9.75	1.45	1.33
3	C	9	ARG	CZ-NH2	-5.31	1.26	1.33
3	D	9	ARG	CZ-NH2	-5.22	1.26	1.33

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3197	0	3149	19	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	3176	0	3121	9	0
3	C	63	0	60	1	0
3	D	63	0	60	0	0
4	A	48	0	32	3	0
4	B	48	0	32	1	0
5	A	12	0	16	1	0
5	B	12	0	16	2	0
6	A	1	0	0	1	0
6	B	2	0	0	0	0
7	C	15	0	27	0	0
7	D	15	0	27	0	0
8	A	271	0	0	1	0
8	B	301	0	0	1	0
8	C	6	0	0	0	0
8	D	7	0	0	0	0
All	All	7237	0	6540	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:501:COA:O4B	4:A:501:COA:C1B	1.66	1.21
4:B:501:COA:O4B	4:B:501:COA:C1B	1.65	1.17
2:B:145:ILE:HD12	2:B:272:HIS:HB3	1.78	0.64
2:B:389:ASN:HB3	2:B:428[A]:GLN:HE22	1.63	0.63
4:A:501:COA:HS1	3:C:2:GLY:N	2.00	0.60
2:B:234:HIS:ND1	8:B:602:HOH:O	2.31	0.56
1:A:258:ARG:NH1	8:A:608:HOH:O	2.40	0.55
2:B:244:GLU:OE1	5:B:503:GOL:H12	2.10	0.52
1:A:410:HIS:CD2	1:A:413:HIS:H	2.27	0.51
2:B:374:TRP:CZ2	5:B:503:GOL:H11	2.46	0.51
1:A:317:ASN:OD1	1:A:317:ASN:N	2.43	0.49
1:A:244:GLU:OE2	5:A:503:GOL:O3	2.29	0.48
2:B:458:ASN:HA	2:B:461:PHE:CE2	2.48	0.48
1:A:106:ALA:HB2	1:A:115:ARG:HD2	1.95	0.47
4:A:501:COA:S1P	4:A:501:COA:N6A	2.81	0.46
1:A:107:LYS:HE2	1:A:107:LYS:HB3	1.83	0.46
1:A:458:ASN:HA	1:A:461:PHE:CE2	2.50	0.46
1:A:206:TRP:CZ3	6:A:504:CL:CL	3.07	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:475:GLN:HB2	2:B:477:TYR:CE2	2.52	0.45
1:A:104:GLY:O	1:A:115:ARG:NE	2.51	0.45
1:A:107:LYS:NZ	1:A:184:ASP:OD1	2.36	0.44
1:A:475:GLN:HB2	1:A:477:TYR:CE2	2.52	0.44
1:A:138:VAL:HG22	1:A:482:LYS:HG3	2.00	0.44
1:A:410:HIS:HD2	1:A:412:THR:H	1.67	0.43
1:A:417[B]:LYS:HD2	1:A:447:PHE:CD1	2.55	0.42
1:A:435:MET:HG3	1:A:461:PHE:CE2	2.55	0.42
1:A:145:ILE:HD12	1:A:272:HIS:HB3	2.02	0.41
1:A:144:ASN:N	1:A:144:ASN:OD1	2.54	0.41
1:A:410:HIS:CD2	1:A:412:THR:H	2.39	0.40
2:B:128:LEU:HD23	2:B:128:LEU:HA	1.88	0.40
1:A:253:LYS:HA	1:A:253:LYS:HD3	1.84	0.40
2:B:458:ASN:O	2:B:462:LEU:HG	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	395/401 (98%)	387 (98%)	8 (2%)	0	100	100
2	B	394/401 (98%)	383 (97%)	11 (3%)	0	100	100
3	C	7/9 (78%)	7 (100%)	0	0	100	100
3	D	7/9 (78%)	7 (100%)	0	0	100	100
All	All	803/820 (98%)	784 (98%)	19 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	348/362 (96%)	343 (99%)	5 (1%)	62	68
2	B	345/362 (95%)	340 (99%)	5 (1%)	62	68
3	C	7/7 (100%)	6 (86%)	1 (14%)	2	1
3	D	7/7 (100%)	6 (86%)	1 (14%)	2	1
All	All	707/738 (96%)	695 (98%)	12 (2%)	58	61

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	146	ARG
1	A	184	ASP
1	A	322	ARG
1	A	355	ARG
1	A	455	LEU
2	B	184	ASP
2	B	206	TRP
2	B	428[A]	GLN
2	B	428[B]	GLN
2	B	455	LEU
3	C	4	CYS
3	D	4	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	351	GLN
1	A	410	HIS
2	B	144	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 3 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	GOL	B	503	-	5,5,5	0.51	0	5,5,5	0.60	0
4	COA	A	501	-	41,50,50	4.25	12 (29%)	52,75,75	1.75	6 (11%)
7	MYR	C	101	3	14,14,15	0.53	0	13,13,15	0.66	0
5	GOL	B	502	-	5,5,5	0.56	0	5,5,5	0.27	0
7	MYR	D	101	3	14,14,15	0.53	0	13,13,15	0.81	0
5	GOL	A	503	-	5,5,5	0.52	0	5,5,5	0.53	0
4	COA	B	501	-	41,50,50	4.18	12 (29%)	52,75,75	1.73	5 (9%)
5	GOL	A	502	-	5,5,5	0.53	0	5,5,5	0.34	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	B	503	-	-	4/4/4/4	-
4	COA	A	501	-	-	8/44/64/64	0/3/3/3
7	MYR	C	101	3	-	0/11/12/13	-
5	GOL	B	502	-	-	0/4/4/4	-
7	MYR	D	101	3	-	0/11/12/13	-
5	GOL	A	503	-	-	4/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	COA	B	501	-	-	4/44/64/64	0/3/3/3
5	GOL	A	502	-	-	0/4/4/4	-

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	501	COA	O4B-C1B	18.12	1.66	1.41
4	B	501	COA	O4B-C1B	17.67	1.65	1.41
4	A	501	COA	C2B-C1B	-13.50	1.33	1.53
4	B	501	COA	C2B-C1B	-13.13	1.33	1.53
4	B	501	COA	C5P-N4P	6.89	1.49	1.33
4	A	501	COA	C9P-N8P	6.84	1.48	1.33
4	B	501	COA	C9P-N8P	6.54	1.47	1.33
4	A	501	COA	C5P-N4P	6.48	1.48	1.33
4	B	501	COA	O4B-C4B	-6.13	1.31	1.45
4	A	501	COA	O4B-C4B	-6.06	1.31	1.45
4	A	501	COA	C6A-N6A	4.56	1.50	1.34
4	B	501	COA	C6A-N6A	4.54	1.50	1.34
4	B	501	COA	P3B-O3B	4.50	1.67	1.59
4	A	501	COA	P3B-O3B	4.29	1.67	1.59
4	A	501	COA	C2A-N3A	3.14	1.37	1.32
4	B	501	COA	C2A-N3A	2.89	1.36	1.32
4	A	501	COA	O3B-C3B	-2.76	1.34	1.44
4	B	501	COA	O3B-C3B	-2.45	1.35	1.44
4	B	501	COA	C3B-C4B	2.38	1.59	1.52
4	A	501	COA	P2A-O6A	2.30	1.68	1.59
4	A	501	COA	C3B-C4B	2.22	1.58	1.52
4	B	501	COA	P2A-O6A	2.19	1.68	1.59
4	B	501	COA	C2A-N1A	2.09	1.37	1.33
4	A	501	COA	C2A-N1A	2.07	1.37	1.33

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	501	COA	C5A-C6A-N6A	7.39	131.58	120.35
4	A	501	COA	C5A-C6A-N6A	7.38	131.57	120.35
4	B	501	COA	N3A-C2A-N1A	-5.49	120.10	128.68
4	A	501	COA	N3A-C2A-N1A	-5.45	120.16	128.68
4	A	501	COA	N6A-C6A-N1A	-5.10	107.98	118.57
4	B	501	COA	N6A-C6A-N1A	-5.01	108.18	118.57
4	B	501	COA	P2A-O3A-P1A	-4.12	118.68	132.83
4	A	501	COA	P2A-O3A-P1A	-4.07	118.86	132.83

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	501	COA	CDP-CBP-CAP	2.50	113.16	108.82
4	A	501	COA	C6P-C7P-N8P	-2.44	106.97	111.90
4	B	501	COA	O6A-CCP-CBP	-2.41	106.67	110.55

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	501	COA	C3B-O3B-P3B-O8A
4	A	501	COA	C5B-O5B-P1A-O1A
4	A	501	COA	C5B-O5B-P1A-O2A
4	A	501	COA	C5B-O5B-P1A-O3A
4	B	501	COA	C3B-O3B-P3B-O8A
5	A	503	GOL	C1-C2-C3-O3
5	B	503	GOL	O1-C1-C2-O2
5	B	503	GOL	O1-C1-C2-C3
5	A	503	GOL	O1-C1-C2-C3
5	B	503	GOL	C1-C2-C3-O3
5	A	503	GOL	O1-C1-C2-O2
5	A	503	GOL	O2-C2-C3-O3
4	A	501	COA	C3B-C4B-C5B-O5B
5	B	503	GOL	O2-C2-C3-O3
4	A	501	COA	P1A-O3A-P2A-O4A
4	A	501	COA	O4B-C4B-C5B-O5B
4	B	501	COA	C3B-O3B-P3B-O7A
4	B	501	COA	C5B-O5B-P1A-O3A
4	B	501	COA	C6P-C7P-N8P-C9P
4	A	501	COA	P1A-O3A-P2A-O5A


There are no ring outliers.



4 monomers are involved in 7 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	503	GOL	2	0
4	A	501	COA	3	0
5	A	503	GOL	1	0
4	B	501	COA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	393/401 (98%)	-0.63	4 (1%) 79 82	9, 19, 45, 63	4 (1%)
2	B	392/401 (97%)	-0.71	4 (1%) 79 82	7, 17, 46, 69	4 (1%)
3	C	8/9 (88%)	-0.81	0 100 100	15, 21, 25, 41	0
3	D	8/9 (88%)	-0.75	0 100 100	14, 17, 23, 32	0
All	All	801/820 (97%)	-0.67	8 (0%) 79 82	7, 18, 45, 69	8 (0%)

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	105	PRO	3.4
1	A	104	GLY	2.9
2	B	316	ARG	2.9
1	A	315	SER	2.8
2	B	411	PRO	2.6
2	B	317	ASN	2.3
1	A	106	ALA	2.2
2	B	108	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	COA	A	501	48/48	0.91	0.09	20,37,49,56	0
5	GOL	A	503	6/6	0.92	0.10	26,33,39,39	0
5	GOL	B	503	6/6	0.92	0.10	24,34,35,35	0
4	COA	B	501	48/48	0.95	0.07	15,28,40,41	0
7	MYR	D	101	15/16	0.96	0.05	6,8,14,14	0
7	MYR	C	101	15/16	0.97	0.05	6,12,17,17	0
5	GOL	A	502	6/6	0.97	0.05	14,22,23,25	0
6	CL	B	505	1/1	0.98	0.06	31,31,31,31	0
5	GOL	B	502	6/6	0.98	0.04	12,13,20,22	0
6	CL	A	504	1/1	0.98	0.05	31,31,31,31	0
6	CL	B	504	1/1	0.99	0.03	23,23,23,23	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around COA A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around COA B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MYR D 101:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around MYR C 101:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.