



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 13, 2024 – 10:12 pm GMT

PDB ID : 6Q78
Title : The structure of GH26C from Muricauda sp. MAR_2010_75
Authors : Robb, C.S.; Hehemann, J.H.
Deposited on : 2018-12-13
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	418	 3% 86% 11%
1	B	418	 3% 85% 5% 11%

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 7047 atoms, of which 150 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GH26C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	374	3173	1947	150	478	584	14	0	0	0
1	B	374	3016	1942	479	581	14	1	0	0	0

- Molecule 2 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: $C_4H_{12}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	8	4	1	3	0	0
2	B	1	8	4	1	3	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	445	Total 445	O 445	0	0
3	B	397	Total 397	O 397	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	94.12Å 60.36Å 149.00Å 90.00° 95.92° 90.00°	Depositor
Resolution (Å)	48.84 – 1.50 48.84 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.5 (48.84-1.50) 99.5 (48.84-1.50)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.12 (at 1.50Å)	Xtrriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.164 , 0.191 0.165 , 0.192	Depositor DCC
R_{free} test set	6753 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	13.9	Xtrriage
Anisotropy	0.308	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 45.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7047	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TRS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.57	0/3110	0.69	0/4221
1	B	0.58	1/3103 (0.0%)	0.69	2/4212 (0.0%)
All	All	0.58	1/6213 (0.0%)	0.69	2/8433 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	379	TYR	CE1-CZ	-6.00	1.30	1.38

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	79	LEU	CA-CB-CG	7.69	132.98	115.30
1	B	108	MET	CG-SD-CE	-5.62	91.21	100.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3023	150	2799	11	0
1	B	3016	0	2788	12	0
2	A	8	0	12	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	8	0	12	0	0
3	A	445	0	0	6	1
3	B	397	0	0	5	4
All	All	6897	150	5611	21	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:405:VAL:O	3:B:601:HOH:O	2.04	0.74
1:A:405:VAL:O	3:A:601:HOH:O	2.06	0.73
1:B:367:ASP:OD2	3:B:602:HOH:O	2.08	0.71
1:A:410:ASP:OD2	3:A:601:HOH:O	2.11	0.69
1:B:76:LEU:HA	1:B:79:LEU:HD13	1.76	0.66
1:A:304:ASP:OD1	3:A:602:HOH:O	2.16	0.61
1:A:142:GLN:HG2	3:A:994:HOH:O	2.08	0.54
1:B:410:ASP:OD2	3:B:601:HOH:O	2.19	0.53
1:A:419:PRO:N	3:A:613:HOH:O	2.46	0.47
1:A:310:GLN:O	1:A:314:GLU:HG3	2.15	0.47
1:A:279:PHE:HE2	1:B:279:PHE:CD1	2.35	0.45
1:B:361:GLU:OE1	3:B:603:HOH:O	2.21	0.44
1:A:279:PHE:CE2	1:B:279:PHE:CD1	3.07	0.43
1:B:139:GLN:NE2	3:B:618:HOH:O	2.52	0.42
1:B:383:TYR:CZ	1:B:390:LEU:HD13	2.54	0.42
1:A:211:LYS:NZ	1:A:263:LEU:O	2.48	0.42
1:B:273:SER:HA	1:B:274:PRO:HD2	1.91	0.42
1:A:177:MET:HB2	1:A:181:GLN:OE1	2.21	0.41
1:A:217:ASN:ND2	3:A:615:HOH:O	2.48	0.41
1:B:54:GLU:CG	1:B:420:ASN:HD21	2.33	0.41
1:B:200:LYS:HE3	1:B:258:TYR:CE1	2.55	0.41

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:743:HOH:O	3:B:959:HOH:O[4_645]	1.92	0.28
3:B:937:HOH:O	3:B:959:HOH:O[4_645]	2.07	0.13
3:B:952:HOH:O	3:B:952:HOH:O[2_655]	2.09	0.11

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:986:HOH:O	3:A:999:HOH:O[4_646]	2.10	0.10
3:B:628:HOH:O	3:B:959:HOH:O[4_645]	2.12	0.08

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	372/418 (89%)	363 (98%)	8 (2%)	1 (0%)	41	18
1	B	372/418 (89%)	365 (98%)	6 (2%)	1 (0%)	41	18
All	All	744/836 (89%)	728 (98%)	14 (2%)	2 (0%)	41	18

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	275	ASP
1	A	275	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	325/371 (88%)	324 (100%)	1 (0%)	92	85
1	B	323/371 (87%)	321 (99%)	2 (1%)	86	74
All	All	648/742 (87%)	645 (100%)	3 (0%)	88	78

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	252	TRP
1	B	79	LEU
1	B	252	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	TRS	B	501	-	7,7,7	0.30	0	9,9,9	0.54	0
2	TRS	A	501	-	7,7,7	0.36	0	9,9,9	0.61	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TRS	B	501	-	-	0/9/9/9	-
2	TRS	A	501	-	-	0/9/9/9	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	374/418 (89%)	-0.00	5 (1%) 77 81	6, 13, 26, 54	14 (3%)
1	B	374/418 (89%)	0.15	13 (3%) 44 48	8, 16, 32, 60	8 (2%)
All	All	748/836 (89%)	0.08	18 (2%) 59 63	6, 15, 30, 60	22 (2%)

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	345	LEU	9.7
1	B	348	ASN	4.6
1	B	350	ALA	4.4
1	B	346	SER	3.8
1	B	419	PRO	3.7
1	B	344	THR	3.6
1	B	420	ASN	3.3
1	A	49	LEU	3.3
1	B	349	GLY	3.2
1	A	420	ASN	3.1
1	B	391	PRO	3.0
1	B	49	LEU	2.9
1	A	419	PRO	2.5
1	B	347	GLU	2.3
1	A	345	LEU	2.2
1	A	348	ASN	2.2
1	B	379	TYR	2.1
1	B	47	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	TRS	B	501	8/8	0.90	0.12	14,19,20,23	0
2	TRS	A	501	8/8	0.92	0.12	13,18,20,22	0

6.5 Other polymers [i](#)

There are no such residues in this entry.