



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 21, 2018 – 07:05 pm GMT

PDB ID : 1QBV  
Title : CRYSTAL STRUCTURE OF THROMBIN COMPLEXED WITH AN  
GUANIDINE-MIMETIC INHIBITOR  
Authors : Bone, R.; Lu, T.; Illig, C.R.; Soll, R.M.; Spurlino, J.C.  
Deposited on : 1999-04-27  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30686


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	122078	6075 (1.80-1.80)
Ramachandran outliers	120005	6009 (1.80-1.80)
Sidechain outliers	119972	6008 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain	
1	L	36		72% 28%
2	H	259		77% 17% . .
3	E	11		82% 9% 9%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	PPX	H	907	-	-	X	-

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2479 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called THROMBIN (LIGHT CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	L	26	214	134	35	44	1	9	0	0


- Molecule 2 is a protein called THROMBIN (HEAVY CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	249	2012	1285	354	359	14	18	0	0

- Molecule 3 is a protein called Hirudin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	E	11	104	64	12	27	1	13	0	0

- Molecule 4 is [PHENYLALANINYL-PROLINYL]-[2-(PYRIDIN-4-YLAMINO)-ETHYL]-AMINE (three-letter code: PPX) (formula: C<sub>21</sub>H<sub>27</sub>N<sub>5</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	H	1	28	21	5	2	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	L	12	Total	O	0	0
			12	12		
5	H	109	Total	O	0	0
			109	109		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.


- Molecule 1: THROMBIN (LIGHT CHAIN)

Chain L:  72% 28%





- Molecule 2: THROMBIN (HEAVY CHAIN)

Chain H:  77% 17%



- Molecule 3: Hirudin

Chain E:  82% 9% 9%



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	70.90Å 71.80Å 72.70Å 90.00° 100.52° 90.00°	Depositor
Resolution (Å)	8.00 – 1.80	Depositor
% Data completeness (in resolution range)	88.1 (8.00-1.80)	Depositor
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	X-PLOR 98.0	Depositor
R, $R_{free}$	0.193 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2479	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PPX, TYS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	L	0.56	0/216	0.62	0/287
2	H	0.51	0/2064	0.60	0/2789
3	E	0.44	0/88	0.41	0/115
All	All	0.51	0/2368	0.60	0/3191

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	H	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	H	165	ARG	Sidechain
2	H	225	TYR	Sidechain

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	214	0	214	0	0
2	H	2012	0	1983	31	0
3	E	104	0	82	1	0
4	H	28	0	27	10	0
5	H	109	0	0	2	0
5	L	12	0	0	0	0
All	All	2479	0	2306	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:H:907:PPX:H6	4:H:907:PPX:C3	1.90	1.01
4:H:907:PPX:H5	4:H:907:PPX:C3	1.96	0.95
4:H:907:PPX:C5	4:H:907:PPX:C6	2.38	0.94
4:H:907:PPX:C6	4:H:907:PPX:C3	2.50	0.89
4:H:907:PPX:C5	4:H:907:PPX:C3	2.53	0.86
4:H:907:PPX:H32	4:H:907:PPX:H5	1.73	0.69
2:H:50:ARG:NH2	2:H:107:LYS:HD3	2.10	0.66
2:H:35:ARG:HB2	2:H:41:LEU:HD13	1.77	0.66
2:H:35:ARG:O	2:H:38:GLN:HA	2.00	0.61
4:H:907:PPX:H25	5:H:458:HOH:O	2.00	0.60
2:H:17:VAL:O	2:H:188:GLY:HA2	2.04	0.57
2:H:51:TRP:CZ2	2:H:107:LYS:HE3	2.40	0.57
4:H:907:PPX:H6	4:H:907:PPX:C2	2.38	0.54
2:H:165:ARG:NH1	2:H:177:THR:O	2.41	0.54
2:H:32:MET:HG3	2:H:40:LEU:CD1	2.39	0.52
2:H:85:LEU:HD13	2:H:106:MET:HE2	1.90	0.52
2:H:143:ASN:ND2	2:H:192:GLU:HB3	2.25	0.52
2:H:235:LYS:HA	2:H:238:ILE:HD12	1.93	0.50
3:E:60:PRO:HB2	3:E:62:GLU:HG3	1.93	0.50
2:H:204(B):ASN:C	2:H:204(B):ASN:HD22	2.15	0.49
2:H:195:SER:OG	4:H:907:PPX:H202	2.14	0.47
2:H:50:ARG:HH22	2:H:107:LYS:HD3	1.80	0.47
4:H:907:PPX:H6	4:H:907:PPX:H2	1.98	0.46
2:H:51:TRP:CH2	2:H:107:LYS:HE3	2.51	0.46
2:H:203:SER:HB3	2:H:204(B):ASN:ND2	2.31	0.45
2:H:32:MET:HG3	2:H:40:LEU:HD13	1.98	0.45
2:H:165:ARG:HB3	2:H:166:PRO:HD3	2.00	0.44
2:H:134:TYR:O	2:H:162:ILE:HG12	2.17	0.43

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:203:SER:HB3	2:H:204(B):ASN:HD21	1.83	0.43
2:H:211:GLY:HA2	2:H:229:THR:O	2.19	0.43
2:H:41:LEU:HD12	2:H:41:LEU:HA	1.86	0.43
2:H:185:LYS:HB2	2:H:186(A):ASP:OD1	2.18	0.42
2:H:143:ASN:HD22	2:H:192:GLU:HB3	1.83	0.42
2:H:204(B):ASN:ND2	2:H:206:ARG:H	2.17	0.42
2:H:35:ARG:HB2	2:H:41:LEU:CD1	2.49	0.42
2:H:144:LEU:HD21	2:H:152:PRO:HB3	2.02	0.41
2:H:151:GLN:HG2	5:H:316:HOH:O	2.21	0.41
2:H:191:CYS:SG	2:H:220:CYS:CB	3.06	0.41
2:H:81:LYS:HA	2:H:81:LYS:HD3	1.92	0.41
2:H:36(A):SER:HA	2:H:37:PRO:C	2.41	0.40
2:H:201:MET:SD	2:H:210:MET:HG3	2.61	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	24/36 (67%)	23 (96%)	1 (4%)	0	100	100
2	H	245/259 (95%)	236 (96%)	8 (3%)	1 (0%)	36	22
3	E	8/11 (73%)	8 (100%)	0	0	100	100
All	All	277/306 (90%)	267 (96%)	9 (3%)	1 (0%)	36	22

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	195	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	24/31 (77%)	24 (100%)	0	100	100
2	H	217/225 (96%)	203 (94%)	14 (6%)	19	6
3	E	10/10 (100%)	9 (90%)	1 (10%)	8	2
All	All	251/266 (94%)	236 (94%)	15 (6%)	21	8

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	H	41	LEU
2	H	50	ARG
2	H	64	LEU
2	H	65	LEU
2	H	107	LYS
2	H	109	LYS
2	H	126	ARG
2	H	127	GLU
2	H	129(B)	SER
2	H	145	LYS
2	H	165	ARG
2	H	182	CYS
2	H	204(B)	ASN
2	H	236	LYS
3	E	62	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
2	H	78	ASN
2	H	204(B)	ASN
2	H	244	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	TYS	E	63	3	15,16,17	2.05	3 (20%)	19,22,24	0.73	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TYS	E	63	3	-	0/9/11/13	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	63	TYS	OH-CZ	-3.22	1.37	1.42
3	E	63	TYS	O2-S	3.38	1.58	1.45
3	E	63	TYS	O1-S	5.92	1.68	1.45

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	PPX	H	907	-	30,30,30	24.51	3 (10%)	39,39,39	17.88	7 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PPX	H	907	-	-	0/23/33/33	1/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	H	907	PPX	C6-C4	47.73	2.39	1.38
4	H	907	PPX	C5-C4	48.45	2.41	1.38
4	H	907	PPX	C3-C4	115.66	4.29	1.51

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	907	PPX	C8-C6-C4	-58.85	29.82	120.63
4	H	907	PPX	C7-C5-C4	-58.74	30.00	120.63
4	H	907	PPX	C3-C4-C6	-45.71	29.41	120.91
4	H	907	PPX	C3-C4-C5	-45.19	30.46	120.91
4	H	907	PPX	C6-C4-C5	-37.28	59.28	118.17
4	H	907	PPX	C4-C3-C2	-3.56	107.01	114.29

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	907	PPX	C22-C21-N27	-2.26	116.34	120.97

There are no chirality outliers.

There are no torsion outliers.

All (1) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	H	907	PPX	C4-C5-C6-C7-C8-C9

1 monomer is involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	H	907	PPX	10	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.