



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 14, 2020 – 03:55 pm BST

PDB ID : 2QB7
Title : Saccharomyces cerevisiae cytosolic exopolyphosphatase, phosphate complex
Authors : White, S.A.; Ugochukwu, E.
Deposited on : 2007-06-16
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	397	
1	B	397	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	ACT	A	903	-	-	X	-
4	ACT	B	901	-	-	X	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 7133 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Exopolyphosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	393	3168	2000	544	612	12	0	9	0
1	B	389	3119	1973	529	605	12	0	4	0

- Molecule 2 is COBALT (II) ION (three-letter code: CO) (formula: Co).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Co	0	0
			2	2		
2	A	2	Total	Co	0	0
			2	2		

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O P 5 4 1	0	0
3	A	1	Total O P 5 4 1	0	0
3	A	1	Total O P 5 4 1	0	0
3	B	1	Total O P 5 4 1	0	0
3	B	1	Total O P 5 4 1	0	0
3	B	1	Total O P 5 4 1	0	0

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	457	Total O 457 457	0	0
6	B	327	Total O 327 327	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.


- Molecule 1: Exopolyphosphatase

Chain A:  93% 6%



- Molecule 1: Exopolyphosphatase

Chain B:  91% 6%



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	79.99Å 83.04Å 118.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	67.42 – 1.60	Depositor
% Data completeness (in resolution range)	94.4 (67.42-1.60)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.171 , 0.201	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	7133	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, CO, EDO, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.44	0/3257	0.64	2/4380 (0.0%)
1	B	0.39	0/3184	0.56	0/4284
All	All	0.41	0/6441	0.60	2/8664 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	250	ARG	NE-CZ-NH1	-6.25	117.17	120.30
1	A	250	ARG	NE-CZ-NH2	5.12	122.86	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	64	GLU	Peptide
1	B	230	ASN	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3168	0	3193	10	0
1	B	3119	0	3133	26	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	15	0	0	0	0
3	B	15	0	0	0	0
4	A	8	0	6	3	0
4	B	4	0	3	4	0
5	A	4	0	6	0	0
5	B	12	0	18	3	0
6	A	457	0	0	2	0
6	B	327	0	0	8	0
All	All	7133	0	6359	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:62:SER:HB2	4:B:901:ACT:H1	1.76	0.67
1:B:382:LYS:NZ	6:B:1221:HOH:O	2.30	0.63
1:B:237:LEU:O	1:B:241[B]:SER:OG	2.19	0.60
1:B:382:LYS:HE3	6:B:1221:HOH:O	2.02	0.57
1:B:339:LEU:HD13	1:B:369:VAL:HG13	1.90	0.54
1:B:382:LYS:CE	6:B:1221:HOH:O	2.56	0.54
1:A:244:TYR:OH	4:A:903:ACT:CH3	2.57	0.53
1:B:63:GLU:H	4:B:901:ACT:CH3	2.21	0.52
1:B:343:ASN:HD22	1:B:346:ARG:HH22	1.58	0.52
1:B:182:GLN:NE2	6:B:1203:HOH:O	2.42	0.51
1:B:382:LYS:HE3	6:B:1163:HOH:O	2.11	0.51
1:A:53:GLN:HE21	1:A:57:ASN:HD21	1.59	0.51
1:A:34:GLY:HA2	1:A:126:VAL:O	2.11	0.50
1:B:174:PHE:CD1	1:B:195:LEU:HD22	2.46	0.50
1:B:162:ILE:HG23	1:B:164:LYS:HE3	1.94	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:34:GLY:HA2	1:B:126:VAL:O	2.13	0.48
1:B:62:SER:HB2	4:B:901:ACT:CH3	2.42	0.48
1:A:114:ASN:ND2	6:A:1250:HOH:O	2.47	0.48
1:B:115:VAL:HG21	1:B:119:THR:HG21	1.97	0.47
1:B:339:LEU:HD12	1:B:369:VAL:HG22	1.95	0.47
1:A:327:ARG:CZ	1:A:332[B]:SER:OG	2.62	0.47
1:A:244:TYR:OH	4:A:903:ACT:H2	2.15	0.47
1:A:244:TYR:OH	4:A:903:ACT:H1	2.17	0.45
1:B:148:HIS:CE1	5:B:907:EDO:H21	2.51	0.45
1:B:166:SER:O	5:B:907:EDO:H22	2.15	0.44
1:A:359:GLN:NE2	6:A:1286:HOH:O	2.50	0.44
1:A:175:ASN:HD22	1:A:220:ARG:HH21	1.66	0.43
1:B:115:VAL:CG2	1:B:119:THR:HG21	2.48	0.43
1:B:63:GLU:N	4:B:901:ACT:CH3	2.81	0.43
1:B:170[B]:SER:HB3	1:B:195:LEU:HD12	2.00	0.43
1:B:128:ASN:HD22	1:B:128:ASN:N	2.16	0.42
5:B:907:EDO:C1	6:B:1110:HOH:O	2.68	0.42
1:A:174:PHE:CD1	1:A:195:LEU:HD22	2.55	0.42
1:B:53:GLN:NE2	6:B:921:HOH:O	2.51	0.41
1:B:339:LEU:CD1	1:B:369:VAL:HG13	2.51	0.41
1:B:382:LYS:NZ	6:B:1223:HOH:O	2.52	0.41
1:B:149:HIS:CD2	1:B:382:LYS:NZ	2.90	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	400/397 (101%)	391 (98%)	8 (2%)	1 (0%)	41	21
1	B	389/397 (98%)	384 (99%)	4 (1%)	1 (0%)	41	21
All	All	789/794 (99%)	775 (98%)	12 (2%)	2 (0%)	41	21

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	230	ASN
1	A	68	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	360/355 (101%)	353 (98%)	7 (2%)	57	34
1	B	352/355 (99%)	343 (97%)	9 (3%)	46	21
All	All	712/710 (100%)	696 (98%)	16 (2%)	53	27

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	61	TYR
1	A	66	LYS
1	A	67	LYS
1	A	148	HIS
1	A	170[A]	SER
1	A	170[B]	SER
1	A	397	GLU
1	B	4	LEU
1	B	18	LEU
1	B	115	VAL
1	B	128	ASN
1	B	134	ASN
1	B	148	HIS
1	B	230	ASN
1	B	339	LEU
1	B	366	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	57	ASN
1	A	114	ASN
1	A	141	ASN
1	A	175	ASN
1	A	230	ASN
1	A	252	ASN
1	B	53	GLN
1	B	57	ASN
1	B	117	GLN
1	B	134	ASN
1	B	175	ASN
1	B	182	GLN
1	B	252	ASN
1	B	343	ASN
1	B	359	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 4 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	EDO	B	906	-	3,3,3	0.44	0	2,2,2	0.06	0
3	PO4	A	502	-	4,4,4	1.07	0	6,6,6	0.54	0
5	EDO	A	904	-	3,3,3	0.42	0	2,2,2	0.17	0
3	PO4	B	503	-	4,4,4	0.85	0	6,6,6	0.53	0
3	PO4	A	503	2	4,4,4	0.63	0	6,6,6	0.93	0
4	ACT	B	901	-	1,3,3	0.10	0	0,3,3	0.00	-
5	EDO	B	907	-	3,3,3	0.26	0	2,2,2	0.40	0
3	PO4	B	502	2	4,4,4	0.72	0	6,6,6	0.86	0
4	ACT	A	902	-	1,3,3	0.90	0	0,3,3	0.00	-
5	EDO	B	905	-	3,3,3	0.48	0	2,2,2	0.07	0
3	PO4	A	501	2	4,4,4	0.84	0	6,6,6	0.93	0
3	PO4	B	501	2	4,4,4	0.77	0	6,6,6	1.07	0
4	ACT	A	903	-	1,3,3	0.09	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	B	906	-	-	0/1/1/1	-
5	EDO	B	907	-	-	1/1/1/1	-
5	EDO	A	904	-	-	1/1/1/1	-
5	EDO	B	905	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	904	EDO	O1-C1-C2-O2
5	B	907	EDO	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	901	ACT	4	0
5	B	907	EDO	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	903	ACT	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.