



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2018 – 08:07 pm GMT

PDB ID : 3QFD
Title : Human Class I MHC HLA-A2 in complex with Mart-1(27-35) nonameric peptide
Authors : Borbulevych, O.Y.; Baker, B.M.
Deposited on : 2011-01-21
Resolution : 1.68 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtrriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.68 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	5816 (1.70-1.66)
Clashscore	122126	6305 (1.70-1.66)
Ramachandran outliers	120053	6198 (1.70-1.66)
Sidechain outliers	120020	6197 (1.70-1.66)
RSRZ outliers	108989	5700 (1.70-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	 3% 93% 6%
1	D	275	 0% 94% 5%
2	B	100	 4% 93% 7%
2	E	100	 2% 96%
3	C	9	 11% 100%
3	F	9	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	278	-	-	-	X

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 7271 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class I histocompatibility antigen, A-2 alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	275	Total	C	N	O	S	0	15	0
			2309	1445	413	442	9			
1	D	275	Total	C	N	O	S	0	8	0
			2280	1427	410	434	9			

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	100	Total	C	N	O	S	0	6	0
			858	547	143	163	5			
2	E	100	Total	C	N	O	S	0	4	0
			853	545	141	162	5			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	INITIATING METHIONINE	UNP P61769
E	0	MET	-	INITIATING METHIONINE	UNP P61769

- Molecule 3 is a protein called Mart-1(27-35) peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	9	Total	C	N	O	0	0	0
			57	37	9	11			
3	F	9	Total	C	N	O	0	1	0
			60	39	9	12			

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	B	1	Total C O 6 3 3	0	0
4	D	1	Total C O 6 3 3	0	0
4	D	1	Total C O 6 3 3	0	0

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total Na 1 1	0	0
5	E	1	Total Na 1 1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	264	Total O 264 264	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	119	Total O 120 120	0	1
6	C	8	Total O 8 8	0	0
6	D	287	Total O 287 287	0	0
6	E	125	Total O 125 125	0	0
6	F	12	Total O 12 12	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: HLA class I histocompatibility antigen, A-2 alpha chain

Chain A: 





- Molecule 1: HLA class I histocompatibility antigen, A-2 alpha chain

Chain D: 




- Molecule 2: Beta-2-microglobulin

Chain B: 




- Molecule 2: Beta-2-microglobulin

Chain E: 



- Molecule 3: Mart-1(27-35) peptide

Chain C: 



- Molecule 3: Mart-1(27-35) peptide

Chain F:  100%

There are no outlier residues recorded for this chain.

4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	58.25Å 84.12Å 83.96Å 90.00° 90.05° 90.00°	Depositor
Resolution (Å)	20.00 – 1.68 19.78 – 1.68	Depositor EDS
% Data completeness (in resolution range)	98.5 (20.00-1.68) 98.6 (19.78-1.68)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.03 (at 1.68Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.170 , 0.211 0.182 , 0.219	Depositor DCC
R_{free} test set	4543 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	15.7	Xtriage
Anisotropy	0.118	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 40.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.017 for -h,-l,-k 0.010 for -h,l,k 0.128 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7271	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.74	1/2414 (0.0%)	0.85	6/3275 (0.2%)
1	D	0.70	0/2372	0.85	2/3220 (0.1%)
2	B	0.74	0/905	0.80	0/1222
2	E	0.73	0/890	0.80	0/1202
3	C	0.82	0/56	0.73	0/74
3	F	0.82	0/63	0.76	0/84
All	All	0.73	1/6700 (0.0%)	0.83	8/9077 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	95	VAL	CB-CG2	-5.75	1.40	1.52

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	97	ARG	NE-CZ-NH1	-6.91	116.85	120.30
1	A	97	ARG	NE-CZ-NH2	6.67	123.64	120.30
1	A	110	LEU	CA-CB-CG	6.00	129.10	115.30
1	A	156	LEU	CB-CG-CD1	5.92	121.07	111.00
1	D	270	LEU	CB-CG-CD1	-5.68	101.34	111.00
1	A	37	ASP	CB-CG-OD1	5.63	123.37	118.30
1	A	215	LEU	CB-CG-CD1	5.08	119.64	111.00
1	D	156	LEU	CB-CG-CD1	5.06	119.60	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2309	0	2174	6	0
1	D	2280	0	2139	7	0
2	B	858	0	814	4	0
2	E	853	0	822	2	0
3	C	57	0	67	0	0
3	F	60	0	70	0	0
4	A	18	0	24	2	0
4	B	6	0	8	0	0
4	D	12	0	16	1	0
5	B	1	0	0	0	0
5	E	1	0	0	0	0
6	A	264	0	0	3	0
6	B	120	0	0	1	0
6	C	8	0	0	0	0
6	D	287	0	0	0	0
6	E	125	0	0	0	0
6	F	12	0	0	0	0
All	All	7271	0	6134	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:157:ARG:NH1	6:A:336:HOH:O	2.26	0.69
2:B:9[B]:VAL:HG11	2:B:80:CYS:HB3	1.87	0.56
1:A:108:ARG:NH1	1:D:174:ASN:OD1	2.43	0.52
2:B:34[A]:ASP:OD2	6:B:803:HOH:O	2.20	0.50
1:D:117:ALA:HB2	2:E:60:TRP:CE2	2.50	0.46
2:E:6:LYS:HD3	2:E:6:LYS:N	2.29	0.46
1:D:11:SER:HB2	1:D:95[A]:VAL:CG2	2.46	0.45
4:A:278:GOL:O3	6:A:281:HOH:O	2.21	0.45
1:D:6:ARG:HD2	1:D:98:MET:SD	2.56	0.45
1:A:126:LEU:HG	1:A:130:LEU:HA	2.00	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:11:SER:HB2	1:D:95[A]:VAL:HG23	2.01	0.43
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.55	0.42
2:B:9[B]:VAL:CG1	2:B:80:CYS:HB3	2.50	0.41
1:D:30:ASP:OD1	4:D:276:GOL:H31	2.21	0.41
1:A:155:GLN:NE2	6:A:613:HOH:O	2.53	0.40
1:A:6:ARG:HD2	4:A:277:GOL:O3	2.22	0.40
1:D:152:VAL:HA	1:D:155:GLN:HE21	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	288/275 (105%)	283 (98%)	5 (2%)	0	100	100
1	D	281/275 (102%)	278 (99%)	3 (1%)	0	100	100
2	B	103/100 (103%)	102 (99%)	1 (1%)	0	100	100
2	E	101/100 (101%)	100 (99%)	1 (1%)	0	100	100
3	C	7/9 (78%)	7 (100%)	0	0	100	100
3	F	8/9 (89%)	8 (100%)	0	0	100	100
All	All	788/768 (103%)	778 (99%)	10 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	244/231 (106%)	232 (95%)	12 (5%)	27	8
1	D	239/231 (104%)	232 (97%)	7 (3%)	45	22
2	B	101/95 (106%)	97 (96%)	4 (4%)	34	12
2	E	99/95 (104%)	95 (96%)	4 (4%)	34	12
3	C	5/5 (100%)	5 (100%)	0	100	100
3	F	6/5 (120%)	6 (100%)	0	100	100
All	All	694/662 (105%)	667 (96%)	27 (4%)	42	13

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	35	ARG
1	A	89[A]	GLU
1	A	89[B]	GLU
1	A	110	LEU
1	A	111	ARG
1	A	156	LEU
1	A	182[A]	THR
1	A	182[B]	THR
1	A	215	LEU
1	A	222[A]	GLU
1	A	222[B]	GLU
1	A	225	THR
2	B	0[A]	MET
2	B	0[B]	MET
2	B	70	PHE
2	B	89	GLN
1	D	2	SER
1	D	35	ARG
1	D	115	GLN
1	D	156	LEU
1	D	215[A]	LEU
1	D	215[B]	LEU
1	D	225	THR
2	E	0[A]	MET
2	E	0[B]	MET
2	E	6	LYS
2	E	70	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	72	GLN
1	A	155	GLN
1	D	115	GLN
1	D	155	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	A	276	-	5,5,5	0.31	0	5,5,5	0.55	0
4	GOL	A	277	-	5,5,5	0.72	0	5,5,5	1.32	0
4	GOL	A	278	-	5,5,5	0.55	0	5,5,5	0.58	0
4	GOL	B	100	-	5,5,5	0.46	0	5,5,5	1.80	1 (20%)
4	GOL	D	276	-	5,5,5	0.26	0	5,5,5	0.93	0
4	GOL	D	277	-	5,5,5	0.26	0	5,5,5	0.90	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	276	-	-	0/4/4/4	0/0/0/0
4	GOL	A	277	-	-	0/4/4/4	0/0/0/0
4	GOL	A	278	-	-	0/4/4/4	0/0/0/0
4	GOL	B	100	-	-	0/4/4/4	0/0/0/0
4	GOL	D	276	-	-	0/4/4/4	0/0/0/0
4	GOL	D	277	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	100	GOL	C3-C2-C1	-3.70	97.37	111.63

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	277	GOL	1	0
4	A	278	GOL	1	0
4	D	276	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	275/275 (100%)	0.16	8 (2%) 51 54	14, 22, 36, 55	0
1	D	275/275 (100%)	0.12	4 (1%) 73 77	15, 23, 39, 45	0
2	B	100/100 (100%)	0.09	4 (4%) 38 41	14, 23, 39, 45	0
2	E	100/100 (100%)	0.10	2 (2%) 65 69	16, 23, 36, 45	0
3	C	9/9 (100%)	0.60	1 (11%) 5 5	20, 22, 35, 39	0
3	F	9/9 (100%)	0.31	0 100 100	18, 24, 32, 33	0
All	All	768/768 (100%)	0.13	19 (2%) 57 60	14, 23, 37, 55	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	196	ASP	4.1
2	B	48[A]	LYS	3.8
1	A	16	GLY	3.8
1	A	17	ARG	3.7
1	D	41	ALA	3.6
1	D	111	ARG	3.1
2	E	75	LYS	3.0
1	D	196	ASP	2.9
1	D	226	GLN	2.8
2	B	47	GLU	2.7
1	A	113	TYR	2.6
2	E	48	LYS	2.6
1	A	75	ARG	2.5
2	B	23	LEU	2.3
3	C	5	GLY	2.3
1	A	15	PRO	2.2
1	A	18	GLY	2.1
2	B	9[A]	VAL	2.1
1	A	267	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GOL	B	100	6/6	0.72	0.28	39,43,45,46	0
4	GOL	A	278	6/6	0.76	0.43	41,42,42,43	6
4	GOL	D	276	6/6	0.79	0.17	36,40,41,44	0
4	GOL	A	277	6/6	0.81	0.20	42,42,44,45	0
4	GOL	D	277	6/6	0.85	0.26	26,29,32,36	0
5	NA	E	100	1/1	0.88	0.10	33,33,33,33	1
4	GOL	A	276	6/6	0.94	0.11	26,27,28,29	0
5	NA	B	101	1/1	0.97	0.17	30,30,30,30	0

6.5 Other polymers [i](#)

There are no such residues in this entry.