



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2018 – 09:21 pm GMT

PDB ID : 3QFJ
Title : The complex between TCR A6 and human Class I MHC HLA-A2 with the modified TAX (Y5F) peptide
Authors : Borbulevych, O.Y.; Baker, B.M.
Deposited on : 2011-01-21
Resolution : 2.29 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtrriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	4477 (2.30-2.30)
Clashscore	122126	5072 (2.30-2.30)
Ramachandran outliers	120053	5022 (2.30-2.30)
Sidechain outliers	120020	5021 (2.30-2.30)
RSRZ outliers	108989	4374 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	
2	B	100	
3	C	9	
4	D	200	
5	E	245	

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6898 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class I histocompatibility antigen, A-2 alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	275	2247	1403	409	426	9	0	0	0

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	100	837	533	141	159	4	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	INITIATING METHIONINE	UNP P61769

- Molecule 3 is a protein called TAX(Y5F) peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	9	76	56	9	11	0	0	0

- Molecule 4 is a protein called A6 alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	200	1552	965	255	325	7	0	0	0

- Molecule 5 is a protein called A6 beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	E	245	1927	1209	338	372	8	0	0	0

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 6 3 3	0	0
6	A	1	Total C O 6 3 3	0	0
6	B	1	Total C O 6 3 3	0	0
6	B	1	Total C O 6 3 3	0	0
6	B	1	Total C O 6 3 3	0	0
6	C	1	Total C O 6 3 3	0	0
6	D	1	Total C O 6 3 3	0	0
6	E	1	Total C O 6 3 3	0	0
6	E	1	Total C O 6 3 3	0	0
6	E	1	Total C O 6 3 3	0	0
6	E	1	Total C O 6 3 3	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	64	Total O 64 64	0	0
7	B	31	Total O 31 31	0	0
7	C	3	Total O 3 3	0	0
7	D	34	Total O 34 34	0	0
7	E	61	Total O 61 61	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: HLA class I histocompatibility antigen, A-2 alpha chain



- Molecule 2: Beta-2-microglobulin




- Molecule 3: TAX(Y5F) peptide





There are no outlier residues recorded for this chain.

- Molecule 4: A6 alpha chain



- Molecule 5: A6 beta chain





4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	224.01Å 48.10Å 92.96Å 90.00° 91.02° 90.00°	Depositor
Resolution (Å)	20.00 – 2.29 19.90 – 2.29	Depositor EDS
% Data completeness (in resolution range)	98.8 (20.00-2.29) 98.8 (19.90-2.29)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.41 (at 2.30Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.218 , 0.271 0.218 , 0.268	Depositor DCC
R_{free} test set	2244 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	45.4	Xtrriage
Anisotropy	0.282	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 44.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.016 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6898	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.59	0/2312	0.72	1/3137 (0.0%)
2	B	0.54	0/860	0.76	1/1162 (0.1%)
3	C	0.86	0/79	0.98	0/106
4	D	0.51	1/1585 (0.1%)	0.72	2/2150 (0.1%)
5	E	0.62	1/1980 (0.1%)	0.79	0/2699
All	All	0.58	2/6816 (0.0%)	0.75	4/9254 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
4	D	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	90	CYS	CB-SG	-6.07	1.72	1.82
5	E	92	CYS	CB-SG	-5.49	1.72	1.81

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	98	MET	CG-SD-CE	-8.60	86.44	100.20
4	D	89	LEU	CA-CB-CG	6.13	129.41	115.30
2	B	85	VAL	CB-CA-C	-5.40	101.13	111.40
4	D	128	LEU	CA-CB-CG	5.32	127.53	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
4	D	131	SER	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2247	0	2096	15	0
2	B	837	0	803	5	0
3	C	76	0	79	0	0
4	D	1552	0	1461	9	0
5	E	1927	0	1830	12	0
6	A	12	0	16	2	0
6	B	18	0	24	0	0
6	C	6	0	8	0	0
6	D	6	0	8	0	0
6	E	24	0	32	0	0
7	A	64	0	0	1	0
7	B	31	0	0	0	0
7	C	3	0	0	0	0
7	D	34	0	0	0	0
7	E	61	0	0	0	0
All	All	6898	0	6357	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:D:164:THR:OG1	5:E:195:ARG:NH2	2.32	0.63
5:E:95:ARG:HG2	5:E:106:GLN:HB2	1.82	0.62
1:A:14:ARG:HA	6:A:277:GOL:H11	1.85	0.59
4:D:5:GLN:NE2	4:D:88:TYR:O	2.38	0.55
1:A:224:GLN:O	1:A:228:THR:OG1	2.24	0.54
1:A:14:ARG:HG2	6:A:277:GOL:H31	1.90	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:223:ASP:OD1	1:A:223:ASP:N	2.34	0.52
1:A:214:THR:HB	1:A:262:GLN:HG3	1.91	0.52
1:A:73:THR:HG22	5:E:98:LEU:HG	1.92	0.52
4:D:33:PHE:HB2	4:D:91:ALA:HB3	1.93	0.51
5:E:231:LYS:HG2	5:E:233:VAL:HG13	1.92	0.50
2:B:11:SER:OG	2:B:13:HIS:O	2.27	0.50
1:A:97:ARG:NH2	7:A:295:HOH:O	2.45	0.49
5:E:6:GLN:HB3	5:E:110:PRO:HD2	1.95	0.49
5:E:132:PRO:HD3	5:E:145:LEU:HG	1.97	0.47
1:A:122:ASP:OD2	2:B:60:TRP:NE1	2.42	0.47
4:D:118:ILE:HD11	4:D:145:ASP:HA	1.94	0.47
4:D:120:ASN:HA	4:D:121:PRO:HD2	1.74	0.46
4:D:63:THR:HB	4:D:76:LEU:HB2	1.98	0.46
5:E:12:VAL:HG13	5:E:154:PRO:HG3	1.97	0.46
1:A:188:HIS:HB2	1:A:204:TRP:HB2	1.98	0.46
5:E:211:ARG:NH2	5:E:213:GLN:OE1	2.51	0.44
1:A:48:ARG:HA	1:A:48:ARG:HD3	1.74	0.44
2:B:19:LYS:HE3	2:B:19:LYS:HB2	1.78	0.43
1:A:163:THR:OG1	4:D:68:LYS:NZ	2.52	0.43
2:B:25:CYS:HB2	2:B:39:LEU:HD21	2.00	0.43
1:A:215:LEU:HD12	1:A:243:LYS:HD3	2.00	0.43
1:A:5:MET:HB2	1:A:168:LEU:HD13	2.00	0.42
4:D:8:GLY:HA2	4:D:9:PRO:HA	1.65	0.42
1:A:150:ALA:O	5:E:101:GLY:HA2	2.19	0.42
5:E:59:GLU:H	5:E:59:GLU:HG3	1.56	0.42
5:E:244:ARG:NH1	5:E:246:ASP:OD2	2.49	0.42
4:D:20:LEU:HD12	4:D:75:LEU:HD23	2.02	0.41
5:E:29:HIS:ND1	5:E:94:SER:OG	2.33	0.40
1:A:23:ILE:HD12	2:B:54:LEU:HB3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	273/275 (99%)	261 (96%)	12 (4%)	0	100	100
2	B	98/100 (98%)	96 (98%)	2 (2%)	0	100	100
3	C	7/9 (78%)	7 (100%)	0	0	100	100
4	D	198/200 (99%)	186 (94%)	11 (6%)	1 (0%)	31	38
5	E	243/245 (99%)	229 (94%)	12 (5%)	2 (1%)	21	25
All	All	819/829 (99%)	779 (95%)	37 (4%)	3 (0%)	36	45

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
5	E	2	ALA
4	D	170	SER
5	E	39	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/231 (100%)	210 (91%)	21 (9%)	10	12
2	B	95/95 (100%)	88 (93%)	7 (7%)	15	19
3	C	8/8 (100%)	8 (100%)	0	100	100
4	D	178/178 (100%)	158 (89%)	20 (11%)	6	7
5	E	209/209 (100%)	182 (87%)	27 (13%)	5	5
All	All	721/721 (100%)	646 (90%)	75 (10%)	8	9

All (75) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	VAL
1	A	23	ILE
1	A	35	ARG
1	A	52	ILE
1	A	72	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	82	ARG
1	A	98	MET
1	A	113	TYR
1	A	128	GLU
1	A	134	THR
1	A	165	VAL
1	A	177	GLU
1	A	178	THR
1	A	182	THR
1	A	218	GLN
1	A	223	ASP
1	A	225	THR
1	A	255	GLN
1	A	256	ARG
1	A	262	GLN
1	A	272	LEU
2	B	4	THR
2	B	34	ASP
2	B	64	LEU
2	B	70	PHE
2	B	75	LYS
2	B	85	VAL
2	B	98	ASP
4	D	1	LYS
4	D	6	ASN
4	D	7	SER
4	D	12	VAL
4	D	22	CYS
4	D	23	THR
4	D	31	SER
4	D	73	VAL
4	D	103	LYS
4	D	110	THR
4	D	120	ASN
4	D	127	GLN
4	D	128	LEU
4	D	135	ASP
4	D	136	LYS
4	D	145	ASP
4	D	151	SER
4	D	186	ASP
4	D	191	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
4	D	197	ILE
5	E	12	VAL
5	E	13	LEU
5	E	19	MET
5	E	20	THR
5	E	25	GLN
5	E	31	TYR
5	E	56	ASP
5	E	59	GLU
5	E	67	VAL
5	E	72	THR
5	E	88	SER
5	E	94	SER
5	E	95	ARG
5	E	98	LEU
5	E	117	GLU
5	E	119	LEU
5	E	133	SER
5	E	146	VAL
5	E	163	VAL
5	E	170	SER
5	E	172	VAL
5	E	195	ARG
5	E	207	ARG
5	E	226	THR
5	E	229	ARG
5	E	244	ARG
5	E	246	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	2	GLN
4	D	105	GLN
4	D	111	GLN
4	D	127	GLN
4	D	191	ASN
5	E	11	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	GOL	A	276	-	5,5,5	0.34	0	5,5,5	0.27	0
6	GOL	A	277	-	5,5,5	0.17	0	5,5,5	0.72	0
6	GOL	B	100	-	5,5,5	0.34	0	5,5,5	0.46	0
6	GOL	B	101	-	5,5,5	0.55	0	5,5,5	0.86	0
6	GOL	B	102	-	5,5,5	0.29	0	5,5,5	0.60	0
6	GOL	C	10	-	5,5,5	0.42	0	5,5,5	0.40	0
6	GOL	D	207	-	5,5,5	0.35	0	5,5,5	0.51	0
6	GOL	E	247	-	5,5,5	0.55	0	5,5,5	0.55	0
6	GOL	E	248	-	5,5,5	0.41	0	5,5,5	0.36	0
6	GOL	E	249	-	5,5,5	0.77	0	5,5,5	0.69	0
6	GOL	E	250	-	5,5,5	0.42	0	5,5,5	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	276	-	-	0/4/4/4	0/0/0/0
6	GOL	A	277	-	-	0/4/4/4	0/0/0/0
6	GOL	B	100	-	-	0/4/4/4	0/0/0/0
6	GOL	B	101	-	-	0/4/4/4	0/0/0/0
6	GOL	B	102	-	-	0/4/4/4	0/0/0/0
6	GOL	C	10	-	-	0/4/4/4	0/0/0/0
6	GOL	D	207	-	-	0/4/4/4	0/0/0/0
6	GOL	E	247	-	-	0/4/4/4	0/0/0/0
6	GOL	E	248	-	-	0/4/4/4	0/0/0/0
6	GOL	E	249	-	-	0/4/4/4	0/0/0/0
6	GOL	E	250	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	277	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	275/275 (100%)	0.58	37 (13%) 3 4	23, 51, 155, 190	0
2	B	100/100 (100%)	0.25	6 (6%) 22 28	23, 45, 86, 98	0
3	C	9/9 (100%)	-0.25	0 100 100	30, 31, 34, 35	0
4	D	200/200 (100%)	0.73	31 (15%) 2 3	31, 58, 110, 134	0
5	E	245/245 (100%)	0.28	19 (7%) 13 17	21, 50, 86, 111	0
All	All	829/829 (100%)	0.48	93 (11%) 5 7	21, 52, 122, 190	0

All (93) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	D	171	MET	8.1
1	A	253	GLN	7.2
4	D	206	SER	7.1
1	A	223	ASP	7.0
2	B	99	MET	6.8
5	E	184	ALA	6.6
1	A	256	ARG	6.2
1	A	195	SER	6.1
4	D	186	ASP	6.0
1	A	257	TYR	5.5
1	A	221	GLY	5.2
1	A	274	TRP	4.9
1	A	247	VAL	4.7
1	A	224	GLN	4.7
1	A	201	LEU	4.7
5	E	246	ASP	4.5
1	A	275	GLU	4.4
4	D	184	LYS	4.0
1	A	217	TRP	4.0
4	D	185	SER	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	252	GLY	3.8
4	D	205	PRO	3.8
1	A	248	VAL	3.7
5	E	228	ASP	3.7
1	A	260	HIS	3.6
1	A	226	GLN	3.6
1	A	227	ASP	3.6
1	A	219	ARG	3.4
1	A	255	GLN	3.4
4	D	133	SER	3.4
1	A	220	ASP	3.4
1	A	196	ASP	3.4
4	D	169	ARG	3.4
5	E	141	GLN	3.3
1	A	273	ARG	3.2
4	D	8	GLY	3.2
1	A	194	VAL	3.1
5	E	2	ALA	3.0
1	A	197	HIS	3.0
4	D	131	SER	2.9
2	B	1	ILE	2.9
5	E	41	MET	2.9
4	D	154	LYS	2.9
5	E	222	ASN	2.9
1	A	264	GLU	2.9
5	E	1	ASN	2.8
5	E	186	ASN	2.8
4	D	187	PHE	2.8
4	D	140	LEU	2.8
4	D	156	SER	2.8
4	D	172	ASP	2.8
5	E	207	ARG	2.7
4	D	179	VAL	2.7
4	D	129	ARG	2.6
1	A	17	ARG	2.6
1	A	222	GLU	2.6
4	D	1	LYS	2.6
1	A	267	PRO	2.6
1	A	199	ALA	2.5
5	E	80	LEU	2.5
1	A	249	VAL	2.5
2	B	48	LYS	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	75	LYS	2.4
5	E	144	THR	2.4
4	D	139	CYS	2.4
1	A	218	GLN	2.4
4	D	203	PHE	2.4
1	A	250	PRO	2.4
4	D	130	ASP	2.3
5	E	166	LYS	2.3
5	E	78	ARG	2.3
1	A	258	THR	2.2
4	D	195	ASN	2.2
1	A	89	GLU	2.2
1	A	261	VAL	2.2
4	D	191	ASN	2.2
1	A	251	SER	2.2
5	E	182	GLN	2.2
4	D	135	ASP	2.2
5	E	119	LEU	2.1
5	E	177	GLN	2.1
4	D	159	TYR	2.1
4	D	197	ILE	2.1
1	A	203	CYS	2.1
2	B	80	CYS	2.1
4	D	137	SER	2.1
5	E	16	GLY	2.1
2	B	23	LEU	2.1
4	D	7	SER	2.1
4	D	204	PHE	2.0
4	D	34	TRP	2.0
5	E	185	LEU	2.0
4	D	132	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	GOL	B	100	6/6	0.77	0.24	76,76,76,76	0
6	GOL	D	207	6/6	0.79	0.37	48,48,49,50	0
6	GOL	B	102	6/6	0.81	0.22	46,47,48,49	0
6	GOL	A	277	6/6	0.81	0.38	47,48,49,50	0
6	GOL	A	276	6/6	0.83	0.26	72,72,72,73	0
6	GOL	E	250	6/6	0.83	0.44	52,53,54,55	0
6	GOL	B	101	6/6	0.85	0.20	47,47,48,48	0
6	GOL	E	249	6/6	0.89	0.45	41,43,44,44	0
6	GOL	E	248	6/6	0.94	0.17	58,59,59,59	0
6	GOL	C	10	6/6	0.94	0.29	47,47,48,49	0
6	GOL	E	247	6/6	0.95	0.14	53,53,53,54	6

6.5 Other polymers [i](#)

There are no such residues in this entry.